

Relation Algebra

§14. Thinning Algorithms

§14.1. Thinning

(14.1a)

For $Q : A \rightarrow A$, $\text{thin}(Q) \triangleq (\exists \exists) \cap (\epsilon \setminus (Q^\circ \epsilon)) : EA \rightarrow EA$ (8.1).


$ys \text{ thin}(Q) xs \iff xs \subseteq ys \wedge (\forall a \in ys. \exists b \in xs. b \ Q \ a)$ the same $^\circ$ as $\triangleq \text{est}$: ϵX runs $a \rightarrow ys \rightarrow xs$, member of ys first, and $Q^\circ \epsilon$ runs $a \rightarrow b \rightarrow xs$, so $(a, b) \in Q^\circ$ reads $b \ Q \ a$; at $Q^\circ \leq$ every $a \in ys$ keeps some $b \leq a$ in xs , the end $\text{est}(\leq)$ picks

(14.1b)


the law	what it says
$X \subseteq \frac{S}{\exists} \text{thin}(Q) \iff X \exists \subseteq S$ and $S^\circ X \subseteq Q^\circ \epsilon$	everything kept is an S-value, and every S-value has a Q-lower bound among the kept ones
$\text{thin}(Q) \exists \exists$	everything a thinning keeps was in the set
$\text{thin}(Q)^\circ \exists \exists \exists Q$	every element of the set has a Q-lower bound among the kept ones
$Q \subseteq R \implies \text{thin}(Q) \subseteq \text{thin}(R)$	the fewer pairs Q relates, the fewer subsets count as thinnings
$1 \subseteq Q \implies 1 \subseteq \text{thin}(Q)$ $Q \subseteq Q \implies \text{thin}(Q) \text{thin}(Q) \subseteq \text{thin}(Q)$	keeping everything is always a legal thinning
$Q \subseteq R \wedge 1 \subseteq Q \wedge R R \subseteq R \implies \text{est}(R) = \text{thin}(Q) \text{est}(R)$ $Q \subseteq R, 1 \subseteq Q, R R \subseteq R$ — (14.1.2a); weaker than the book's “both preorders”: Q transitive is never used and $1 \subseteq R$ follows	thin-introduction : thinning first cannot lose an R-minimum
$\text{thin}(Q) \supseteq \text{est}(Q) \frac{1}{\exists}$ (8.2) — (14.1.2d)	thin-elimination : keeping one element is a thinning, but its domain is the sets est (Q) is defined on
$\frac{S}{\exists} \text{thin}(Q) \supseteq \frac{S}{\exists} \text{est}(R) \frac{1}{\exists}$ (8.3), $R \cap (S^\circ S) \subseteq Q$ — thin-elimination variant	the usable variant: R need only refine Q between values S gives one argument
$P(\text{thin}(Q)) \text{union} \subseteq \text{union thin}(Q)$ (8.4)	thinning each member set is a thinning of the union

§14.1.1. $X \subseteq (S^\circ \exists) \text{thin}(Q) \iff X \exists \subseteq S$ and $S^\circ X \subseteq Q^\circ \epsilon$

(14.1.1a)



(14.1.1b)



(14.1.1a) at $X \triangleq \text{thin}(Q)$, $S \triangleq \exists$: the left side is $\text{thin}(Q) \subseteq \text{thin}(Q)$, since $\exists \exists = \wedge(\exists) = 1$

§14.1.2. $\text{est}(R) = \text{thin}(Q) \text{est}(R)$ given $Q \subseteq R, 1 \subseteq Q, R R \subseteq R$

$\epsilon \text{ thin}(Q) \text{ est}(R) \sqsubseteq R^\circ$ (14.1.2c)	
	the $\exists$ half, second condition — $Q \sqsubseteq R$, R transitive
$A \text{---} \boxed{\epsilon} \text{---} \boxed{\text{thin}(Q)} \text{---} \epsilon A \text{---} \boxed{\text{est}(R)} \text{---} A$	
$\sqsubseteq$ $A \text{---} \boxed{Q} \text{---} \boxed{\epsilon} \text{---} \epsilon A \text{---} \boxed{\text{est}(R)} \text{---} A$	$\epsilon \text{ thin}(Q) \sqsubseteq Q^\circ \epsilon$ — (14.1.1a) at $S \triangleq \exists, X \triangleq \text{thin}(Q)$, second half
$\sqsubseteq$ $A \text{---} \boxed{Q} \text{---} \boxed{R} \text{---} A$	$\epsilon \text{ est}(R) \sqsubseteq R^\circ$ — est-laws first row at $X \triangleq \text{est}(R)$, second half, conversed
$\sqsubseteq$ $A \text{---} \boxed{R} \text{---} \boxed{R} \text{---} A$	$Q \sqsubseteq R$, conversed
$\sqsubseteq$ $A \text{---} \boxed{R} \text{---} A$	R transitive, conversed

$\text{est}(Q) \stackrel{1}{\exists} \text{thin}(Q)$ (14.1.2d)	
the singleton holding a Q -least member of a set is a thinning of that set	
circuit the $\exists/\exists$ half of $\triangleq$ thin at $X \triangleq \text{est}(Q)$ $\frac{1}{3}$	Hinze–Marsden
$\frac{1}{3} \exists = 1$ — pow-laws — then $\text{est}(Q) \sqsubseteq \exists$ — est-laws	
the $\epsilon \setminus (Q^\circ \epsilon)$ half of $\triangleq$ thin, $\neg \sqsubseteq Q^\circ \epsilon$	
$\epsilon \text{ est}(Q) \sqsubseteq Q^\circ$ — est-up at $X \triangleq \text{est}(Q)$, conversed	
$\frac{1}{3} \sqsubseteq \epsilon$, since $\frac{1}{3} \exists = 1$ with $\frac{1}{3}$ a map — pow-laws	

$\mathbf{RnS^\circ S \sqsubseteq Q \implies \frac{S}{S} \text{ est}(R) \frac{1}{S} \sqsubseteq \frac{S}{S} \text{ thin}(Q)}$ given $\mathbf{Rn(S^\circ S) \sqsubseteq Q}$, Q a preorder	
circuit	Hinze–Marsden
$\leftarrow B \text{---} \boxed{\frac{1}{S}} \text{---} EB \text{---} \boxed{E(S)} \text{---} EA \text{---} \boxed{\text{est}(R)} \text{---} A \text{---} \boxed{\frac{1}{S}} \text{---} EA \text{---} \boxed{\exists} \text{---} A$ the first of (14.1b)‘ two conditions at $X \stackrel{S}{=} \text{est}(R) \frac{1}{S}$	
$\sqsubseteq B \text{---} \boxed{S} \text{---} A$ $\frac{1}{S} \exists = 1$ — pow-laws — then $\frac{S}{S} \text{ est}(R) = S n(S^\circ \setminus R^\circ) \sqsubseteq S$ — est-laws	
$\leftarrow A \text{---} \boxed{S} \text{---} B \text{---} \boxed{\frac{1}{S}} \text{---} EB \text{---} \boxed{E(S)} \text{---} EA \text{---} \boxed{\text{est}(R)} \text{---} A \text{---} \boxed{\frac{1}{S}} \text{---} EA$ the second condition, $\neg \sqsubseteq Q^\circ \sqsubseteq$	
$= A \text{---} \boxed{S} \text{---} B \text{---} \boxed{\frac{1}{S}} \text{---} EB \text{---} \boxed{E(S)} \text{---} EA \text{---} \boxed{\text{est}(RnS^\circ S)} \text{---} A \text{---} \boxed{\frac{1}{S}} \text{---} EA$ $\frac{S}{S} \text{ est}(R) = \frac{S}{S} \text{ est}(RnS^\circ S)$ — est-laws	
$\sqsubseteq A \text{---} \boxed{\epsilon} \text{---} EA \text{---} \boxed{\text{est}(RnS^\circ S)} \text{---} A \text{---} \boxed{\frac{1}{S}} \text{---} EA$ $S^\circ \frac{S}{S} \sqsubseteq \epsilon$, since $\frac{S}{S} \exists = S$ with $\frac{S}{S}$ a map — pow-laws	
$\sqsubseteq A \text{---} \boxed{Q} \text{---} \boxed{\frac{1}{S}} \text{---} EA$ $\epsilon \text{ est}(RnS^\circ S) \sqsubseteq (RnS^\circ S)^\circ$ — est-up at $X \stackrel{A}{=} \text{est}(RnS^\circ S)$, conversed — then $\mathbf{Rn(S^\circ S) \sqsubseteq Q}$	
$\sqsubseteq A \text{---} \boxed{Q} \text{---} \boxed{\epsilon} \text{---} EA$ $\frac{1}{S} \sqsubseteq \epsilon$, since $\frac{1}{S} \exists = 1$ with $\frac{1}{S}$ a map — pow-laws	

$\text{PreservesRecip}(F) \wedge Q \sqsubseteq Q \wedge \text{MonotonicAlg}(S, Q) \implies (\frac{F(\exists)S}{\exists} \text{thin}(Q)) \sqsubseteq \frac{(S)}{\exists} \text{thin}(Q)$ (14.1.2f) thinning at every step of the reduce is a thinning of the whole candidate set — S monotonic on Q , Q a preorder	
circuit	Hinze–Marsden
$(\frac{F(\exists)S}{\exists} \text{thin}(Q)) \sqsubseteq (S)$ and $(S) \circ (\frac{F(\exists)S}{\exists} \text{thin}(Q)) \sqsubseteq Q^\circ \epsilon$ (14.1b) at $X \triangleq (\frac{F(\exists)S}{\exists} \text{thin}(Q))$, (S) for its S	
$\leftarrow A \text{---} \boxed{S} \text{---} FA \text{---} \boxed{F(Q^\circ \epsilon)} \text{---} FEA \text{---} \boxed{\frac{1}{\exists}} \text{---} EFEA \text{---} \boxed{E(F(\exists)S)} \text{---} \boxed{\text{thin}(Q)} \text{---} EA$ the first by fusion; hylo-least at the bound $Q^\circ \epsilon$ reduces the second to $\sqsubseteq Q^\circ \epsilon$	
$\sqsubseteq A \text{---} \boxed{Q} \text{---} \boxed{S} \text{---} FA \text{---} \boxed{F(\epsilon)} \text{---} FEA \text{---} \boxed{\frac{1}{\exists}} \text{---} EFEA \text{---} \boxed{E(F(\exists)S)} \text{---} \boxed{\text{thin}(Q)} \text{---} EA$ $S^\circ F(Q^\circ) \sqsubseteq Q^\circ S^\circ$ — mon-str at S, conversed; $F(R)^\circ = F(R^\circ)$ — relator-laws	
$\sqsubseteq A \text{---} \boxed{Q} \text{---} \boxed{\epsilon} \text{---} \boxed{\text{thin}(Q)} \text{---} EA$ $S^\circ F(\epsilon) \xrightarrow{\frac{F(\exists)S}{\exists}} \epsilon$, since $\frac{F(\exists)S}{\exists} \exists = F(\exists)S$ with $\frac{F(\exists)S}{\exists}$ a map — pow-laws	
$\sqsubseteq A \text{---} \boxed{Q} \text{---} \boxed{Q} \text{---} \boxed{\epsilon} \text{---} EA$ $\epsilon \text{thin}(Q) \sqsubseteq Q^\circ \epsilon$, the $\epsilon \setminus (Q^\circ \epsilon)$ half of $\triangleq \text{thin}$ — adj-all	
$\sqsubseteq A \text{---} \boxed{Q} \text{---} \boxed{\epsilon} \text{---} EA$ $Q^\circ Q^\circ = Q^\circ$, Q a preorder	

$\text{PreservesRecip}(F) \wedge Q \in R \wedge 1 \in Q \wedge Q \circ Q \in Q \wedge R \circ R \in R \wedge \text{MonotonicAlg}(S, Q) \Rightarrow \langle \frac{F(\exists)S}{\exists} \text{thin}(Q) \rangle \text{est}(R) \in \langle \frac{QS}{\exists} \text{est}(R) \rangle$ (14.1.2g) the thinning fold refines the optimisation problem itself — S monotonic on Q , $Q \in R$, both preorders	
circuit	Hinze–Marsden
$\sqsubseteq$ thinning theorem	
$=$ $\text{est}(R) = \text{thin}(Q) \text{ est}(R) \quad (14.1b), Q \in R$	

§14.2. Paths in a layered network

$F(A, X) = A + A \times X$, $L = \text{list}^+$ with initial algebra $\alpha \triangleq [\text{wrap}, \text{cons}] : F(A, LA) \rightarrow LA$. $\text{wrapz} \triangleq (\text{wrap}, \text{zero})$, $\text{consw}(a, (xs, n)) = (\text{cons}(a, xs), \text{wt}(a, \text{head}(xs)) + n)$. $\text{cost} \triangleq \langle [\text{wrapz}, \text{consw}] \rangle \pi_2$, $\langle [\text{wrapz}, \text{consw}] \rangle = \langle 1, \text{cost} \rangle$, $R \triangleq \text{cost} \leq \text{cost}^\circ$. $Q \triangleq R \cap (\text{head} \text{ head}^\circ)$, $S \triangleq F(1, \exists) \alpha$, $\frac{F(\exists, 1)}{\exists} = 1 + \text{cpl}$, $\frac{F(1, \exists)}{\exists} = 1 + \text{cpr}$, $\text{step} \triangleq \text{cpr } P(\text{cons}) \text{ est}(R)$.	(14.2a)
---	---------

the law	what it says
$F(\exists, Q) \alpha \in F(\exists, 1) \alpha Q$	$F(\exists, 1) \alpha$ is monotonic on Q ; on R it is not, since the next edge can cost arbitrarily much
$S \text{ head} \in [1, \pi_1] \quad S \triangleq F(1, \exists) \alpha$	$S \text{ head}$ is simple, which gives $R \cap (S^\circ S) \in Q$: between two paths S builds from one argument, equal cost and equal head already means Q

$\text{PreservesRecip}(F) \wedge Q \in R \wedge 1 \in Q \wedge Q \circ Q \in Q \wedge R^\circ R^\circ \in R^\circ \wedge \text{MonotonicAlg}(F(\exists, 1) \alpha, Q) \wedge R \cap (F(1, \exists) \alpha)^\circ F(1, \exists) \alpha \in Q \Rightarrow$ (14.2c) $\langle \frac{F(\exists, 1)}{\exists} P(\frac{F(1, \exists) \alpha}{\exists} \text{est}(R)) \rangle \text{est}(R) \in \langle \frac{F(\exists, 1) \alpha}{\exists} \text{est}(R) \rangle$ a least-cost path in a layered network, as a fold over the layers	
circuit	Hinze–Marsden
$T(EA) \rightarrow \left[\frac{1}{\exists} \right] \rightarrow E(T(EA)) \rightarrow E(\langle F(\exists, 1) \alpha \rangle) \rightarrow EB \rightarrow \text{est}(R) \rightarrow B$ $= \frac{L(\exists)}{\exists} \text{est}(R)$	
$\sqsubseteq$ (14.1.2g), at $F(\exists, 1) \alpha$ monotonic on Q — (14.2b).	
$\sqsubseteq$ (14.2d) under $\langle \rangle$ monotonic: the whole chain runs inside the reduce, and the $\text{est}(R)$ behind it never moves.	

$\text{Rn}(F(1, \exists)\alpha) \circ F(1, \exists)\alpha \sqsubseteq Q \implies \frac{F(\exists, 1)}{\exists} P(\frac{F(1, \exists)\alpha}{\exists} \text{est}(R)) \sqsubseteq \frac{F(\exists, \exists)\alpha}{\exists} \text{thin}(Q)$ (14.2d)	
thinning the algebra of a layered network costs no more than taking the program's two cases	
circuit	Hinze–Marsden
$F(EA)(EB) \xrightarrow{\frac{1}{\exists}} E(F(EA)(EB)) \xrightarrow{E(F(\exists, \exists)\alpha)} \text{thin}(Q) \xrightarrow{} EB$ the algebra of (14.2c) row 2	
$C \xrightarrow{\frac{1}{\exists}} EC \xrightarrow{E(V)} EW \xrightarrow{P(\frac{S}{\exists})} EEA \xrightarrow{\text{union}} EA$ $= F(\exists, \exists) = F(\exists, 1)F(1, \exists); \frac{F(\exists, 1)F(1, \exists)\alpha}{\exists} = \frac{F(\exists, 1)}{\exists} P(\frac{F(1, \exists)\alpha}{\exists}) \text{union}.$	
$\sqsubseteq EEA \xrightarrow{P(\text{thin}(Q))} \text{union} \xrightarrow{} EA$ union thin(Q) $\sqsupseteq$ P(thin(Q)) union — (14.1b).	
$\sqsubseteq B \xrightarrow{\frac{1}{\exists}} EB \xrightarrow{E(S)} EA \xrightarrow{\text{est}(R)} A \xrightarrow{\frac{1}{\exists}} EA$ $\frac{S}{\exists} \text{thin}(Q) \sqsupseteq \frac{S}{\exists} \text{est}(R) \xrightarrow{\frac{1}{\exists}}$ — (14.1b) at $S \triangleq F(1, \exists)\alpha$, $\text{Rn}(S \circ S) \sqsubseteq Q$ — (14.2b).	
$= F(EA)(EB) \xrightarrow{\frac{1}{\exists}} E(F(EA)(EB)) \xrightarrow{E(F(\exists, 1))} E(FA)(EB) \xrightarrow{P(\frac{F(1, \exists)\alpha}{\exists} \text{est}(R))} EB$ $P(\frac{1}{\exists}) \text{union} = 1.$	
$= V + V \times \text{pow}(\text{list}^+(V)) \xrightarrow{} \text{list}^+(V)$ $\begin{array}{c} V \xrightarrow{\text{wrap}} \\ \text{pow}(\text{list}^+(V)) \xrightarrow{\text{step}(wt)} \end{array}$ α a map, so $\frac{F(1, \exists)\alpha}{\exists} = \frac{F(1, \exists)}{\exists} P(\alpha); \frac{F(1, \exists)}{\exists} P(\alpha) \text{est}(R) = [\text{wrap}, \text{step}] \triangleq \text{cost}.$ $= \frac{F(\exists, 1)}{\exists} = 1 + \text{cpl} \triangleq \text{cost}$	

§14.3. Implementing thin

(14.3a)

$\text{setify} : [A] \rightarrow EA$, $\text{cup} : EA \times EA \rightarrow EA$, $\text{cp}(F) \triangleq \frac{F(\exists)}{\exists}$, $\text{listcp} : F(L) \rightarrow LF$, $\text{sort}(P) \triangleq \text{setify}^\circ \text{ ordered } P$ for P a connected preorder.

$\text{thinlist}(Q)$ is any $\text{thinlist}(Q) \sqsubseteq \text{subseq}$ with $\text{thinlist}(Q) \text{ setify} \sqsubseteq \text{setify thin}(Q)$; one is $([\text{nil}, \text{bump } Q])$, $\text{bump } Q (a, []) = [a]$, $\text{bump } Q (a, [b] \# xs) = (b \sqsubseteq a \rightarrow [a] \# xs, a \sqsubseteq b \rightarrow [b] \# xs, [a] \# [b] \# xs)$.

Binary thinning data: $S = (f_1 p_1) \cup (f_2 p_2)$ with p_1, p_2 coreflexive; Q a preorder with $Q \sqsubseteq R$ and both $f_1 p_1, f_2 p_2$ monotonic on Q ; P a connected preorder with both f_1, f_2 monotonic on P ; $g_i \triangleq \text{list}(f_i) \text{ filter}(p_i)$.

(14.3b)

the law	what it says
$\text{thinlist}(Q) \text{ xs} = [\text{minlist}(Q) \text{ xs}]$ (8.5), Q connected, xs non-empty	what thinning should come to when it can: one element
$\text{sort}(P) \text{ thinlist}(Q) \sqsubseteq \text{thin}(Q) \text{ sort}(P)$ (8.6) — (14.3c)	thinning a sorted list is a thinning of the set — this is what $\text{thinlist}(Q) \sqsubseteq \text{subseq}$ buys
$\text{sort}(P) \text{ minlist}(Q) \sqsubseteq \text{est}(Q)$ (8.7)	a minimum of the sorted list is a minimum of the set
$\text{sort}(f P f^\circ) \text{ list}(f) \sqsubseteq P(f) \text{ sort}(P)$ (8.8)	shunt a function through a sort
$\text{sort}(P) \text{ filter}(p) \sqsubseteq E(p) \text{ sort}(P)$ (8.9), p coreflexive	filtering a sorted list sorts the restricted set
$(\text{sort}(P) \times \text{sort}(P)) \text{ merge}(P) \sqsubseteq \text{cup sort}(P)$ (8.10)	merging two sorted lists sorts their union
$F(\text{sort}(P)) \text{ listcp} \sqsubseteq \text{cp}(F) \text{ sort}(FP)$ (8.11), F linear	listcp is the list implementation of the cartesian product $\text{cp}(F)$

Map(setify) $\wedge$ Coreflexive(ordered) $\wedge$ thinlist $\sqsubseteq$ subseq $\wedge$ ordered subseq $\sqsubseteq$ subseq ordered $\wedge$ thinlist (14.3c)
setify $\sqsubseteq$ setify thin(Q) $\Rightarrow$ sortRel(L, setify, ordered) thinlist $\sqsubseteq$ thin(Q) sortRel(L, setify, ordered)
 a thinning of the sorted list lists a thinning of the set — P a connected preorder, $\text{thinlist}(Q) \sqsubseteq \text{subseq}$.

circuit	Hinze–Marsden
$EA \text{---} \boxed{\text{setify}} \text{---} \boxed{\text{ordered}} \text{---} \boxed{\text{thinlist}} \text{---} LA$	
$\begin{aligned} &EA \text{---} \boxed{\text{setify}} \text{---} \boxed{\text{ordered}} \text{---} \boxed{\text{thinlist}} \text{---} LA \\ &= \text{sort}(P) \triangleq \text{setify}^\circ \text{ ordered } P \text{---} \triangleq \text{thinlist} \end{aligned}$	
$EA \text{---} \boxed{\text{setify}} \text{---} \boxed{\text{thinlist}} \text{---} \boxed{\text{ordered}} \text{---} LA$ $\sqsubseteq \text{ordered } P \text{ thinlist}(Q) \sqsubseteq \text{thinlist}(Q) \text{ ordered } P$, since $\text{thinlist}(Q) \sqsubseteq \text{subseq}$ — $\triangleq \text{thinlist}$ — and a subsequence of a P -ordered list is P -ordered	
$EA \text{---} \boxed{\text{thin}(Q)} \text{---} \boxed{\text{setify}} \text{---} \boxed{\text{ordered}} \text{---} LA$ $\sqsubseteq \triangleq \text{thinlist's thinlist}(Q) \text{ setify} \sqsubseteq \text{setify thin}(Q)$ after setify° , at $\text{setify}^\circ \text{ setify} \sqsubseteq 1$ for setify simple — dom-laws — then $\cdot \text{setify} \cdot \text{setify}^\circ$ — triple-chains	
$\begin{aligned} &EA \text{---} \boxed{\text{thin}(Q)} \text{---} \boxed{\text{setify}} \text{---} \boxed{\text{ordered}} \text{---} LA \\ &= \text{sort}(P) \triangleq \text{setify}^\circ \text{ ordered } P \text{---} \triangleq \text{thinlist} \end{aligned}$	

$\text{Map}(f) \wedge (\forall X. \forall Y. X \sqsubseteq Y \Rightarrow \text{F}(X) \sqsubseteq \text{sortF}(Y)) \wedge \text{MonotonicAlg}(f, P) \wedge \text{sortF}(fP f^\circ) \text{listf} \sqsubseteq P(f) \text{sort}(P) \wedge \text{sort}(P) \text{filterp} \sqsubseteq E(p) \text{sort}(P) \wedge \text{F}(\text{sort}(P)) \text{listcp} \sqsubseteq \text{cp}(F, A) \text{sortF}(F(P))$ $\Rightarrow \text{F}(\text{sort}(P)) \text{listcp} \text{listf} \text{filterp} \sqsubseteq \frac{\text{F}(\exists)fp}{\exists} \text{sort}(P)$ one sorted list built from sorted arguments, instead of a set built and then sorted — $f : FA \rightarrow A$ monotonic on P , p coreflexive, F linear.	
circuit	Hinze–Marsden
$\text{FEA} \rightarrow \boxed{\frac{1}{\exists}} \rightarrow \text{EFA} \rightarrow \boxed{E(F(\exists)fp)} \rightarrow \text{EA} \rightarrow \boxed{\text{sort}(P)} \rightarrow \text{LA}$	
$= \text{FEA} \rightarrow \boxed{\text{cp}(F, A)} \rightarrow \text{EFA} \rightarrow \boxed{E(fp)} \rightarrow \text{EA} \rightarrow \boxed{\text{sort}(P)} \rightarrow \text{LA}$ $= \frac{\text{F}(\exists)fp}{\exists} = \frac{\text{F}(\exists)}{\exists} E(fp) \text{ — pow-laws; } \text{cp}(F) \triangleq \frac{\text{F}(\exists)}{\exists} \text{ — } \triangleq \text{thinlist}$	
$= \text{FEA} \rightarrow \boxed{\text{cp}(F, A)} \rightarrow \text{EFA} \rightarrow \boxed{P(f)} \rightarrow \boxed{E(p)} \rightarrow \text{EA} \rightarrow \boxed{\text{sort}(P)} \rightarrow \text{LA}$ $= E(fp) = E(f)E(p); E(f) = P(f) \text{ for } f \text{ a map — powrel-laws}$	
$\sqsubseteq \text{FEA} \rightarrow \boxed{\text{cp}(F, A)} \rightarrow \text{EFA} \rightarrow \boxed{P(f)} \rightarrow \text{EA} \rightarrow \boxed{\text{sort}(P)} \rightarrow \boxed{\text{filterp}} \rightarrow \text{LA}$ $\text{sort}(P) \text{ filter}(p) \sqsubseteq E(p) \text{ sort}(P) \text{ — (14.3b)}$	
$\sqsubseteq \text{FEA} \rightarrow \boxed{\text{cp}(F, A)} \rightarrow \text{EFA} \rightarrow \boxed{\text{sortF}(fP f^\circ)} \rightarrow \text{LFA} \rightarrow \boxed{\text{listf}} \rightarrow \boxed{\text{filterp}} \rightarrow \text{LA}$ $\text{sort}(fP f^\circ) \text{ list}(f) \sqsubseteq P(f) \text{ sort}(P) \text{ — (14.3b)}$	
$\sqsubseteq \text{FEA} \rightarrow \boxed{\text{cp}(F, A)} \rightarrow \text{EFA} \rightarrow \boxed{\text{sortF}(F(P))} \rightarrow \text{LFA} \rightarrow \boxed{\text{listf}} \rightarrow \boxed{\text{filterp}} \rightarrow \text{LA}$ $FP \sqsubseteq fP f^\circ \text{ — mon-str at } f \text{ a map; } \text{sort}(P) \triangleq \text{setify}^\circ \text{ ordered } P \text{ grows with } P \text{ — } \triangleq \text{thinlist}$	
$\sqsubseteq \text{FEA} \rightarrow \boxed{F(\text{sort}(P))} \rightarrow \text{FLA} \rightarrow \boxed{\text{listcp}} \rightarrow \text{LFA} \rightarrow \boxed{\text{listf}} \rightarrow \boxed{\text{filterp}} \rightarrow \text{LA}$ $F(\text{sort}(P)) \text{ listcp} \sqsubseteq \text{cp}(F) \text{ sort}(FP) \text{ — (14.3b), } F \text{ linear}$	

$$\text{PreservesRecip}(F) \wedge \text{Map}(f_1) \wedge \text{Map}(f_2) \wedge Q \sqsubseteq R \wedge 1 \sqsubseteq Q \wedge Q \sqsubseteq Q \wedge R^\circ R^\circ \sqsubseteq R^\circ \wedge \text{MonotonicAlg}(f_1, p_1, Q) \wedge \text{MonotonicAlg}(f_2, p_2, Q) \wedge (\forall X. \forall Y. X \sqsubseteq Y \Rightarrow \text{sort}F(X) \sqsubseteq \text{sort}F(Y)) \wedge \text{MonotonicAlg}(f_1, P) \wedge \text{MonotonicAlg}(f_2, P) \wedge \text{sort}F(f_1$$

$$P f_1^\circ) \text{list}f_1 \sqsubseteq P(f_1) \text{sort}(P) \wedge \text{sort}F(f_2$$

$$P f_1^\circ) \text{list}f_1 \sqsubseteq P(f_1) \text{sort}(P) \wedge \text{sort}F(f_2$$

$$(f_2) \text{sort}(P) \wedge \text{sort}(P) \text{filter}p_1 \sqsubseteq E(p_1) \text{sort}(P) \wedge \text{sort}(P) \text{filter}p_2 \sqsubseteq E(p_2) \text{sort}(P) \wedge F(\text{sort}(P)) \text{list}cp \sqsubseteq cp(F, A) \text{sort}F(F(P)) \wedge (\text{sort}(P) \times \text{sort}(P))$$

$$\text{sort}(P) \wedge \text{sort}(P) \text{thinlist}(Q) \sqsubseteq \text{thin}(Q) \text{sort}(P) \wedge \text{sort}(P) \text{minlist}(R) \sqsubseteq \text{est}(R) \wedge g_1 = \text{list}f_1$$









$$\text{filter}p_1 \wedge g_2 = \text{list}f_2 \quad \text{filter}p_2 \wedge S = f_1 \quad p_1 \cup f_2 \quad p_2 \Rightarrow (\text{list}cp(g_1, g_2) \text{merge}(P) \text{thinlist}(Q))$$

$$\text{minlist}(R) \sqsubseteq \frac{(S)}{S} \text{est}(R)$$

a fold on sorted lists of partial solutions, thinned at every step — at $\triangleq$ **thinlist**'s binary thinning data.

circuit



Hinze–Marsden




$\begin{aligned} & \text{Map}(f_1) \wedge \text{Map}(f_2) \wedge (\forall X. \forall Y. X \sqsubseteq Y \implies \\ & \text{sortF}(X) \sqsubseteq \text{sortF}(Y)) \wedge \text{MonotonicAlg}(f_1, P) \wedge \text{MonotonicAlg}(f_2, P) \wedge \text{sortF}(f_1 \\ & P f_1^\circ) \text{listf}_1 \sqsubseteq P(f_1) \text{sort}(P) \wedge \text{sortF}(f_2 \\ & (f_2) \text{sort}(P) \wedge \text{sort}(P) \text{filterp}_1 \sqsubseteq E(p_1) \text{sort}(P) \wedge \text{sort}(P) \text{filterp}_2 \sqsubseteq E(p_2) \text{sort}(P) \wedge F(\text{sort}(P)) \text{listcp} \sqsubseteq \text{cp}(F, A) \text{sortF}(F(P)) \wedge (\text{sort}(P) \times \text{sort}(P)) \\ & \text{sort}(P) \wedge \text{sort}(P) \text{thinlist}(Q) \sqsubseteq \text{thin}(Q) \text{sort}(P) \implies F(\text{sort}(P)) \text{listcp}(\text{listf}_1 \text{filterp}_1, \text{listf}_2 \\ & \text{filterp}_2) \text{merge}(P) \text{thinlist}(Q) \sqsubseteq \frac{F(\exists)(f_1 p_1 \cup f_2 p_2)}{\exists} \text{thin}(Q) \text{sort}(P) \end{aligned} \quad (14.3f)$ sorting the candidate set is what turns the thinning algebra into an algebra on lists — the side condition of binary thinning theorem's last step.	
---	--

circuit

Hinze–Marsden



$$\begin{aligned} & \text{FEA} \rightarrow \boxed{\frac{1}{3}} \rightarrow \text{EFEA} \rightarrow \boxed{E(F(\exists)S)} \rightarrow \text{EA} \rightarrow \text{sort}(P) \rightarrow \text{thinlist}(Q) \rightarrow \text{LA} \\ & \text{sort}(P) \text{thinlist}(Q) \sqsubseteq \text{thin}(Q) \text{sort}(P) \quad (14.3b) \end{aligned}$$



$$\begin{aligned} & \text{FEA} \rightarrow \boxed{\frac{1}{3}} \rightarrow \text{EFEA} \rightarrow \boxed{E(F(\exists)f_1 p_1)} \rightarrow \text{EA} \rightarrow \text{cup} \rightarrow \text{EA} \rightarrow \text{sort}(P) \rightarrow \text{thinlist}(Q) \rightarrow \text{LA} \\ & = \left\langle \frac{F(\exists)f_1 p_1}{\exists}, \frac{F(\exists)f_2 p_2}{\exists} \right\rangle \text{cup sort}(P) \text{thinlist}(Q) \\ & S = (f_1 p_1) \cup (f_2 p_2) \triangleq \text{thinlist}, \text{ then } \triangleq \text{cup} \end{aligned}$$

$$\begin{aligned} & \text{FEA} \rightarrow \boxed{\frac{1}{3}} \rightarrow \text{EFEA} \rightarrow \boxed{E(F(\exists)f_1 p_1)} \rightarrow \text{EA} \rightarrow \text{sort}(P) \rightarrow \text{merge}(P) \rightarrow \text{thinlist}(Q) \rightarrow \text{LA} \\ & = \left\langle \frac{F(\exists)f_1 p_1}{\exists}, \frac{F(\exists)f_2 p_2}{\exists} \right\rangle (\text{sort}(P) \times \text{sort}(P)) \text{merge}(P) \text{thinlist}(Q) \\ & (\text{sort}(P) \times \text{sort}(P)) \text{merge}(P) \sqsubseteq \text{cup sort}(P) \quad (14.3b) \end{aligned}$$

$$\begin{aligned} & \text{FEA} \rightarrow \boxed{\frac{1}{3}} \rightarrow \text{EFEA} \rightarrow \boxed{E(F(\exists)f_1 p_1)} \rightarrow \text{EA} \rightarrow \text{sort}(P) \rightarrow \text{merge}(P) \rightarrow \text{thinlist}(Q) \rightarrow \text{LA} \\ & = \left\langle \frac{F(\exists)f_1 p_1}{\exists} \text{sort}(P), \frac{F(\exists)f_2 p_2}{\exists} \text{sort}(P) \right\rangle \text{merge}(P) \text{thinlist}(Q) \\ & (X, Y) (\text{sort}(P) \times \text{sort}(P)) = (X \text{sort}(P), Y \text{sort}(P)) \quad \text{bdm-prod-laws} \end{aligned}$$

$$\begin{aligned} & \text{FEA} \rightarrow \boxed{F(\text{sort}(P))} \rightarrow \text{FLA} \rightarrow \text{listcp} \rightarrow \text{LFA} \rightarrow \left\langle \text{listf}_1, \text{listf}_2 \right\rangle \rightarrow \left\langle \text{filterp}_1, \text{filterp}_2 \right\rangle \rightarrow \text{merge}(P) \rightarrow \text{thinlist}(Q) \rightarrow \text{LA} \\ & \text{F}(\text{sort}(P)) \text{listcp} \langle g_1, g_2 \rangle \text{merge}(P) \text{thinlist}(Q) \\ & (14.3d) \text{ at } f_1, p_1 \text{ and at } f_2, p_2, \text{ then } X \langle g_1, g_2 \rangle \sqsubseteq Xg_1, Xg_2 \quad \text{bdm-prod-laws; } g_i \triangleq \text{list}(f_i) \\ & \text{filter}(p_i) \triangleq \text{thinlist} \end{aligned}$$

§14.4. The knapsack problem

(14.4a)

```

vol, wt : Item → Real, value ≐ list(vol) sum, weight ≐ list(wt) sum .

subseq = ([nil, cons] ∪ [nil, π₂]) ∩ , within w the coreflexive on xs with weight xs ≤ w, 0 ≤ w.

R ≐ value ≥ value°, Q ≐ R ∩ (weight ≤ weight°), P ≐ R, .

FA = 1 + Item × A, listcp = wrap + cpr, g₁ ≐ list([nil, cons]) filter(within w) = [list(nil), h₁],
g₂ ≐ list([nil, π₂]) = [list(nil), h₂].

h₁ ≐ list(cons) filter(within w), h₂ ≐ list(π₂).

```

(14.4b)

the law	what it says
$(1 \times R) (\text{cons } (\text{within } w)) \sqsubseteq \text{cons } (\text{within } w) R$ FALSE	not monotonic on R: a selection of greater value need not still fit once one more item goes in
$(1 \times Q) (\text{cons } (\text{within } w)) \sqsubseteq \text{cons } (\text{within } w) Q$ $(1 \times Q) \pi_2 \sqsubseteq \pi_2 Q$,	both halves are monotonic on Q once ties in value are broken by weight

$$0 \leq w \wedge (\forall i. 0 \leq \text{wt}(i)) \wedge (\forall X. \forall Y. X \sqsubseteq Y \Rightarrow \quad (14.4c)$$

$$F(X) \sqsubseteq \text{sort} F(Y) \wedge \text{sort} F([\text{nil}, \text{cons}] R [\text{nil}, \text{cons}]^\circ) \text{list} f_1 \sqsubseteq P([\text{nil}, \text{cons}]) \text{sort} P \wedge \text{sort} F([\text{nil}, \pi_2] R [\text{nil}, \pi_2]^\circ) \text{list} f_2 \sqsubseteq P([\text{nil}, \pi_2]) \text{sort} P \wedge \text{sort}$$

$$\text{filter} p_1 \sqsubseteq E(\text{within}(w)) \text{sort} P \wedge F(\text{sort} P) \text{list} \text{cp} \sqsubseteq \text{cp}(F, [\text{Item}]) \text{sort} F(F(R)) \wedge (\text{sort} P \times \text{sort} P) \text{merge} P \sqsubseteq \text{cup} \\ \text{sort} P \wedge \text{sort} P \text{ thinlist} \sqsubseteq \text{thin}(Q(\text{vol}, \text{wt})) \text{sort} P \wedge \text{sort} P \text{ minlist} \sqsubseteq \text{est}(R) \Rightarrow \langle \text{list} \text{cp}(\text{list} f_1 \\ \text{filter} p_1, \text{list} f_2) \text{merge} P \text{ thinlist} \rangle \text{minlist} \sqsubseteq \frac{\text{subseq } \text{within}(w)}{\supset} \text{est}(R)$$

the knapsack problem, as a fold that thins the packings kept at each item

circuit	Hinze–Marsden
$[\text{Item}] \xrightarrow{\frac{1}{\supset}} E(\text{subseq } \text{within}(w)) E[\text{Item}] \xrightarrow{\text{est}(R)} [\text{Item}]$ $\frac{\text{subseq } (\text{within } w)}{\supset} \text{est}(R)$	
$[\text{Item}] \xrightarrow{\frac{1}{\supset}} E(\langle [\text{nil}, \text{cons}] \text{within}(w) \cup [\text{nil}, \pi_2] \rangle) E[\text{Item}] \xrightarrow{\text{est}(R)} [\text{Item}]$ $= \frac{\langle [\text{nil}, \text{cons}] (\text{within } w) \cup [\text{nil}, \pi_2] \rangle}{\supset} \text{est}(R)$ fusion, weights non-negative.	
$\sqsubseteq \langle \text{list} \text{cp} \langle g_1, g_2 \rangle \text{merge } R \text{ thinlist}(Q) \rangle \text{minlist}(R)$ binary thinning theorem, at $P \triangle R$, F linear, Q from (14.4b)	
$\langle [\text{nil}, \text{cpr} \langle h_1, h_2 \rangle \text{merge } R \text{ thinlist}(Q) \rangle \rangle \text{minlist}(R)$ $= \text{list} \text{cp} = \text{wrap} + \text{cpr}, g_i = [\text{list}(\text{nil}), h_i] \text{ — } \triangle \text{ within};$ minlist(R) is head, packings coming out in descending value	

§14.5. The paragraph problem

(14.5a)

```

Line=list+ Word, Para=list+ Line, FA=Word+Word×A, listcp=wrap+cpr.

new(a,xs)=[[a]]#xs, glue(a,xs)=[[a]#head(xs)]#tail(xs), partition≐([wrap wrap,new u glue]) : list+
Word→Para.

width≐([length,(length×1) plus succ]), 0≤length a, fits w the coreflexive on a line x with width x≤w, ok w the
coreflexive on [x]#xs with width x≤w.

white w x=w-width x, collect≐list(sqr) sum, waste w≐init list(white w) collect.

R≐(waste w)≤(waste w)°, Q≐Rn(head head°), P≐T, .

g1≐list([wrap wrap,new]), g2≐list([wrap wrap,glue]) filter(ok w), start≐wrap wrap wrap, h1≐list(new),
h2≐list(glue) filter(ok w).

```

(14.5b)

the law	what it says
(1×R) glue⊆glue R FALSE	glue is not monotonic on R: the waste of a paragraph depends on its whole first line, so no greedy algorithm solves this
(1×Q) new⊆new Q (1×Q) (glue (ok w))⊆glue (ok w)Q cons monotonic on collect≤collect°,	both halves are monotonic on Q once ties in waste are broken by the first line
merge T=cat; P≐head prefix head° also serves	T needs no sorting at all, and prefix is a linear order on first lines of paragraphs of one input

$$\begin{aligned}
 & (\forall a. 0 \leq \text{len}(a)) \wedge (\forall a. \text{len}(a) \leq w) \wedge (\forall X. \forall Y. X \sqsubseteq Y \Rightarrow \text{sortF}(X) \sqsubseteq \text{sortF}(Y)) \wedge \text{sortF}([\text{wrap} \\
 & \text{wrap}, \text{new}] \tau [\text{wrap} \text{ wrap}, \text{new}]^\circ) \text{listf}_1 \sqsubseteq P([\text{wrap} \text{ wrap}, \text{new}]) \text{sortP} \sqsubseteq \text{sortF}([\text{wrap} \text{ wrap}, \text{glue}] \tau [\text{wrap} \\
 & \text{wrap}, \text{glue}]^\circ) \text{listf}_2 \sqsubseteq P([\text{wrap} \text{ wrap}, \text{glue}]) \text{sortP} \sqsubseteq \text{sortP} \\
 & \text{filterp}_2 \sqsubseteq E(\text{ok}(w)) \text{sortP} \sqsubseteq F(\text{sortP}) \text{listcp} \sqsubseteq \text{cp}(F, \text{list}^+((\text{list}^+ \\
 & \text{Word}))) \text{sortF}(F(\tau)) \wedge (\text{sortP} \times \text{sortP}) \text{mergeP} \sqsubseteq \text{cup} \text{ sortP} \sqsubseteq \text{sortP} \text{ thinlist} \sqsubseteq \text{thin}(Q(\text{len}, w)) \text{sortP} \sqsubseteq \text{sortP} \\
 & \text{minlist} \sqsubseteq \text{est}(R) \Rightarrow (\text{listcp}(\text{listf}_1, \text{listf}_2 \text{ filterp}_2) \text{mergeP} \text{ thinlist}) \text{minlist} \sqsubseteq \frac{\text{partition fits}(w)}{3} \\
 & \text{est}(R)
 \end{aligned}
 \tag{14.5c}$$

a paragraph laid out as a fold that thins the layouts kept at each word

circuit	Hinze–Marsden
$ \begin{aligned} & \text{list}^+(\text{Word}) \xrightarrow{\frac{1}{3}} E(\text{list}^+(\text{Word})) \xrightarrow{E(\text{partition fits}(w))} E(\text{list}^+(\text{list}^+(\text{Word}))) \xrightarrow{\text{est}(R)} \text{list}^+(\text{list}^+(\text{Word})) \\ & \frac{\text{partition list}^+(\text{fits } w)}{3} \text{est}(R) \end{aligned} $	
$ \begin{aligned} & \text{list}^+(\text{Word}) \xrightarrow{\frac{1}{3}} E(\text{list}^+(\text{Word})) \xrightarrow{E([\text{wrap wrap}, \text{new}] \cup [\text{wrap wrap}, \text{glue}] \text{ok}(w))} E(\text{list}^+(\text{list}^+(\text{Word}))) \xrightarrow{\text{est}(R)} \text{list}^+(\text{list}^+(\text{Word})) \\ & = \frac{([\text{wrap wrap}, \text{new}] \cup [\text{wrap wrap}, \text{glue}] \text{ok}(w))}{3} \text{est}(R) \end{aligned} $ fusion, every word fits on a line by itself.	
$ \begin{aligned} & \text{list}^+(\text{Word}) \xrightarrow{\frac{1}{3}} E(\text{list}^+(\text{Word})) \xrightarrow{E([\text{wrap wrap}, \text{new}] \cup [\text{wrap wrap}, \text{glue}] \text{ok}(w))} E(\text{list}^+(\text{list}^+(\text{Word}))) \xrightarrow{\text{est}(R)} \text{list}^+(\text{list}^+(\text{Word})) \\ & = \frac{([\text{wrap wrap}, \text{new}] \cup ([\text{wrap wrap}, \text{glue}] \text{ok } w))}{3} \text{est}(R) \end{aligned} $ the algebra as $(f_1 p_1) \cup (f_2 p_2)$, $p_1 \triangleq 1 \text{ — } \triangleq \text{thinlist}$.	
$ \begin{aligned} & \text{listcp} \langle g_1, g_2 \rangle \text{cat thinlist}(Q) \text{minlist}(R) \\ & \text{binary thinning theorem, at } P \triangleq T \text{ with merge } \tau = \text{cat}, Q \text{ from} \\ & (14.5b) \end{aligned} $	
$ \begin{aligned} & ([\text{start}, \text{cpr} \langle h_1, h_2 \rangle \text{cat thinlist}(Q)]) \\ & = \text{minlist}(R) \\ & \text{listcp} = \text{wrap} + \text{cpr}, g_i \text{ along the coproduct — } \triangleq \text{partition} \end{aligned} $	

§14.6. Bitonic tours

$\text{FA} = (\text{City} \times \text{City}) + (\text{City} \times A)$, the base functor of cons-lists of length at least two; $\text{listcp} = \text{wrap} + \text{cpr}$; $\text{tc} : \text{City} \times \text{City} \rightarrow \text{Real}$, neither positive nor symmetric.

$\text{start } (a, b) = ([a, b], [a, b])$, $\text{dropl } (a, ([b] \# xs, ys)) = ([a] \# xs, [a] \# ys)$, $\text{dropr } (a, (xs, [b] \# ys)) = ([a] \# xs, [a] \# ys)$, $\text{tour} \triangleq ([\text{start}, \text{dropl} \cup \text{dropr}])$.

$\text{cost } (xs, ys) = \text{outcost } xs + \text{incost } ys$, $\text{outcost } [a_0, \dots, a_n] = \text{tc } (a_0, a_1) + \dots + \text{tc } (a_{n-1}, a_n)$, $\text{incost } [a_0, \dots, a_n] = \text{tc } (a_1, a_0) + \dots + \text{tc } (a_n, a_{n-1})$.

$\text{next} \triangleq \text{tail head}$, $\text{next2} \triangleq \text{next} \times \text{next}$, $\text{head2} \triangleq \text{head} \times \text{head}$, $R \triangleq \text{cost} \leq \text{cost}^\circ$, $Q \triangleq R \cap (\text{next2 next2}^\circ) \cap (\text{head2 head2}^\circ)$, $P \triangleq T$, $g_1 \triangleq \text{list}([\text{start}, \text{dropl}])$, $g_2 \triangleq \text{list}([\text{start}, \text{dropr}])$.

(14.6b)

the law	what it says
$(1 \times R) \text{ dropl} \sqsubseteq \text{dropl } R$ FALSE	neither drop is monotonic on R: the two edges it adds and removes depend on head and next of both lists
$(1 \times R) \text{ dropr} \sqsubseteq \text{dropr } R$ FALSE	
$(1 \times Q) \text{ dropl} \sqsubseteq \text{dropl } Q$	both are, once ties in cost are broken by the two second cities — the heads already agree among tours of one input
$(1 \times Q) \text{ dropr} \sqsubseteq \text{dropr } Q$,	

$$(\forall X. \forall Y. X \sqsubseteq Y \Rightarrow$$

(14.6c)

$$, \text{dropLFn}]^\circ) g_1 \sqsubseteq P([\text{start}, \text{dropLFn}]) \text{sortP} \wedge \text{sortF}([\text{start}, \text{dropRFn}] \tau [\text{start}, \text{dropRFn}]^\circ) g_2 \sqsubseteq P([\text{start}, \text{dropRFn}]) \text{sortP} \wedge F(\text{sortP}) \text{listcp} \sqsubseteq \text{cp}(F$$

$$\text{sortP} \wedge \text{sortP} \text{ thinlist} \sqsubseteq \text{thin}(Qc(tc)) \text{sortP} \wedge \text{sortP} \text{ minlist} \sqsubseteq \text{est}(R) \Rightarrow (\text{listcp}(g_1, g_2) \text{cat thinlist})$$

$$\text{minlist} \sqsubseteq \frac{(\text{tourAlg})}{\exists} \text{est}(R)$$


a least-cost bitonic tour, as a fold that thins the tours kept at each city

circuit

Hinze–Marsden

$$\text{dCL}(\text{City} \times \text{City}, \text{City}) \xrightarrow{\frac{1}{3}} E(\text{dCL}(\text{City} \times \text{City}, \text{City})) \xrightarrow{E((\text{tourAlg}))} E(\text{Journey}(\text{City})^2) \xrightarrow{\text{est}(R)} \text{Journey}(\text{City})$$

$$\frac{\text{tour}}{\exists} \text{est}(R)$$




$$\text{dCL}(\text{City} \times \text{City}, \text{City}) \xrightarrow{\frac{1}{3}} E(\text{dCL}(\text{City} \times \text{City}, \text{City})) \xrightarrow{E([\text{start}, \text{dropLFn}] \cup [\text{start}, \text{dropRFn}])} E(\text{Journey}(\text{City})^2) \xrightarrow{\text{est}(R)} \text{Journey}(\text{City})$$

$$= \frac{([\text{start}, \text{dropL}] \cup [\text{start}, \text{dropR}])}{\exists} \text{est}(R)$$

$$\text{tour} \triangleq ([\text{start}, \text{dropL}] \cup [\text{start}, \text{dropR}]) \triangleq \text{tour}$$

$$\text{dCL}(\text{City} \times \text{City}, \text{City}) \xrightarrow{\left[\frac{([\text{start}, \text{dropL}] \cup [\text{start}, \text{dropR}])}{\exists} \right]} \text{listcp} \cdot \text{list}(F(\text{Journey}(\text{City})^2)) \xrightarrow{\left[\frac{g_1}{\exists} \right]} \text{list}(\text{Journey}(\text{City})^2) \xrightarrow{\left[\frac{g_2}{\exists} \right]} \text{list}(\text{Journey}(\text{City})^2) \xrightarrow{\text{cat}} \text{thinlist} \xrightarrow{\text{list}(\text{Journey}(\text{City})^2)} \text{minlist} \xrightarrow{\text{Journey}(\text{City})} \text{Journey}(\text{City})$$

$$\sqsubseteq (\text{listcp}(g_1, g_2) \text{cat thinlist}(Q)) \text{minlist}(R)$$

 binary thinning theorem, at $P \triangleq T$ with merge $\tau = \text{cat}$, Q from (14.6b)


$$([\text{start wrap}, \text{cpr}(\text{list}(\text{dropL}), \text{list}(\text{dropR})) \text{cat thinlist}(Q)] \text{minlist}(R)$$

$$\text{listcp} = \text{wrap} + \text{cpr}, g_i = [\text{list}(\text{start}), \text{list}(\text{drop}_i)] \triangleq \text{tour};$$

quadratic, two tours added per step