

Companion to Bird & de Moor, Algebra of Programming

§10. Greedy Algorithms

§10.1. Theory


(10.1a)

h, T, R, H, M as in $\triangleq H$; additionally Q a **connected** preorder on the sets $\frac{T^\circ}{\exists}$ returns, so that $\frac{T^\circ}{\exists} \text{ est}(Q)$ is entire.

**Map(h) $\wedge$ MonoAlg(h,R) $\wedge$ R R $\subseteq$ R $\wedge$ ThinCondition(T,h,R,Q) $\Rightarrow$ ($\mu X : \frac{T^\circ}{\exists} \text{ est}(Q) F(X)$ Theorem 10.1
 $h) \subseteq \frac{H}{\exists} \text{ est}(R)$**


the same optimum reached by keeping ONE decomposition at each step, so that no set is ever carried
 and the recursion runs on values alone

$H \triangleq (T)^\circ (h) \triangleq H, Q$



(a)	$\frac{H}{\exists} \text{ est}(R)$	
(b)	$\sqsubseteq \frac{T^\circ}{\exists} \text{ est}(Q) F(X) h$	


the body at T a coproduct of two arms



(a)	$\frac{T^\circ}{\exists} \text{ est}(Q) F(X) U$	Proposition 10.1, $V_2 V_1^\circ = \emptyset$
(b)	$\sqsubseteq \frac{\text{arm}_2^\circ}{\exists} \text{ est}(\text{arm}_2) (X \times 1) \text{ arm}_2$	

$T^\circ F(H) h = H \Rightarrow \frac{T^\circ}{\exists} \text{ est}(Q) F(\frac{H}{\exists} \text{ est}(R)) h \subseteq H$ (10.1b)


taking the input apart in one Q -extreme way, solving the parts by M and assembling by h returns only what H
 returns




(a)	$\frac{T^\circ}{\exists} \text{ est}(Q) F(\frac{H}{\exists} \text{ est}(R)) h$	$\text{est}(Q) \sqsubseteq \exists \triangleq \text{est}$
(b)	$\sqsubseteq \frac{T^\circ}{\exists} \exists F(\frac{H}{\exists} \text{ est}(R)) h$	$\frac{T^\circ}{\exists} \exists = T^\circ$ — pow-laws
(c)	$= T^\circ F(\frac{H}{\exists} \text{ est}(R)) h$	$M \sqsubseteq \frac{H}{\exists} \exists = H$ — est-up
(d)	$\sqsubseteq T^\circ F(H) h$	$T^\circ F(H) h = H$ — hylomorphism theorem
(e)	$= H$	

$$\text{Map}(h) \wedge \text{MonoAlg}(h, R^\circ) \wedge R^\circ R^\circ \subseteq R^\circ \wedge T^\circ F(H) h = H \wedge Q F(H) h \subseteq F(H) h R \implies H^\circ \xrightarrow{T^\circ} \text{est}(Q) F\left(\frac{H}{3} \text{est}(R)\right) h \subseteq R^\circ \quad (10.1c)$$

H° followed by the greedy body at M is $\subseteq R^\circ$: an answer of the body is never worse than an answer of H to the same input




(a)	$H^\circ \xrightarrow{T^\circ} \text{est}(Q) F\left(\frac{H}{3} \text{est}(R)\right) h$	$H^\circ = h^\circ F(H^\circ) T$ — hyломorphism theorem
(b)	$= h^\circ F(H^\circ) T \xrightarrow{T^\circ} \text{est}(Q) F\left(\frac{H}{3} \text{est}(R)\right) h$	$T \xrightarrow{T^\circ} \subseteq \epsilon$
(c)	$\subseteq h^\circ F(H^\circ) \epsilon \text{est}(Q) F\left(\frac{H}{3} \text{est}(R)\right) h$	$\epsilon \text{est}(Q) \subseteq Q^\circ$ — est-up
(d)	$\subseteq h^\circ F(H^\circ) Q^\circ F\left(\frac{H}{3} \text{est}(R)\right) h$	$QF(H) h \subseteq F(H) h R$ conversed



(a)	$\subseteq R^\circ h^\circ F(H^\circ) F\left(\frac{H}{3} \text{est}(R)\right) h$	$H^\circ M \subseteq R^\circ$ under F — est-up
(b)	$\subseteq R^\circ h^\circ F(R^\circ) h$	$h^\circ F(R^\circ) h \subseteq R^\circ$
(c)	$\subseteq R^\circ R^\circ$	R transitive
(d)	$\subseteq R^\circ$	

$$V_2 \ V_1^\circ = \emptyset \implies \frac{[V_1, V_2]^\circ}{3} \text{est}(Q_1 + Q_2) [U_1, U_2] = (\text{ran}(V_1) \ \frac{V_1^\circ}{3} \text{est}(Q_1) \ U_1) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{3} \text{est}(Q_2) \ U_2) \quad \text{Proposition 10.1}$$

when V_1 and V_2 have disjoint ranges, the greedy step over $[V_1, V_2]$ runs the V_1 step on inputs V_1 reaches and the V_2 step on inputs V_2 reaches



(a)	$\frac{[V_1, V_2]^\circ}{3} \text{est}(Q_1 + Q_2) [U_1, U_2]$	empty off $\text{ran } V_1 \cup \text{ran } V_2$
(b)	$= (\text{ran}(V_1) \ \frac{V_1^\circ}{3} \text{est}(Q_1 + Q_2) [U_1, U_2]) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{3} \text{est}(Q_1 + Q_2) [U_1, U_2])$	$\frac{[V_1, V_2]^\circ}{3} = \frac{V_1^\circ}{3} P(\text{inl})$ on $\text{ran } V_1$ — dp-disjoint
(c)	$= (\text{ran}(V_1) \ \frac{V_1^\circ}{3} P(l) \text{est}(Q_1 + Q_2) [U_1, U_2]) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{3} P(r) \text{est}(Q_1 + Q_2) [U_1, U_2])$	$P(\text{inl})\text{est}(Q_1 + Q_2) = \text{est}(Q_1)\text{inl}$
(d)	$= (\text{ran}(V_1) \ \frac{V_1^\circ}{3} \text{est}(Q_1) \ l [U_1, U_2]) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{3} \text{est}(Q_2) \ r [U_1, U_2])$	$\text{inl}[U_1, U_2] = U_1$
(e)	$= (\text{ran}(V_1) \ \frac{V_1^\circ}{3} \text{est}(Q_1) \ U_1) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{3} \text{est}(Q_2) \ U_2)$	

§10.2. The detab-entab problem

(10.2a)

$\text{detab} \triangleq ([\text{nil}, \text{expand}]) : \text{String} \rightarrow \text{String}$ over snoc -lists, $\alpha \triangleq [\text{nil}, \text{snoc}]$, $H = \text{detab}^\circ$; $\text{expand } (xs, a) = (a = \text{TB} \rightarrow \text{fill } xs, xs \# [a]), \text{ fill } xs = xs \# \text{blanks } (n - (\text{col } xs) \bmod n)$.


$\text{col} \triangleq ([\text{zero}, \text{count}])$, $\text{count } (c, a) = (a = \text{NL} \rightarrow 0, c + 1)$; TB the tab, BL the blank, NL the newline, tab stops every n columns.

$R \triangleq \text{length} \leq \text{length}^\circ$, U the preorder with $a \cup b \iff a = \text{TB} \vee a = b$, $V \triangleq \text{prefix}^\circ n (\text{fill } \text{fill}^\circ)$, $Q \triangleq 1 + (V \times U)$.


$\text{unfill } xs$ the shortest prefix of xs with fill ($\text{unfill } xs = \text{fill } xs$); tbc the trailing blank count, $\text{triple} \triangleq (\text{unfill } \text{entab}, (\text{tbc}, \text{col}))$.

$$0 < n \wedge \text{blank} \neq n\text{l} \Rightarrow \text{expand } V^\circ \sqsubseteq \text{expand } u (\pi_1 V^\circ) \quad (10.2b)$$

shortening the output of one **expand** step to a V -smaller string either keeps the whole step (**expand**) or drops its character and shortens its input string ($\pi_1 V^\circ$)




(a)	$\text{expand } V^\circ$	$\text{expand} = \text{istab outl fillunottab snocR}$ — definition of expand
(b)	$= ((\text{istab } \pi_1 \text{ fill}) \cup (\text{nottab snoc})) V^\circ$	$(\text{SuT})R = \text{SRuTR}$ — conditionals distribute
(c)	$= (\text{istab } \pi_1 \text{ fill } V^\circ) \cup (\text{nottab snoc } V^\circ)$	$\text{fill } V^\circ = \text{fill}$ (Ex. 10.4)




(a)	$= (\text{istab } \pi_1 \text{ fill}) \cup (\text{nottab snoc } V^\circ)$	$\text{snoc } V^\circ \sqsubseteq \text{snoc} \cup (\pi_1 V^\circ)$ (Ex. 10.4)
(b)	$\sqsubseteq (\text{istab } \pi_1 \text{ fill}) \cup (\text{nottab (snoc } \cup (\pi_1 V^\circ)))$	$\text{expand} = \text{istab outl fillunottab snocR}$ — definition of expand; $\text{nottab } R \sqsubseteq R$ — guard dropped
(c)	$\sqsubseteq \text{expand } u (\pi_1 V^\circ)$	

$$0 < n \wedge \text{blank} \neq n\mathbf{l} \Rightarrow \text{detab } V^\circ \sqsubseteq R^\circ \text{ detab}$$

(10.2c)

any V -shortening of **detab**'s output is **detab**'s output on an input no longer than the given one

(a)	$\text{detab } V^\circ$	detab is a fold: $\text{detab} = \alpha^\circ F(\text{detab}) [\text{nil}, \text{expand}] \triangleq \text{detab}$
(b)	$= [\text{nil}, \text{snoc}]^\circ F(\text{detab}) [\text{nil}, \text{expand}] V^\circ$	coproducts, and $\text{nil } V^\circ = \text{nil}$ (Exercise 10.4)
(c)	$= [\text{nil}, \text{snoc}]^\circ F(\text{detab}) [\text{nil}, \text{expand } V^\circ]$	the claim — (10.2b)
(d)	$\sqsubseteq [\text{nil}, \text{snoc}]^\circ F(\text{detab}) [\text{nil}, \text{expand } u (\pi_1 V^\circ)]$	distributing u ; the fold again, and the definition of F




(a)	$= \text{detab } u (\text{snoc}^\circ (\text{detab} \times 1) \pi_1 V^\circ)$	naturality of π_1 : $(\text{detab} \times 1) \pi_1 = \pi_1 \text{ detab}$; $\text{snoc}^\circ \pi_1$ is init
(b)	$= \text{detab } u (\text{snoc}^\circ \pi_1 \text{ detab } V^\circ)$	init is inductive, so $X \triangleq \text{detab } V^\circ$, a solution of $X \sqsubseteq \text{detab } u (\text{init } X)$, lies below the greatest one, prefix detab (induction on the input); $\text{prefix} \sqsubseteq R^\circ$
(c)	$\sqsubseteq R^\circ \text{ detab}$	

$$0 < n \wedge \text{blank} \neq n1 \Rightarrow (\mu X : \frac{[\text{nil}, \text{expand}]^\circ}{3} \text{est}(Q) F(X) [\text{nil}, \text{snoc}]) \sqsubseteq \frac{\text{detab}^\circ}{3} \text{est}(R) \quad (10.2d)$$

the shortest input **detab** expands to the given output is one pass along that output, holding each blank back and cashing the held blanks in for a tab wherever the column reaches a tab stop


the specification — $\triangleq \text{detab}; \text{detab entab} = 1$ and nothing shorter does



(a)	$\frac{\text{detab}^\circ}{3} \text{est}(R)$	at $Q \triangleq 1 + (V \times U)$: one character of input is decided at each step. $F(T, R) \alpha \alpha R$. $\text{detab prefix} \sqsubseteq R^\circ$ detab is FALSE, — at $n=8$, detab $[a, b, c, d, e, TB] = [a, b, c, d, e, BL, BL, BL]$, whose prefix $[a, b, c, d, e, BL, BL]$ is longer than any input giving it, and $\text{detab } V^\circ \sqsubseteq R^\circ$ detab holds. $\text{expand } V^\circ \sqsubseteq \text{expand } U$ ($\pi_1 V^\circ$) — shortening the output either leaves the last step alone or discards it
(b)	$\sqsubseteq \frac{[\text{nil}, \text{expand}]^\circ}{3} \text{est}(Q) F(X) [\text{nil}, \text{snoc}]$	Proposition 10.1: nil and expand have disjoint ranges. The greedy step is to emit a tab whenever a tab is legal, consuming all the blanks back to the previous tab stop
(c)	$\sqsubseteq \frac{\text{expand}^\circ}{3} \text{est}(V \times U) (X \times 1) \text{snoc}$	

$$\text{blank} \neq n1 \wedge \text{tb} \neq n1 \Rightarrow \langle \text{detab}, \text{detab col} \rangle = \langle [\text{base}, \text{step}] \rangle \quad (10.2e)$$

one pass over the input carries the output so far together with its column



(a)	$\langle \text{detab}, \text{detab col} \rangle$	both components are folds over the same input; step on $(\text{detab } x, \text{col}(\text{detab } x))$ is $(\text{detab}, \text{col detab})$ one character further — not a tabulated row
(b)	$= \langle [\text{base}, \text{step}] \rangle$	

Exercise 10.1

$$\text{blank} \neq n1 \wedge \text{tb} \neq n1 \Rightarrow \langle [\text{base}, \text{step}] \rangle (\text{convert}(l), r) \Leftrightarrow r = \text{loop}(\text{step}, (([]), 0), l)$$

the fold of $[\text{base}, \text{step}]$ over the input converted to a snoc-list is the left loop of **step** from **base**

$$\pi_1(\text{loop}(\text{step}, (([]), c_0), x)) = \text{loop}'((f, g), (c_0, x))$$

Exercise 10.1: when **step** only appends $f(c, a)$ to the output and moves the state by g , the output of the loop is $\text{loop}'(f, g)$, which never carries the output

$$a \neq \text{tb} \Rightarrow \text{expand}(y, b) = \text{snoc}(x, a) \Leftrightarrow \text{fill}(y) = \text{snoc}(x, a) \wedge b = \text{tb} \vee (y, b) = (x, a) \quad (10.2f)$$

expand produces $x \# [a]$ from a tab after a string that fills to it, or from x and a

$$0 < n \wedge \text{blank} \neq n1 \Rightarrow (\exists y. \text{fill}(y) = \text{snoc}(x, a)) \Leftrightarrow a = \text{blank} \wedge \text{col}(\text{snoc } x \text{ } a) \% n = 0$$

a string is a **fill** exactly when it ends in a blank on a tab stop

$$\text{entab}(x) = \text{entab}(\text{unfill } x) ++ \text{blanks}(\text{tbc } x) \quad (10.1)$$

(10.1): the output of **entab** is that of **unfill** x followed by the trailing blanks held back

$$\text{tbc}[] = 0$$


the empty string has no trailing blanks

$$\text{tbc}(\text{snoc}(x, a)) = \text{if } a = \text{blank} \wedge \text{col}(\text{snoc } x \text{ } a) \% n \neq 0 \text{ then } \text{tbc } x + 1 \text{ else } 0$$

a blank off a tab stop adds one held blank; anything else releases them all

$$\text{blank} \neq \text{nl} \Rightarrow \langle \text{tbc}, \text{col} \rangle = \langle \llbracket \text{base}, \text{op} \rrbracket \rangle \quad (10.2g)$$

one pass counts the held blanks and the column together




(a) $\langle \text{tbc}, \text{col} \rangle$

(b) $= \langle \llbracket \text{base}, \text{op} \rrbracket \rangle$

op on $\langle \text{tbc } x, \text{col } x \rangle$ is $\langle \text{tbc}, \text{col} \rangle$ one character further, from the equations for **tbc** — (10.1)

$$\text{blank} \neq \text{nl} \Rightarrow \text{triple} = \langle \llbracket \text{base}, \text{op} \rrbracket \rangle \quad (10.2h)$$

one pass carries the output up to the held blanks, their count and the column




(a) **triple**

(b) $= \langle \llbracket \text{base}, \text{op} \rrbracket \rangle$

op on **triple** x is **triple** one character further: (10.1) where the blanks are released — (10.1)

$$\text{entab} = \text{triple assocl } \pi_1 (1 \times \text{blanks}) \text{ cat} \quad (10.2i)$$

entab is **triple** with the held blanks appended to its output



(a) **entab**

(b) $= \text{triple assocl } \pi_1 (1 \times \text{blanks}) \text{ cat}$

(10.1) — (10.1)

§10.3. The minimum tardiness problem

(10.3a)

$\text{FX} = 1 + (X \times \text{Job})$, $\alpha \triangleq [\text{nil}, \text{snoc}]$ on schedules, $\beta \triangleq [\text{nil}, \text{snag}]$ on bags, **snag** putting a job into a bag; $\text{bagify} \triangleq \langle \beta \rangle : [\text{Job}] \rightarrow \text{Bag Job}$, $\text{H} = \text{bagify}^\circ$.

$\text{ct}, \text{dt}, \text{wt} : \text{Job} \rightarrow \text{Real}$ the completion, due and weighting quantities of a job; $\text{penalty}(\text{xs}, j) = (\text{sum}(\text{list}(\text{ct})(\text{xs})) + \text{ct}(j) - \text{dt}(j)) \times \text{wt}(j)$.

$\text{cost} \triangleq \frac{\text{prefix}}{\sup} P(\alpha^\circ [\text{zero}, \text{penalty}]) \text{ est}(\geq)$, $\text{cost} [] = 0$, $\text{cost} (\text{xs} \# [j]) = \text{bmax} (\text{cost } \text{xs}, \text{penalty} (\text{xs}, j))$, $R \triangleq \text{cost} \leq \text{cost}^\circ$.


$\text{perm} \triangleq \text{bagify } \text{bagify}^\circ = \langle [\text{nil}, \text{add}] \rangle$, $\text{add} (\text{xs}, j) = \text{ys} \# [j] \# \text{zs}$ for some $\text{xs} = \text{ys} \# \text{zs}$.

$k \triangleq [\text{zero}, \text{assocr } (1 \times ((\text{bagify}^\circ \times 1) \text{ penalty})) \text{ bmax}]$, $f \triangleq [\text{zero}, (\text{bagify}^\circ \times 1) \text{ penalty}]$, $Q \triangleq f \leq f^\circ$, $Q' \triangleq (\text{bagify}^\circ \times 1) \text{ penalty} \leq \text{penalty}^\circ (\text{bagify} \times 1)$.

$g \triangleq [\text{zero}, \text{penalty}]$ (10.5), $m \triangleq [\text{zero}, \pi_1 \text{ cost}]$ (B&dM's h , renamed as in (10.3e)) (10.6).

$$[\text{nil}, \text{snoc}] \text{ cost} = F(\langle \text{cost}, \text{bagify} \rangle) k \quad (10.3b)$$

the cost of a schedule is got from the cost and the bag of the schedule before its last job, and that job




(a)	$[\text{nil}, \text{snoc}] \text{ cost}$	$\text{cost}(xs \# [j]) = \text{bmax}(\text{cost } xs, \text{penalty}(xs, j))$, and $\text{penalty}(xs, j)$ reads only the bag of xs — $\triangleq \text{bagify}$
(b)	$= F(\langle \text{cost}, \text{bagify} \rangle) k$	

$$F(\geq \times 1) k \sqsubseteq k \geq$$

Exercise 10.5


a larger cost before the last job gives a larger cost after it



(a)	$F(\geq \times 1) k$	bmax is monotone in its first argument
(b)	$\sqsubseteq k \geq$	

$$F(R^\circ \circ n(\text{bagify } \text{bagify}^\circ)) [\text{nil}, \text{snoc}] \sqsubseteq [\text{nil}, \text{snoc}] R^\circ \quad (10.2)$$


improving the schedule before the last job, among schedules of the same bag, improves the whole schedule



(a)	$F(R^\circ \circ n(\text{bagify } \text{bagify}^\circ)) [\text{nil}, \text{snoc}]$	Proposition 9.3 at $S \triangleq \text{bagify}$, $\leq \triangleq$ — dp-context-mono, with (10.3b) and Exercise 10.5
(b)	$\sqsubseteq [\text{nil}, \text{snoc}] R^\circ$	

$$[\text{nil}, \text{snoc}] \text{ cost} = \langle g, m \rangle \text{ bmax} \quad (10.4)$$


the cost of a schedule is the larger of the penalty of its last job and the cost of the schedule before it



(a)	$[\text{nil}, \text{snoc}] \text{ cost}$	$\text{cost}(xs \# [j]) = \text{bmax}(\text{cost } xs, \text{penalty}(xs, j))$ and bmax commutes — $\triangleq \text{bagify}$
(b)	$= \langle g, m \rangle \text{ bmax}$	

$$(\forall j. 0 \leq ct(j)) \wedge (\forall j. 0 \leq wt(j)) \Rightarrow add \sqsubseteq \pi_1 R \quad (10.7)$$


putting one more job anywhere into a schedule never lowers its cost



(a)	add	induction on where j goes: every job after it starts later, so no penalty falls
(b)	$\sqsubseteq \pi_1 R$	

$$[nil, snag] \text{ bagify}^\circ = [nil, (\text{bagify}^\circ \times 1) \text{ add}] \quad (10.8)$$


the orderings of a bag with one more job are the orderings of the bag with that job put in anywhere



(a)	$[nil, snag] \text{ bagify}^\circ$	take out the last copy of j and put it back $\triangleq \text{bagify}$
(b)	$= [nil, (\text{bagify}^\circ \times 1) \text{ add}]$	

$$(\forall j. 0 \leq ct(j)) \wedge (\forall j. 0 \leq wt(j)) \Rightarrow [nil, snag] \text{ bagify}^\circ \sqsubseteq F(\text{bagify}^\circ) m \leq \text{cost}^\circ \quad (10.3c)$$


an ordering of a bag with one more job costs at least the cost of the ordering of the bag before its last job




(a)	$[nil, snag] \text{ bagify}^\circ$	(10.8) — (10.8)
(b)	$= [nil, (\text{bagify}^\circ \times 1) \text{ add}]$	(10.7) — (10.7)
(c)	$\sqsubseteq [nil, (\text{bagify}^\circ \times 1) \pi_1 R]$	definition of R , and $nil \sqsubseteq \text{zero} \leq \text{cost}^\circ$
(d)	$\sqsubseteq F(\text{bagify}^\circ) [zero, \pi_1 \text{ cost}] \leq \text{cost}^\circ$	definition of m
(e)	$= F(\text{bagify}^\circ) m \leq \text{cost}^\circ$	

$$(\forall j. 0 \leq ct(j)) \wedge (\forall j. 0 \leq wt(j)) \Rightarrow (Q \ n \ ([nil, snag] \ [nil, snag]^\circ)) \ F(bagify^\circ) \ [nil, snoc] \sqsubseteq F(bagify^\circ) \ [nil, snoc] \ R \quad (10.3)$$

a schedule of a bag ending in a job of least penalty is no worse than any schedule of the same bag




(a)	$(Q \ n \ ([nil, snag] \ [nil, snag]^\circ)) \ F(bagify^\circ) \ [nil, snoc]$	monotonicity of composition
(b)	$\sqsubseteq (Q \ F(bagify^\circ) \ [nil, snoc]) \ n \ ([nil, snag] \ [nil, snag]^\circ \ F(bagify^\circ) \ [nil, snoc])$	$\beta^\circ F(bagify^\circ) \alpha = bagify^\circ$, since $bagify = (\beta)$
(c)	$= (Q \ F(bagify^\circ) \ [nil, snoc]) \ n \ ([nil, snag] \ bagify^\circ)$	(10.3c)
(d)	$\sqsubseteq (Q \ F(bagify^\circ) \ [nil, snoc]) \ n \ (F(bagify^\circ) \ m \leq cost^\circ)$	modular law




(a)	$\sqsubseteq F(bagify^\circ) \ ((F(bagify) \ Q \ F(bagify^\circ)) \ n \ (m \leq cost^\circ \ [nil, snoc]^\circ)) \ [nil, snoc]$	choice of Q : $F(bagify) \ Q \ F(bagify^\circ) = g \leq g^\circ$
(b)	$= F(bagify^\circ) \ ((g \leq g^\circ) \ n \ (m \leq cost^\circ \ [nil, snoc]^\circ)) \ [nil, snoc]$	products: $\langle R, S \rangle \langle T, U \rangle^\circ = RT^\circ \ n \ SU^\circ$
(c)	$= F(bagify^\circ) \ \langle g \leq, m \leq \rangle \langle g, [nil, snoc] \ cost^\circ \rangle^\circ \ [nil, snoc]$	(10.3d)
(d)	$\sqsubseteq F(bagify^\circ) \ [nil, snoc] \ R$	

$$\langle g \leq, m \leq \rangle \langle g, [nil, snoc] \ cost^\circ \rangle^\circ \ [nil, snoc] \sqsubseteq [nil, snoc] \ R \quad (10.3d)$$

a job whose penalty is at most the last penalty, put after a schedule costing at most the cost before the last job, gives a schedule costing at most the whole



(a)	$\langle g \leq, m \leq \rangle \langle g, [nil, snoc] \ cost^\circ \rangle^\circ \ [nil, snoc]$	cost a map, so $1 \sqsubseteq cost \ cost^\circ$
(b)	$\sqsubseteq \langle g \leq, m \leq \rangle \langle g, [nil, snoc] \ cost^\circ \rangle^\circ \ [nil, snoc] \ cost \ cost^\circ$	$\alpha \ cost = (g, \alpha \ cost) \ bmax$
(c)	$= \langle g \leq, m \leq \rangle \langle g, [nil, snoc] \ cost^\circ \rangle^\circ \ \langle g, [nil, snoc] \ cost \rangle \ bmax \ cost^\circ$	$\langle g, \alpha \ cost \rangle$ simple
(d)	$\sqsubseteq \langle g \leq, m \leq \rangle \ bmax \ cost^\circ$	monotonicity of $bmax$




(a)	$\sqsubseteq \langle g, m \rangle \ bmax \leq cost^\circ$	(10.4) — (10.4)
(b)	$= [nil, snoc] \ cost \leq cost^\circ$	definition of R
(c)	$= [nil, snoc] \ R$	

$$(\forall j. 0 \leq \text{ct}(j)) \wedge (\forall j. 0 \leq \text{wt}(j)) \Rightarrow (\mu X : \frac{[\text{nil}, \text{snag}]^\circ}{3} \text{est}(Q) F(X) [\text{nil}, \text{snoc}]) \sqsubseteq \frac{\text{bagify}^\circ}{3} \quad (10.3e)$$

$\text{est}(R)$, $\text{schedule} = (\text{null} \rightarrow \text{nil}, \text{pick}(\text{schedule} \times 1) \text{snoc})$

an ordering of the given bag with least maximum penalty is got by taking a job of least penalty out of the bag, putting it last, and scheduling what is left the same way

the specification — $\triangleq \text{bagify}$



(a)	$\frac{\text{bagify}^\circ}{3} \text{est}(R)$	No greedy reduce exists — one would also solve every prefix of the input, and the best schedule of a prefix need not extend to a best schedule of the whole
(b)	$\sqsubseteq \frac{[\text{nil}, \text{snag}]^\circ}{3} \text{est}(Q) F(X) [\text{nil}, \text{snoc}]$	Proposition 10.1: nil and snag have disjoint ranges
(c)	$\sqsubseteq \frac{\text{snag}^\circ}{3} \text{est}(Q') (X \times 1) \text{snoc}$	$\text{pick} \sqsubseteq \frac{\text{snag}^\circ}{3} \text{est}(Q')$, a partial function, quadratic in the number of jobs
(d)	$\sqsubseteq \text{pick} (X \times 1) \text{snoc}$	

§10.4. The TeX problem

(10.4a)

$\text{intern} \triangleq \text{val round} : \text{Decimal} \rightarrow [0, 2^{16})$, $\text{val} \triangleq ([\text{zero}, \text{shift}])$, $\text{shift}(d, r) = (d+r)/10$, round r rounds $2^{16}r$ to the nearest integer: $\text{round } r = n \iff 2n-1 < 2^{17}r < 2n+1$.

$\text{interval } n = ((2n-1)/2^{17}, (2n+1)/2^{17})$, $r \text{ inrange } (a, b) \iff a < r < b$, $\text{round}^\circ = \text{interval inrange}$, $R \triangleq \text{length} \leq \text{length}^\circ$.


Interval the pairs (a, b) with $\text{Legal}(a, b) \iff 0 < b < 1$ and $a < b$ (10.9); step_legal is what types $\text{step} [\text{arb}, \text{step}] : 1 + (\text{Digit} \times \text{Interval}) \rightarrow \text{Interval}$, $\text{step}(d, (a, b)) = ((d+a)/10, (d+b)/10)$.

$\text{FX} = 1 + (\text{Digit} \times X)$, $\alpha \triangleq [\text{nil}, \text{cons}]$, $H \triangleq ([\text{arb}, \text{step}])^\circ$, $! : \text{Digit} \times \text{Interval} \rightarrow 1$, $Q \triangleq (l^\circ !^\circ r) \cup 1$ l, r are coprod-laws's injections into $\text{FX} = 1 + (\text{Digit} \times X)$, so $l : 1 \rightarrow \text{FX}$ and $r : \text{Digit} \times X \rightarrow \text{FX}$, $w \triangleq 2^{17}$.

$$([[\text{arb}, \text{step}]] \sqsubseteq \text{val inrange}^\circ \quad (10.4b)$$

every interval got by folding a decimal's digits with $[\text{arb}, \text{step}]$ has the decimal's value strictly inside it

fusion: it suffices that $[\text{zero}, \text{shift}] \text{ inrange}^\circ \sqsubseteq F(\text{inrange}^\circ) [\text{arb}, \text{step}]$




(a)	$[\text{zero}, \text{shift}] \text{ inrange}^\circ$	$[T, U]Z = [TZ, UZ]$, a coproduct law not tabulated in the note
(b)	$= [\text{zero inrange}^\circ, \text{shift inrange}^\circ]$	$\text{zero inrange}^\circ = \text{arb}$: the first condition, which determines arb
(c)	$= [\text{arb}, \text{shift inrange}^\circ]$	arithmetic: $10a-d < r < 10b-d \implies a < (d+r)/10 < b$ for $(a, b) = \text{step}(d, (10a-d, 10b-d))$; only $\exists$, since $(10a-d, 10b-d)$ satisfies (10.9) only when $d < 10b < d+1$
(d)	$\sqsubseteq [\text{arb}, (1 \times \text{inrange}^\circ) \text{step}]$	$F(S) [T, U] = [T, (1 \times S)U]$ read right to left: definition of F
(e)	$= F(\text{inrange}^\circ) [\text{arb}, \text{step}]$	


$$Q F(X) \alpha \sqsubseteq F(X) \alpha R$$

(10.4c)

choosing by Q among the one-step decompositions before building with $F(X)$ and α yields only decimals that are R -related to one built without choosing



(a)	$Q F(X) \alpha$	$(S \cup T)U = SU \cup TU$ — definition of Q ; composition distributes over $\cup$
(b)	$= (\mathbf{l}^\circ \mathbf{!}^\circ r F(X) \alpha) \cup (F(X) \alpha)$	$\mathbf{1} \sqsubseteq R$ — R is reflexive
(c)	$\sqsubseteq (\mathbf{l}^\circ \mathbf{!}^\circ r F(X) \alpha) \cup (F(X) \alpha R)$	$rF(X)\alpha \sqsubseteq \mathbf{!}\alpha R$ — $\mathbf{l}\alpha = \text{nil}$, and $\text{length}(\text{nil}) = 0$ is at most any length
(d)	$\sqsubseteq (\mathbf{l}^\circ \mathbf{!}^\circ \mathbf{!} \mathbf{l} \alpha R) \cup (F(X) \alpha R)$	$\mathbf{!}^\circ \mathbf{!} \sqsubseteq \mathbf{1}$ on $\mathbf{1}$ — universal property of $\mathbf{!}$



(a)	$\sqsubseteq (\mathbf{l}^\circ \mathbf{l} \alpha R) \cup (F(X) \alpha R)$	$\mathbf{l}F(X) = \mathbf{l}$ — definition of F
(b)	$= (\mathbf{l}^\circ \mathbf{l} F(X) \alpha R) \cup (F(X) \alpha R)$	$\mathbf{l}^\circ \mathbf{l} \sqsubseteq \mathbf{1}$ — $\mathbf{l}$ is simple
(c)	$\sqsubseteq (F(X) \alpha R) \cup (F(X) \alpha R)$	$S \cup S = S$ — $\cup$ is idempotent
(d)	$= F(X) \alpha R$	

(a)	$\frac{\text{intern}^\circ}{\exists} \text{ est}(R)$	round° is not a map, but interval is, so it comes out of the transpose
(b)	$= \text{interval } \frac{\text{inrange val}^\circ}{\exists} \text{ est}(R)$	the type restriction (10.9): a shortest decimal H gives an interval is a shortest one among all decimals inside it, and is inside it by (10.4b)
(c)	$\sqsupseteq \text{interval } \frac{H}{\exists} \text{ est}(R)$	$\frac{[\text{arb}, \text{step}]}{\exists}^\circ$ returns at most two elements — stop, or take one more digit — and $! \text{ nil} \sqsubseteq \text{cons } R^\circ$ makes it stop whenever stopping is legal
(d)	$\sqsubset \text{interval } \frac{[\text{arb}, \text{step}]}{\exists}^\circ \text{ est}(Q) \text{ F}(X) \alpha$	

- (10.4e)

§10. Appendix

§10.1. $P(S) \text{ est}(R) = (\exists S) n(\epsilon \setminus (SR^\circ))$

(10.1a)

(p, q) tabulates $W \triangleq (\exists S) n(\epsilon \setminus (SR^\circ))$, and $y \triangleq \frac{(p \exists S) n(qR)}{\exists}$, a map.

(10.1b)

$$\begin{array}{c} \boxed{\begin{array}{c} W \\ P(S) \text{ est}(R) \end{array}} \xleftarrow{p^\circ q = W, 1 \sqsubseteq y y^\circ} \begin{array}{c} \boxed{\begin{array}{c} p^\circ y \\ P(S) \end{array}} \\ \boxed{\begin{array}{c} y^\circ q \\ \text{est}(R) \end{array}} \end{array}$$

(10.1c)

$$\begin{array}{c} \boxed{\begin{array}{c} y^\circ q \\ \exists \end{array}} \xleftarrow{f^\circ \cdot \neg f^\circ, \cdot \exists \neg \exists} \boxed{\begin{array}{c} q \\ (p \exists S) n(qR) \end{array}} \xleftarrow{\Delta \neg n, f^\circ \cdot \neg f^\circ} \begin{array}{c} \boxed{\begin{array}{c} p^\circ q \\ \exists S \end{array}} \\ \boxed{\begin{array}{c} 1 \\ R \end{array}} \end{array}$$

R reflexive

$$\boxed{\begin{array}{c} \epsilon y^\circ q \\ R^\circ \end{array}} \xleftarrow{\cdot \exists \neg \exists, \circ, \text{meet}} \boxed{\begin{array}{c} R^\circ q^\circ q \\ R^\circ \end{array}} \quad \text{q simple}$$

(10.1d)

$$\begin{array}{c} \boxed{\begin{array}{c} p^\circ y \\ P(S) \end{array}} \xleftarrow{\Delta \neg n, \cdot T \neg / T, \circ, f^\circ \cdot \neg f^\circ} \begin{array}{c} \boxed{\begin{array}{c} p^\circ y \exists \\ \exists S \end{array}} \\ \boxed{\begin{array}{c} p \exists \\ y \exists S^\circ \end{array}} \end{array}$$

$$\boxed{\begin{array}{c} p^\circ y \exists \\ \exists S \end{array}} \xleftarrow{\cdot \exists \neg \exists, \text{meet}} \boxed{\begin{array}{c} p^\circ p \exists S \\ \exists S \end{array}} \quad \text{p simple}$$

$$\boxed{\begin{array}{c} p \exists \\ y \exists S^\circ \end{array}} \xleftarrow{\text{modular law}} \boxed{\begin{array}{c} p \exists \\ (p \exists) n(qRS^\circ) \end{array}} \xleftarrow{1 \sqsubseteq q q^\circ} \boxed{\begin{array}{c} q^\circ p \exists \\ RS^\circ \end{array}} \xleftarrow{\circ, T \cdot \neg T \setminus} \boxed{\begin{array}{c} W \\ \epsilon \setminus (SR^\circ) \end{array}} \quad \text{W's right half}$$

§10.2. $P(\text{est}(R)) \text{ est}(R) = P(\text{dom}(\text{est}(R))) \text{ union est}(R)$

(10.2a)

$$\begin{array}{c} \boxed{P(\text{est}(R)) \text{ est}(R)} \xrightarrow{\text{dom}(\text{est}(R)) \text{ est}(R) = \text{est}(R)} \boxed{P(\text{dom}(\text{est}(R)) \text{ est}(R)) \text{ est}(R)} \\ \xrightarrow{P \text{ a relator}} \boxed{P(\text{dom}(\text{est}(R))) P(\text{est}(R)) \text{ est}(R)} \xrightarrow{\text{est-711}} \boxed{P(\text{dom}(\text{est}(R))) \text{ union est}(R)} \end{array}$$

(10.2b)

$$\begin{array}{c} \boxed{\begin{array}{c} P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ P(\text{est}(R)) \text{ est}(R) \end{array}} \xleftarrow{(10.1b) \text{ at } S := \text{est}(R), \Delta \neg n, T \cdot \neg T \setminus} \boxed{\begin{array}{c} P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ \exists \text{est}(R) \end{array}} \\ \boxed{\begin{array}{c} \epsilon P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ \text{est}(R) R^\circ \end{array}} \xleftarrow{\text{modular law}} \boxed{\begin{array}{c} \exists (\exists n(\epsilon \setminus (\epsilon \setminus R^\circ))) \\ \exists (\exists n(\epsilon \setminus R^\circ)) \end{array}} \quad \text{count of } T \cdot \neg T \setminus \\ \boxed{\begin{array}{c} P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ \exists \text{est}(R) \end{array}} \xleftarrow{P(\text{dom}(\text{est}(R))) \sqsubseteq 1} \boxed{\begin{array}{c} \text{union est}(R) \\ \exists \text{est}(R) \end{array}} \\ \boxed{\begin{array}{c} \epsilon P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ \text{est}(R) R^\circ \end{array}} \xleftarrow{\epsilon \text{ lax natural}} \boxed{\begin{array}{c} \text{dom}(\text{est}(R)) \epsilon \text{ union est}(R) \\ \text{est}(R) R^\circ \end{array}} \\ \boxed{\begin{array}{c} \text{est}(R) \text{ est}(R)^\circ \epsilon \text{ union est}(R) \\ \text{est}(R) R^\circ \end{array}} \xleftarrow{\text{dom}(\text{est}(R)) \sqsubseteq \text{est}(R) \text{ est}(R)^\circ} \boxed{\begin{array}{c} \text{est}(R) \epsilon \text{est}(R) \\ \text{est}(R) R^\circ \end{array}} \quad \text{UP of est} \end{array}$$

§10.3. Shortest paths on a cylinder, on lists

(10.3a)

definition	type	note
$F(A, X) = A + A \times X$	$\mathcal{A} \rightarrow \mathcal{A}$	The base functor: one square of the new column, alone or in front of the path so far.
$L = \text{list}^+$	$\mathcal{A} \rightarrow \mathcal{A}$	A path is a non-empty list of squares, one per column crossed.
α the initial algebra	$F(\text{dCL}(L, E)) \rightarrow \text{dCL}(L, E)$	$\alpha(5) = [5]$ and $\alpha(1, [5]) = [1, 5]$: a path is started by one square, or extended by one.
N the n-tuple relator	$\mathcal{A} \rightarrow \mathcal{A}$	One component per row of a column, so the fold carries n answers at once.
$R \triangleq \text{sum} \leq \text{sum}^\circ$	$[\text{Nat}] \rightarrow [\text{Nat}]$	The cost of a path, which the cheapest minimises.
setify	$A[n] \rightarrow PA$	$\text{setify}(1, 2, 3, 4) = \{1, 2, 3, 4\}$ — which row a component came from is forgotten.
moves	$A[n] \rightarrow P(A[n])$	$\text{moves}(x) = \{\text{up}(x), x, \text{down}(x)\}$ — rotated up, unrotated, rotated down.
trans	$P(A[n]) \rightarrow (P A)[n]$	$\text{trans}\{(a, b, c), (x, y, z)\} = (\{a, x\}, \{b, y\}, \{c, z\})$ — component k of the result is the set of the k-th components.
zip	$F(a[n]) \rightarrow (F a)[n]$	$\text{zip}((1, 2, 3, 4), (\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\})) = ((1, \{[5]\}), (2, \{[6]\}), (3, \{[7]\}), (4, \{[8]\}))$.
$\text{cp} \triangleq \frac{F(1, \exists)}{\exists}$	$F(PA) \rightarrow PFA$	$\text{cp}(1, \{[5], [6], [8]\}) = (\{1, [5]\}, \{1, [6]\}, \{1, [8]\})$.
$\text{gen} \triangleq F(1, \text{moves trans } N(\text{union}))$ $\text{zip } N(\text{cp}(F(A, -)) E(\alpha))$	$F(NA, N(P(TA))) \rightarrow N(P(TA))$	$\text{gen}((1, 2, 3, 4), (\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\}))$ is worked out in (10.3.1b).
$\text{paths} \triangleq (\text{gen}) \text{ setify union}$	$T(NA) \rightarrow P(TA)$	$\text{paths}[(1, 2, 3, 4), (5, 6, 7, 8)]$ is the union of (10.3.1b)'s four sets: 12 paths, 3 from each entry row.
the specification $\text{paths est}(R)$	$[\text{Nat}[n]] \rightarrow [\text{Nat}]$	A cheapest path from the entry side to the exit side.

§10.3.1. $N(E(L A))$

```

1  5
2  6      [(1,2,3,4), (5,6,7,8)] : L N Nat
3  7
4  8

```

(10.3.1a)

$$(\text{gen})[(5, 6, 7, 8)] = (\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\})$$

(10.3.1b)

$$\begin{aligned} \text{gen}((1, 2, 3, 4), (\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\})) = \\ (\{ [1, 5], [1, 6], [1, 8] \}, \\ \{ [2, 5], [2, 6], [2, 7] \}, \\ \{ [3, 6], [3, 7], [3, 8] \}, \\ \{ [4, 5], [4, 7], [4, 8] \}) \end{aligned}$$


(10.3.1c)

N	the 4-tuple, one component per row of the entry column
E	the set of paths that can start in that row
L	a path is the non-empty list of the squares it crosses


row 1's set has no $[1,7]$ — from row 1 only rows 4, 1 and 2 are reachable — and it does have $[1,8]$, because the cylinder glues the bottom row to the top. That is **moves trans N(union)** of $\triangleq$ **paths**: component k collects the paths of rows $k-1, k, k+1$.

§10.3.2. The cross product $\text{cp} \triangleq \frac{F(1, \exists)}{\exists}$


(10.3.2a)



cp on all of $F(A, E(L A))$



the $A \times -$ summand: $\exists$ picks one path, $1\% \exists$ collects the results



the A summand, no E to distribute: $1\% \exists$ alone, $a \mapsto \{a\}$

(10.3.2b)


type	note
$F(A, -) = A + A \times -$	one wire carries the whole functor, both summands with it; the $\exists$ under it is $F(1, \exists)$
$\exists : E(L A) \rightarrow L A$	the $\exists$ inside $F(1, \exists)$: a set of paths, one of them — $L A$ is one path, A one square
$F(1, \exists) = 1 + 1 \times \exists$	$: A + A \times E(L A) \rightarrow A + A \times L A$, the relator acting on each summand
$\exists : E(A + A \times L A) \rightarrow A + A \times L A$	the $\exists$ under the bar: a set of cells, one of them — a different $\exists$
$\text{cp} : A + A \times E(L A) \rightarrow E(A + A \times L A)$	$\frac{R}{\exists}$ turns $R : X \rightarrow Y$ into $X \rightarrow E Y$
$\text{cp} = \frac{1}{\exists} E(F(1, \exists))$	what the picture draws: the unit makes the outer E , leaving one $\exists$ — pow-laws

(10.3.2c)

$w = (1, \{[5], [6], [8]\}) : A \times E(L A)$ the right summand
 $1 \times \exists : A \times E(L A) \rightarrow A \times L A$ 1 keeps the square 1, $\exists$ picks one path
 $w \mapsto (1, [5]), (1, [6]), (1, [8]) : A \times L A$ one output per path in w
 $\text{cp}(w) = \{(1, [5]), (1, [6]), (1, [8])\} : E(A + A \times L A)$ the three of them, collected

§10.3.3. $\text{gen} = F(1, \text{moves trans N(union)}) \text{ zip N(cp P}(\alpha))$

(10.3.3a)




(10.3.3b)

$u = ((1,2,3,4),(\{[5]\},\{[6]\},\{[7]\},\{[8]\}))$: $F(N A, N(E(L A)))$
 $F(1, \text{moves trans } N(\text{union}))$: 1 keeps the column, the path SETS move
 $\text{moves}(\{[5]\},\{[6]\},\{[7]\},\{[8]\})$
 $= (\{[6]\},\{[7]\},\{[8]\},\{[5]\}),$
 $(\{[5]\},\{[6]\},\{[7]\},\{[8]\}),$
 $(\{[8]\},\{[5]\},\{[6]\},\{[7]\})$: $E(N(E(L A)))$ down, unmoved, up
 $\text{trans}(\text{that})$
 $= (\{[6]\},\{[5]\},\{[8]\},\{[7]\},\{[6]\},\{[5]\}),$
 $\{[8]\},\{[7]\},\{[6]\},\{[5]\},\{[8]\},\{[7]\})$: $N(E(E(L A)))$ row k gets rows k-1, k, k+1
 $N(\text{union})(\text{that})$
 $= (\{[5],[6],[8]\},\{[5],[6],[7]\},$
 $\{[6],[7],[8]\},\{[5],[7],[8]\})$: $N(E(L A))$ every path into row k, none dropped
 $\text{zip}(\text{that})$
 $= ((1,\{[5],[6],[8]\}), (2,\{[5],[6],[7]\}),$
 $(3,\{[6],[7],[8]\}), (4,\{[5],[7],[8]\}))$: $N(F(A, E(L A)))$ each row: its square, and the paths it may be put in front of
 $N(\text{cp } P(\alpha))(\text{that})$
 $= (\{[1,5],[1,6],[1,8]\},\{[2,5],[2,6],[2,7]\},$
 $\{[3,6],[3,7],[3,8]\},\{[4,5],[4,7],[4,8]\})$: $N(E(L A))$ cp pairs the square with each path, α prefixes it: $\alpha(1,[5])=[1,5]$

§10.3.4. gen is an F-algebra; $(\text{gen}): \alpha(\text{gen})=F(1, (\text{gen}))\text{gen}$

(10.3.4a)

 α puts the column back on the list, then the fold reads all of itthe fold reads the rest under F , then one gen puts the column in front

(10.3.4b)

$$xs = [(1,2,3,4), (5,6,7,8)] = \alpha((1,2,3,4), [(5,6,7,8)]) : L(N A)$$

$$(\text{gen})(xs) = \text{gen}((1,2,3,4), (\text{gen})[(5,6,7,8)])$$

both halves are in (10.3.1b): the tail folds to $(\{[5]\},\{[6]\},\{[7]\},\{[8]\})$, one path per row and each of them one square long, and gen on it is (10.3.3b)'s walk.

§10.3.5. $Q : F(N A, N(L A)) \rightarrow N(L A)$, an $F(N A, -)$ -algebra: $Q=F(1, \text{moves trans } N(\text{est}(R))) \text{ zip } N(\alpha)$

(10.3.5a)

$Q : F(N A, N(L A)) \rightarrow N(L A)$: the fold's algebra: a new column, one path per row
 $u = ((1,2,3,4),([5],[6],[7],[8]))$: $F(N A, N(L A))$
 $F(1, \text{moves trans } N(\text{est}(R)))$: 1 keeps the column, the paths move
 $\text{moves}([5],[6],[7],[8])$
 $= (\{[6],[7],[8],[5]\},$
 $\{[5],[6],[7],[8]\},$
 $\{[8],[5],[6],[7]\})$: $E(N(L A))$ down, unmoved, up
 $\text{trans}(\text{that})$
 $= (\{[6],[5],[8]\},\{[7],[6],[5]\},$
 $\{[8],[7],[6]\},\{[5],[8],[7]\})$: $N(E(L A))$ row k gets rows k-1, k, k+1
 $N(\text{est}(R))(\text{that})$
 $= ([5],[5],[6],[5])$: $N(L A)$ the cheapest into each row – chosen BEFORE the new square
 $\text{zip}(\text{that})$
 $= ((1,[5]), (2,[5]), (3,[6]), (4,[5]))$: $N(F(A, L A))$ each row: its square, and the one predecessor that survived
 $N(\alpha)(\text{that})$
 $= ([1,5],[2,5],[3,6],[4,5])$: $N(L A)$ $\alpha(1,[5])=[1,5]$

$Q = [N(\text{wrap}), (1 \times \text{moves trans } N(\text{est}(R))) \text{ zip}' N(\text{cons})]$ (10.3.5b)	
at the last column Q starts one path per row, and at each earlier one it puts each square in front of the cheapest of the three kept paths it can step to — the algebra (10.3.5c)'s last step folds	
circuit — the fork is $F(NA, N(LA)) = NA + NA \times N(LA)$	Hinze–Marsden
$F(N A) (N (T A)) \text{---} \boxed{Q} \text{---} N(T A)$	
$FN(T A) \text{---} \boxed{F(1, \text{moves trans } N(\text{est}(R)))} \text{---} FN(T A) \text{---} \boxed{\text{zip}} \text{---} N(F A (T A)) \text{---} \boxed{N(\alpha)} \text{---} N(T A)$ the fusion condition (10.3.5d) read as a definition,	
$F(N A) (N (T A)) \text{---} \left\{ \begin{array}{l} \boxed{l} \text{---} N(p_0) \text{---} \boxed{N(\text{wrapA})} \\ \boxed{r} \text{---} q \text{---} \boxed{pm} \text{---} q \text{---} \boxed{\text{zip}'} \text{---} N(q_0) \text{---} \boxed{N(\text{consA})} \end{array} \right\} \text{---} N(T A)$ $\text{zip} = 1 + \text{zip}', \alpha = [\text{wrap}, \text{cons}]$	

paths $\text{est}(R) \exists (Q)$ setify $\text{est}(R)$ (10.3.5c)	
shortest paths on a cylinder: the dynamic programming approach to exhaustive search allows a path of least cost to be found in $O(n \times m)$ time; Q is (10.3.5b)'s algebra.	
circuit	Hinze–Marsden
$T(N\ A) \text{---} \boxed{\text{paths}} \text{---} P(T\ A) \text{---} \boxed{\text{est}(R)} \text{---} T\ A$	
$= T(N\ A) \text{---} \boxed{F(N\ A)\ (N\ (P\ (T\ A)))} \text{---} \boxed{\text{gen}} \text{---} NP(T\ A) \text{---} \boxed{\text{setify}} \text{---} PP(T\ A) \text{---} \boxed{\text{union}} \text{---} P(T\ A) \text{---} \boxed{\text{est}(R)} \text{---} T\ A$ $\triangleq \text{paths at paths}$	
$\sqsubseteq T(N\ A) \text{---} \boxed{F(N\ A)\ (N\ (P\ (T\ A)))} \text{---} \boxed{\text{gen}} \text{---} NP(T\ A) \text{---} \boxed{\text{setify}} \text{---} PP(T\ A) \text{---} \boxed{P(\text{est}(R))} \text{---} P(T\ A) \text{---} \boxed{\text{est}(R)} \text{---} T\ A$ $P(\text{est}(R))\ \text{est}(R) \sqsubseteq \text{union}\ \text{est}(R) \text{--- a minimum in each set, then a minimum of those; } R \text{ transitive}$	
$\sqsubseteq T(N\ A) \text{---} \boxed{F(N\ A)\ (N\ (P\ (T\ A)))} \text{---} \boxed{\text{gen}} \text{---} NP(T\ A) \text{---} \boxed{N(\text{est}(R))} \text{---} N(T\ A) \text{---} \boxed{\text{setify}} \text{---} P(T\ A) \text{---} \boxed{\text{est}(R)} \text{---} T\ A$ $\text{setify lax natural}$	
$\sqsubseteq T(N\ A) \text{---} \boxed{F(N\ A)\ (N\ (T\ A))} \text{---} \boxed{Q} \text{---} N(T\ A) \text{---} \boxed{\text{setify}} \text{---} P(T\ A) \text{---} \boxed{\text{est}(R)} \text{---} T\ A$ $\text{fusion at (10.3.5d)}$	

gen $N(\text{est}(R)) \exists F(1, N(\text{est}(R))) Q$ (10.3.5d)	
fusion: the condition for fusion in (10.3.5c)'s last step, used to derive a definition of Q .	
circuit	Hinze–Marsden
$F(N\ A)\ (N\ (P\ (T\ A))) \text{---} \boxed{\text{gen}} \text{---} NP(T\ A) \text{---} \boxed{N(\text{est}(R))} \text{---} N(T\ A)$	
$\sqsubseteq FNP(T\ A) \text{---} \boxed{F(1, N(\text{est}(R)))} \text{---} FN(T\ A) \text{---} \boxed{Q} \text{---} N(T\ A)$ $(7.13), \text{ then zip, trans, moves lax natural}$	

(7.13) is $F(1, \text{est}(R)) \alpha \sqsubseteq \text{cp } P(\alpha) \text{ est}(R)$, mon-thm71 at the map α with $\frac{F(1, \exists) \alpha}{\exists} = \text{cp } P(\alpha)$: extending every path in a set and then taking a minimum is beaten by extending one minimum. It is the crux here, not the greedy theorem.