

Companion to Bird & de Moor, Algebra of Programming

§5. Datatypes in Allegories

§5.1. Relators

(5.1a)

Every hom-set of an allegory is a poset, so an allegory is a **locally posetal 2-category**: the 2-cell from R to S IS $R \sqsubseteq S$. A **relator** $F : \mathcal{C} \rightarrow \mathcal{D}$ is a 2-functor between allegories:

$$F(1) = 1 \quad F(RS) = F(R)F(S) \quad R \sqsubseteq S \implies F(R) \sqsubseteq F(S)$$

Preserving $\circ$ is **not** asked for — $\circ$ is an identity-on-objects involution $\mathcal{C}^{\circ} \rightarrow \mathcal{C}$, no part of the 2-category.

(5.1b)

the statement

- 1 $\text{Map}(f) \implies \text{Map}(F(f))$ and $\text{Map}(f) \implies F(f^\circ) = F(f)^\circ$. Lemma 5.1
- 2 Over a **tabular** allegory a functor is a relator $\iff$ it preserves $\circ$. Theorem 5.1 the book's reverse direction (preserving $\circ \implies$ relator) needs the source to be **Rel**: the counterexample over a tabular allegory that is not **Rel** shows it can fail
- 3 $F(R^\circ) = F(R)^\circ$ for every R , so $F(R)^\circ$ needs no bracket.
- 4 Two relators agreeing on maps are equal. Corollary 5.1
- 5 $\text{Coreflexive}(X) \wedge \text{Coreflexive}(Y) \implies F(X \cap Y) = F(X) \cap F(Y)$ Ex 5.2
- 6 $F(R \cap S) \sqsubseteq F(R) \cap F(S)$, and strictly. Ex 5.2
- 7 $\text{PreservesRecip}(F) \implies F(\text{dom}(R)) = \text{dom}(F(R))$ Ex 5.5

The **power relator** $P \multimap xs \ P(R) \ ys \iff (\forall a \in xs. \exists b \in ys. a \ R \ b) \wedge (\forall b \in ys. \exists a \in xs. a \ R \ b)$ — is where the fourth is strict: for $R = \{(a_1, b_1), (a_2, b_2)\}$ and $S = \{(a_1, b_2), (a_2, b_1)\}$ the pair $(\{a_1, a_2\}, \{b_1, b_2\})$ is in $P(R) \cap P(S)$, while $R \cap S = \emptyset$.

§5.2. Relational products


(5.1)

The **fork** of $R : \mathcal{C} \rightarrow \mathcal{A}$ and $S : \mathcal{C} \rightarrow \mathcal{B}$ is $\langle R, S \rangle \triangleq R \pi_1^\circ \cap S \pi_2^\circ$, where (π_1, π_2) is the tabulation of τ .

(5.2a)

$$\langle R, S \rangle \pi_1 = \text{dom}(S) \ R \quad \langle R, S \rangle \pi_2 = \text{dom}(R) \ S \quad (5.6) \quad (5.7)$$


(5.2b)



A domain is coreflexive, so $\langle R, S \rangle \pi_1 \sqsubseteq R$, with equality exactly when S is entire; for maps both triangles commute and $\langle f, g \rangle$ is unique. In **Rel**, $c \in \langle R, S \rangle (a, b)$ iff $c \ R \ a$ and $c \ S \ b$ — copy c , then R on one strand and S on the other, which is $\triangleleft (R \otimes S)$ on the right.

No $\circ$ survives the translation. $\pi_1 = 1 \otimes \bullet$ discards the second component, so $\pi_1^\circ = 1 \otimes \bullet$ **creates** one out of nothing, and $\cap$ is copy, run both, merge. Draw that and the created strands — the two dots with no left end, and the crossing they force — are merged against real ones, which is the monoid's unit law:

(5.2c)




§5.2.1. Relational product $R \times S$

(5.2)

$R \times S \triangleq \langle \pi_1 R, \pi_2 S \rangle$, a relator in each argument but no longer a categorical product.


(5.2.1a)



Right-then-up is $(R \times S)\pi_1$, up-then-right is $\pi_1 R$, and $(R \times S)\pi_1 \sqsubseteq \pi_1 R$, equality when S is entire. In $\mathbf{Rel}$, $(c, d) \in (R \times S) \iff (a, b) \in R$ and $d \in S \iff b$ — two strands side by side, no copy dot: $R \times S = R \otimes S$.

§5.2.2. Absorption

For $X : E \rightarrow C$ and $Y : E \rightarrow D$, $\langle X, Y \rangle (R \times S) = \langle XR, YS \rangle$. Both sides are this picture:



(5.3)

(5.2.2a)

	the law	picture
1	$\langle X, Y \rangle (R, S)^\circ = (X R^\circ) \cap (Y S^\circ)$	
2	$\langle R, S \rangle^\circ \langle U, V \rangle \sqsubseteq (R^\circ U) \times (S^\circ V)$	
3	$\text{Map}(f) \implies f \langle R, S \rangle = \langle f R, f S \rangle$ f a map; it fails for an arbitrary arrow;	
4	$X \langle R, S \rangle \sqsubseteq \langle X R, X S \rangle$ the half of the row above that holds for every X	
5	$F(R \times S) \text{ unzip}(F) \sqsubseteq \text{unzip}(F) (F(R) \times F(S))$ $\text{unzip}(F) \triangleq \langle F(\pi_1), F(\pi_2) \rangle$; only $\sqsubseteq$ for an arbitrary R, S	
6	$g = \text{curry}(f) \iff (g \times 1) \text{ eval} = f$ reading $\times$ as the relational product, does $\mathbf{Rel}$ have exponentials?	

§5.3. Relational coproducts

(5.3a)

	the statement	picture
	$R : a_1 \rightarrow c, S : a_2 \rightarrow c$	
1	$[R, S] \triangleq (l^\circ R) \cup (r^\circ S)$ (5.9) The tape is the union — a particle entering at $A+B$ takes exactly one branch — and the two mirrored boxes are what makes the branches disjoint.	
2	$[R, S] = [\frac{R}{\exists}, \frac{S}{\exists}] \exists$	
3	$l [R, S] = R, r [R, S] = S$ $l T = R \wedge r T = S \implies T = [R, S]$	
4	$U : a_1 \rightarrow d, V : a_2 \rightarrow d$ $[U, V]^\circ [R, S] = (U^\circ R) \cup (V^\circ S)$ (5.11)	
5	$Z : c \rightarrow d$ $[R, S] Z = [R Z, S Z]$	


(5.3b)

	the statement	picture
	$R : a_1 \rightarrow b_1, S : a_2 \rightarrow b_2$	
1	$R + S \triangleq [R l, S r]$ (5.10)	
2	$l l^\circ = 1, r r^\circ = 1$ Ex 5.12	$a_1 \text{---} l \text{---} a \text{---} l \text{---} a_1 = a_1 \text{---} a_1 \quad a_2 \text{---} r \text{---} a \text{---} r \text{---} a_2 = a_2 \text{---} a_2$
3	$l r^\circ = 0, r l^\circ = 0$ Ex 5.12	$a_1 \text{---} l \text{---} a \text{---} r \text{---} a_2 = a_1 \text{---} 0 \text{---} a_2$
4	$(l^\circ l) \cup (r^\circ r) = 1$ Ex 5.12	


§5.3.1. $[R, S] \triangleq [\frac{R}{\exists}, \frac{S}{\exists}] \exists$

The universal-property row is not free: l, r were only ever asked to be a coproduct of **maps**. They stay one once every arrow is allowed because $\exists$ sends an arrow $A \rightarrow C$ to a map $A \rightarrow PC$ reversibly, so the map coproduct can be applied underneath it. For any $T : A+B \rightarrow C$,

(5.3.1a)



(5.3.1b)



Nothing here holds only up to $\cong$: every triangle commutes on the nose, which is the difference from the fork above. The border spells $[R, S] = [\frac{R}{\tilde{l}}, \frac{S}{\tilde{r}}] \exists$, and pushing $\exists$ into the union that $[\cdot, \cdot]$ on maps already is turns that back into the definition, $(\tilde{l} \circ \frac{R}{\tilde{l}} \cup \tilde{r} \circ \frac{S}{\tilde{r}}) \exists = \tilde{l} \circ R \cup \tilde{r} \circ S$.

(5.3.1c)

the law	picture
1 prove (5.11), and say why duality does not carry it over from the product law	
2 $(R + S) \cap ([P, Q] [U, V]^\circ)$ $= (R \cap (P \cup^\circ)) + (S \cap (Q \cup^\circ))$ $[P, Q] [U, V]^\circ$ is a full 2×2 of composites; $R+S$ is diagonal, so the meet cuts the two off-diagonal branches	

§5.4. The power relator $P(R)$

(5.4a)

For $R : A \rightarrow B$,

$$P(R) \triangleq (\epsilon \setminus (R \epsilon)) \cap ((\exists R) / \exists)$$

$$E(R) \triangleq \frac{\exists R}{\exists} = ((\exists R) / \exists) \cap (\exists / (\exists R))^\circ$$


$$xs \ P(R) \ ys \iff (\forall a \in xs. \exists b \in ys. a \ R \ b) \wedge (\forall b \in ys. \exists a \in xs. a \ R \ b)$$

Every element of xs is related by R to some element of ys , and conversely.

(5.4b)

	in Rel	in words	$\text{im}(xs)$	partner
1 $(\exists R) / \exists$	$\forall y \in ys. \exists x \in xs. x \ R \ y$	$\forall y. \text{ some } xs \ R \ y$	$ys \subseteq \text{im}(xs)$	every y has a partner
2 $(\exists / (\exists R))^\circ$	$\forall y. (\exists x \in xs. x \ R \ y) \rightarrow y \in ys$	every y with some $xs \ R \ y$ is in ys	$ys \supseteq \text{im}(xs)$	ys leaves out no partner
3 $E(R)$	$ys = \{y \mid \exists x \in xs. x \ R \ y\}$	$ys = \text{every } y \text{ with some } xs \ R \ y$	$ys = \text{im}(xs)$	every y has a partner, and none is left out
4 $((\exists R^\circ) / \exists)^\circ$	$\forall x \in xs. \exists y \in ys. x \ R \ y$	$\forall x. x \ R \ \text{some } ys$	—	every x has a partner
5 $P(R)$	$\forall y \in ys. \exists x \in xs. x \ R \ y$ and $\forall x \in xs. \exists y \in ys. x \ R \ y$	$\forall y. \text{ some } xs \ R \ y$ and $\forall x. x \ R \ \text{some } ys$	—	every x and every y has a partner

(5.4c)



xs	$(\exists R) / \exists$	$(\exists / (\exists R))^\circ$	$E(R)$	$((\exists R^\circ) / \exists)^\circ$	$P(R)$
1 $\emptyset$	$\emptyset$	all 8 subsets of abc	$\emptyset$	all 8 subsets of abc	$\emptyset$
2 $\{1\}$	$\emptyset, a, b, ab$	ab, abc	ab	a, b, ab, ac, bc, abc	a, b, ab
3 $\{2\}$	$\emptyset$	all 8 subsets of abc	$\emptyset$	none	none
4 $\{3\}$	$\emptyset, c$	c, ac, bc, abc	c	c, ac, bc, abc	c
5 $\{1, 2\}$	$\emptyset, a, b, ab$	ab, abc	ab	none	none
6 $\{1, 3\}$	all 8 subsets of abc	abc	abc	ac, bc, abc	ac, bc, abc
7 $\{2, 3\}$	$\emptyset, c$	c, ac, bc, abc	c	none	none
8 $\{1, 2, 3\}$	all 8 subsets of abc	abc	abc	none	none

(5.4d)

	law	the reading
1	$X \sqsubseteq P(R) \iff X \sqsubseteq \exists \exists R \text{ and } X^\circ \sqsubseteq \exists \exists R^\circ$	One containment, and the same one at R° — which is the definition read off the two divisions. Hence $P(R^\circ) = P(R)^\circ$, and $R \sqsubseteq S \implies P(R) \sqsubseteq P(S)$.
2	$P(1) = 1$	The straightness axiom verbatim: extensionality is P 's unit law.
3	$\text{Map}(f) \implies P(f) = E(f)$	In Rel, $xs \ P(f) \ ys \iff ys = \{f(a) \mid a \in xs\}$. The half at f° says every $a \in xs$ has its $f(a)$ on ys ; f has just the one image per a , so that already says ys contains everything xs reaches, which is the fraction's second half. For a map the two definitions coincide.
4	$P(R \ S) = P(R) \ P(S)$	$\sqsupseteq$ is the division cancellation laws. $\sqsubseteq$ is the one law in this section that is not a calculation: it needs a tabulation of $P(RS)$.

§5.4.1. Type relator

(5.4.1a)

Let F be a binary relator with initial type (α, T) , so T is a type functor. $F(R, S)$ is its action on a pair, and $F(X)$ abbreviates $F(1, X)$, the F of the reduce section. For every object A the initial algebra is $\alpha : F(A, TA) \rightarrow TA$, among the maps. T acts on an arrow $R : A \rightarrow B$ by

$$T(R) = \langle F(R, 1) \ \alpha \rangle$$

(5.4.1b)

	name	law	what it says
1	the defining equation	$T(R) = \langle F(R, 1) \ \alpha \rangle$	Rebuild the structure with α , applying R to the parameter on the way.
2	type relator	$T(R)^\circ = T(R^\circ)$	A datatype acts on relations, not only on maps — the map of the converse is the converse of the map.


§5.5. Relational catamorphisms

(5.5a)

let F be a relator and has **initial algebra** $\alpha : F(T) \rightarrow T$ in the subcategory of functions. α is also initial in the allegory:

§5.5.1. The defining equation

(5.5.1a)



An F -algebra is a **weakened** natural transformation. A transformation $F \Rightarrow 1$ on $\mathcal{A}$ would need a component $FX \rightarrow X$ at every object and a commuting square at every arrow, but F -Algebra only need it works on T and A .

(5.5.1b)


$$([c, f]) = \text{reduce}(c, f)$$

catamorphism: arrows of the form $([f])$ are called catamorphisms, and these two examples reveal the notion to be a familiar idea in abstract clothing

1	part	Nat	[A]
2	datatype	$\text{Nat} ::= \text{zero} \mid \text{succ Nat}$	$[A] ::= \text{nil} \mid \text{cons}(A, [A])$
3	base functor F	$F(X) = 1 + X$	$F(X) = 1 + A \times X$
4	initial algebra α	$\alpha = [\text{zero}, \text{succ}]$ $: 1 + \text{Nat} \rightarrow \text{Nat}$	$\alpha = [\text{nil}, \text{cons}]$ $: 1 + A \times [A] \rightarrow [A]$
5	the fold, pointwise	$([c, f])(\text{zero}) = c$ $([c, f])(\text{succ}(n)) = f([c, f](n))$	$([c, f])(\text{nil}) = c$ $([c, f])(\text{cons}(a, x)) = f(a, [c, f](x))$

§5.5.2. $(R) = ([\frac{F(\exists)R}{\exists}])\exists$

(5.5.2a)



(5.12)


$$\begin{array}{c}
 \boxed{\frac{\alpha X}{F(X)R}} \iff \boxed{\frac{\alpha X}{\frac{F(X)R}{\exists}}} \iff \boxed{\frac{\alpha X}{\frac{F(\frac{X}{\exists})R}{\exists}}} \\
 \iff \boxed{\frac{\alpha \frac{X}{\exists}}{F(\frac{X}{\exists}) \frac{F(\exists)R}{\exists}}} \iff \boxed{\frac{\frac{X}{\exists}}{([\frac{F(\exists)R}{\exists}]})} \iff \boxed{\frac{X}{([\frac{F(\exists)R}{\exists}])\exists}}
 \end{array}$$

relator, fusion twice reduce of maps $\cdot \exists \dashv \frac{_}{\exists}$

the last two rows at $X := (R)$:

$$\alpha X = F(X) R \iff \frac{X}{\exists} = \langle \frac{F(\exists) R}{\exists} \rangle \quad (5.5.2b)$$

in a tabular allegory, with F a relator and $\alpha : F(T) \rightarrow T$ its initial algebra: a relation X out of T satisfies the fold equation $\alpha X = F(X) R$ of R exactly when $\frac{X}{\exists}$ is the fold of the map $\frac{F(\exists) R}{\exists}$



(a)	$\frac{X}{\exists}$	α iso; $\alpha X = F(X) R$
(b)	$= \frac{\alpha^0 F(X) R}{\exists}$	α^0 is a map because α is an iso, so it leaves Λ : $\frac{\alpha^0 F(X) R}{\exists} = \alpha^0$ $\frac{F(X) R}{\exists} \text{ --- Map}(f)$ $\implies \frac{f R}{\exists} = f \frac{R}{\exists}$
(c)	$= \alpha^0 \frac{F(X) R}{\exists}$	$X = \frac{X}{\exists} \text{ --- } \frac{R}{\exists} \text{ --- } \exists = R$
(d)	$= \alpha^0 \frac{F(\frac{X}{\exists}) R}{\exists}$	$F(\frac{X}{\exists}) = F(\frac{X}{\exists}) F(\exists)$ as F is a functor, and $F(\frac{X}{\exists})$ is a map (a relator sends maps to maps), so it leaves Λ --- $\triangleq$ relator, $\text{Map}(f)$ $\implies \frac{f R}{\exists} = f \frac{R}{\exists}$
(e)	$= \alpha^0 F(\frac{X}{\exists}) \frac{F(\exists) R}{\exists}$	fold uniqueness, $\frac{F(\exists) R}{\exists}$ being a map --- (2.6a)
(f)	$= \langle \frac{F(\exists) R}{\exists} \rangle$	

§5.6. Combinatorial functions

(5.6a)


	definition	type	note
1	$[A] ::= \text{nil} \mid \text{cons}(A, [A])$	$\mathcal{A} \rightarrow \mathcal{A}$	The list type, under the short name it keeps.
2	$\text{list}(R) = ([\text{nil}, (R \times 1) \text{ cons}])$	$[A] \rightarrow [B]$	The relator's action on $R : A \rightarrow B$: one R per element, the shape untouched.
3	$\text{subseq} = ([\text{nil}, \text{cons} \cup \pi_2])$	$[A] \rightarrow [A]$	$xs \text{ subseq } ys$: ys is xs with elements dropped — cons keeps the head, π_2 drops it.
4	$\text{prefix} = ([\text{nil}, \text{nil} \cup \text{cons}])$ $\text{prefix} = \text{cat}^\circ \pi_1 = \text{init}^*$	$[A] \rightarrow [A]$	ys is an initial segment of xs ; the first nil is where it stops early. $\text{init} \triangleq \text{snoc}^\circ \pi_1$.
5	$\text{suffix} = \text{cat}^\circ \pi_2 = \text{tail}^*$	$[A] \rightarrow [A]$	The dual, $\text{tail} \triangleq \text{cons}^\circ \pi_2$; as a reduce it needs snoc -lists.
6	$\text{partition} = \text{concat}^\circ$	$[A] \rightarrow [[A]]$	This cat is restricted to $[A]^+ \times [A] \rightarrow [A]$, so ys is a list of non-empty segments of xs .
7	$\text{concat} = ([\text{nil}, \text{cat}])$	$[[A]] \rightarrow [A]$	Joins the segments back up, which is why its converse splits a list.
8	inits	$[A] \rightarrow [[A]]$	Implements $\frac{\text{prefix}}{\exists}$, listing the prefixes by increasing length.
9	tails	$[A] \rightarrow [[A]]$	Implements $\frac{\text{suffix}}{\exists}$ by decreasing length — the opposite order.
10	$\text{cpp}, \text{cpl}, \text{cpr}$	$[A] \times [B] \rightarrow [A \times B]$ $[A] \times B \rightarrow [A \times B]$ $A \times [B] \rightarrow [A \times B]$	Every pair (a, b) with a from the first list and b from the second; cpl , cpr keep one side a single element.
11	$\text{cpp setify} = (\text{setify} \times \text{setify})$ $\frac{\exists \times \exists}{\exists}$ $\text{cpl setify} = (\text{setify} \times 1) \frac{\exists \times 1}{\exists}$ $\text{cpr setify} = (1 \times \text{setify}) \frac{1 \times \exists}{\exists}$	$[A] \times [B] \rightarrow P(A \times B)$	Forgetting the order with setify , cpp is the transpose of $\exists \times \exists$.
12	$\text{cp}(F) \triangleq \frac{F(\exists)}{\exists}$	$F(PA) \rightarrow PFA$	$\text{cp}(F)$, the transpose of a relator's action on $\exists$; cpp implements it at $F(A) = A \times A$.
13	$\text{cp}(\text{list}) = ([\frac{[\text{nil}, (\exists \times \exists) \text{ cons}]}{\exists}])$ $\frac{[\text{nil}, (\exists \times \exists) \text{ cons}]}{\exists} = [\text{nil} \frac{1}{\exists}, \frac{\exists \times \exists}{\exists} \text{ E}(\text{cons})]$	$\text{list}(PA) \rightarrow P(\text{list}(A))$	$\text{cp}(\text{list})$: (5.12) at $\text{list}(\exists) = ([\text{nil}, (\exists \times 1) \text{ cons}])$, then the algebra expanded.
14	$\text{cplist} = ([\text{nil wrap}, \text{cpp list}(\text{cons})])$	$[[A]] \rightarrow [[A]]$	$\text{cp}(\text{list})$ with every set a list: cpp in place of $\frac{\exists \times \exists}{\exists}$.

glue concat $\sqsubseteq (1 \times \text{concat}) \text{ cons}$


(5.6b)

putting **a** on the front of the first segment and flattening (**glue concat**) gives at most flattening the segments and putting **a** on the front of the result (**(1×concat) cons**)

the starting composite



(a)	glue concat	definition of glue
(b)	$\stackrel{\Delta}{=} (1 \times \text{cons}^\circ) \text{ assocl } \text{cons} \times 1 \text{ cons } \text{concat}$	since $\text{concat} = ([\text{nil}, \text{cat}])$
(c)	$= (1 \times \text{cons}^\circ) \text{ assocl } \text{cons} \times 1 (1 \times \text{concat}) \text{ cat}$	naturality of assocl
(d)	$= (1 \times (\text{cons}^\circ (1 \times \text{concat}))) \text{ assocl } \text{cons} \times 1 \text{ cat}$	since $\text{concat} = ([\text{nil}, \text{cat}])$



(a)	$\sqsubseteq (1 \times \text{concat}) (1 \times \text{cat}^\circ) \text{ assocl } \text{cons} \times 1 \text{ cat}$	since $(\text{cons} \times 1) \text{ cat} = \text{assocr } (1 \times \text{cat}) \text{ cons}$
(b)	$= (1 \times \text{concat}) (1 \times \text{cat}^\circ) \text{ assocl } \text{assocr } (1 \times \text{cat}) \text{ cons}$	since $\text{assocl } \text{assocr} = 1$ and $\text{cat}^\circ \text{ cat} \sqsubseteq 1$
(c)	$\sqsubseteq (1 \times \text{concat}) \text{ cons}$	

§5.6.1. $\frac{\text{subseq}}{\ni} = ([\text{nil } \frac{1}{\ni}, (\frac{1 \times \ni}{\ni} \text{ E}(\text{cons}), \pi_2) \text{ cup}])$

(5.6.1a)



$\text{cup} \stackrel{\Delta}{=} \frac{\pi_1 \ni \cup \pi_2 \ni}{\ni} : \text{PA} \times \text{PA} \rightarrow \text{PA}$, so $\frac{R \cup S}{\ni} = (\frac{R}{\ni}, \frac{S}{\ni}) \text{ cup}$.

$$\frac{F(\exists) [\text{nil}, \text{cons} \cup \pi_2]}{\exists} = [\text{nil} \frac{1}{\exists}, \frac{(1 \times \exists) (\text{cons} \cup \pi_2)}{\exists}] \quad (5.6.1b)$$

the set of lists the algebra builds is, from nothing, just **nil**, and from a head and a set of tails, every tail in the set with the head put on or left off — (5.12) at **subseq** = $([\text{nil}, \text{cons} \cup \pi_2])$, **subseq** = $([\text{nil}, \text{cons} \cup \pi_2])$.



circuit — the fork is $F([A]) = 1 + A \times [A]$: **nil** above, the pair below

Hinze–Marsden



$$\frac{F(\exists) [\text{nil}, \text{cons} \cup \pi_2]}{\exists}$$



the π_2 operand of $\text{cons} \cup \pi_2$ under the $1 \times \exists$ summand of $F(\exists)$, i.e. $(1 \times \exists) \pi_2$



$$\frac{(1 + 1 \times \exists) [\text{nil}, \text{cons} \cup \pi_2]}{\exists}$$

$$F(X) = 1 + A \times X \quad (5.6a.1)$$



the same operand under $1 + 1 \times \exists$, whose $1 \times \exists$ summand it sits in



$$\frac{[\text{nil}, (1 \times \exists) (\text{cons} \cup \pi_2)]}{\exists}$$

$$\begin{aligned} R + S &\triangleq [Rl, Sr], \quad l[R, S] = R, \quad r[R, S] = S \quad [R, S] \triangleq \\ &(l^\circ R) \cup (r^\circ S), \quad l[R, S] = R, \quad r[R, S] = S, \quad lT = R \wedge rT = S \\ &\Rightarrow T = [R, S] \end{aligned}$$



the π_2 operand of the second arm $(1 \times \exists) (\text{cons} \cup \pi_2)$



$$\frac{[\text{nil} \frac{1}{\exists}, (1 \times \exists) (\text{cons} \cup \pi_2)]}{\exists}$$

$$(5.3.1a) \text{ at } T := [\text{nil}, (1 \times \exists) (\text{cons} \cup \pi_2)]$$

the π_2 operand under its $\frac{1}{\exists}$, the arm (5.6.1c)'s square rewrites, $(1 \times \exists) \pi_2 = \pi_2 \exists$



$$[\text{nil} \frac{1}{\exists}, \frac{(1 \times \exists) (\text{cons} \cup \pi_2)}{\exists}]$$

$$\text{Map}(f) \wedge 1 \square = \exists \square$$


$$\Rightarrow \frac{f}{\exists} = f \frac{1}{\exists}, \quad \frac{f}{\exists} = f \frac{1}{\exists} \text{ for } f \text{ a map, at } f := \text{nil}$$

the same operand; the two rows differ only in the **nil** arm

(5.6.1c)

$$\begin{array}{ccc} A \times P[A] & \xrightarrow{\pi_2} & P[A] \\ \downarrow 1 \times \exists & & \downarrow \exists \\ A \times [A] & \xrightarrow{\pi_2} & [A] \end{array}$$

$\frac{(1 \times \exists) (\text{cons} \cup \pi_2)}{\exists} = \left(\frac{1 \times \exists}{\exists} E(\text{cons}), \pi_2 \right) \text{cup}$ power transpose of join: the power transpose of the join of two relations is $\left(\frac{R}{\exists}, \frac{S}{\exists} \right) \text{cup}$, where cup is the function that returns the union of two sets	(5.6.1d)
circuit $\frac{(1 \times \exists) (\text{cons} \cup \pi_2)}{\exists}$ (5.6.1b) $\frac{F(\exists) [\text{nil}, \text{cons} \cup \pi_2]}{\exists} = [\text{nil} \frac{1}{\exists}, \frac{(1 \times \exists) (\text{cons} \cup \pi_2)}{\exists}]$'s second branch	Hinze–Marsden the cons operand of cons $\cup$ π_2
$\frac{(1 \times \exists) \text{cons} \cup (1 \times \exists) \pi_2}{\exists}$ $T(X_1 \cup X_2) = TX_1 \cup TX_2$ — (0.2a.8)	the π_2 operand of $(1 \times \exists) \text{cons} \cup (1 \times \exists) \pi_2$
$\frac{(1 \times \exists) \text{cons} \cup \pi_2 \exists}{\exists}$ $(1 \times \exists) \pi_2 = \pi_2 \exists$ — (5.2.1a) at π_2, an equality because 1 is entire	the π_2 operand of $(1 \times \exists) \text{cons} \cup \pi_2 \exists$
$\frac{(1 \times \exists) \text{cons} \cup \pi_2 \exists}{\exists} \text{cup}$ $\frac{R \cup S}{\exists} = \left(\frac{R}{\exists}, \frac{S}{\exists} \right) \text{cup} \triangleq \text{cup}$	the cons operand under its $\frac{1}{\exists}$
$\frac{(1 \times \exists) E(\text{cons}), \pi_2}{\exists} \text{cup}$ $\frac{S}{\exists} E(R) = \frac{S}{\exists} R$, absorption $\frac{S}{\exists} E(R) = \frac{SR}{\exists}$ at $S := 1 \times \exists, R := \text{cons}$; fusion and $\frac{\exists}{\exists} = 1$ on the π_2 operand	the π_2 operand, bare π_2



$$[\frac{\text{nil}}{\exists}, \frac{(1 \times \exists) E(\text{cons}), \pi_2}{\exists} \text{cup}]$$

which writes **Pcons**; **cons** is a map, and there $P(\text{cons}) = E(\text{cons})$ — $\text{Map}(f)$

$$\Rightarrow P(f) = E(f) .$$

(5.6.1e)

§5.7. Lax natural transformations (LaT)


(5.7a)

For relators $G, F : \mathcal{C} \rightarrow \mathcal{D}$ and components $\varphi_A : GA \rightarrow FA$, φ is **lax at** $R : A \rightarrow B$ when $G(R)\varphi_B \sqsubseteq \varphi_A F(R)$ — both sides $GA \rightarrow FB$, the left through the component at B , the right through the one at A .










$\varphi : G \Rightarrow F$ is a **lax natural transformation** (LaT) when it is lax at **every** R . (5.13)

Lax at every **map** already gives LaT, and at a map the inequation is an equality $G(f)\varphi = \varphi F(f)$: laxness is about relations only.

(5.7b)




(5.7c)

closed under	commutative diagram	string diagram
1 composition $\psi\phi$	 $H(R)\psi_B \sqsubseteq \psi_A G(R)$ and $G(R)\phi_B \sqsubseteq \phi_A F(R)$ give $H(R)(\psi_B\phi_B) \sqsubseteq (\psi_A\phi_A)F(R)$	
2 horizontal composition $\chi\circ\phi$	 $\phi : G \Rightarrow F$ with $G, F : \mathcal{C} \rightarrow \mathcal{D}$ and $\chi : L \Rightarrow K$ with $L, K : \mathcal{D} \rightarrow \mathcal{E}$ give $\chi\circ\phi : L\circ G \Rightarrow K\circ F$, the family $A \mapsto \chi_{GA}K(\phi_A)$ $L(G(R))\chi_{GB} \sqsubseteq \chi_{GA}K(G(R))$ is χ lax at $G(R)$; then $K(G(R)\phi_B) \sqsubseteq K(\phi_A F(R))$ is K applied to ϕ's own inequation; $\chi := \mathbb{1}_K$ gives $K(\phi) : K\circ G \Rightarrow K\circ F$, K composed on the outside; $\phi := \mathbb{1}_G$ gives $\chi G : L\circ G \Rightarrow K\circ G$, G composed on the inside two candidates, $\chi_{GA}K(\phi_A)$ and $L(\phi_A)\chi_{FA}$, ordered by $\sqsubseteq$; this row is the first	
3 union $\phi \cup \psi$	 $G(R)\phi_B \sqsubseteq \phi_A F(R)$ and $G(R)\psi_B \sqsubseteq \psi_A F(R)$ give $G(R)(\phi_B \cup \psi_B) \sqsubseteq (\phi_A \cup \psi_A)F(R)$	
4 a relator K $K(\phi)$	 $G(R)\phi_B \sqsubseteq \phi_A F(R)$ gives $K(G(R))K(\phi_B) \sqsubseteq K(\phi_A)K(F(R))$	
5 product $\phi \times \psi$	 $\phi : G \Rightarrow F$ and $\psi : G' \Rightarrow F'$ give $\phi \times \psi : G \times G' \Rightarrow F \times F'$ $(R \times S)(U \times V) = (RU) \times (SV)$ and monotonicity in both slots — the row above's K applied to the inequation, run on a bifunctor; the fork is the derived case $\langle \phi, \psi \rangle = \triangleleft(\phi \times \psi)$, and its ONE cost is the lax copy law $R \triangleleft \triangleleft (R \times R)$ — (0.4a)	
6 coproduct $\phi + \psi$	 $\phi : G \Rightarrow F$ and $\psi : G' \Rightarrow F'$ give $\phi + \psi : G + G' \Rightarrow F + F'$ $(R + S)(U + V) = (RU) + (SV)$ and monotonicity in both slots; the co-fork is the derived case $[\phi, \psi] = (\phi + \psi) \triangleright$, and $\triangleright$ costs nothing the step it would need is $(\phi_A F(R)) \sqcap (\psi_A F(R)) \sqsubseteq (\phi \sqcap \psi)_A F(R)$, the wrong direction of (0.6a) a counterexample: (5.7.1a)	
7 meet — fails $\phi \sqcap \psi$		
8 Each row is the general slide $R_A X \sqsubseteq X' R_B$ and $R_B Y \sqsubseteq Y' R_C$ give $R_A(XY) \sqsubseteq (X'Y')R_C$ at $R_A, R_B := G(R), F(R)$ and $X, X' := \phi_B, \phi_A$, quantified over R ; at $X' = X$ with R_A, R_B endorelations it is the monotonicity reading instead — the $\forall R$ sits outside the law, and is the only difference		

§5.7.1. meet is not closed in LaT

counterexample — $\varphi, \psi : \mathbf{G} \Rightarrow \mathbf{F}$ lax natural does NOT give $\varphi \cap \psi$ lax natural

(5.7.1a)




§5.7.2. Two stacked towers

(5.7.2a)

	lower — inside ONE allegory	upper — between allegories
1 0-cell	an object A	an allegory $\mathcal{C}$ — a whole lower tower
2 1-cell	a relation $R : A \rightarrow B$	a relator $F : \mathcal{C} \rightarrow \mathcal{D}$
3 2-cell	$R \sqsubseteq S$	a LaT $\varphi : \mathbf{G} \Rightarrow \mathbf{F} \triangleq \text{lax}$
4 order	$R \sqsubseteq S$	$\varphi \sqsubseteq \psi$ componentwise
5 union	$R \cup S$	$\varphi \cup \psi$, survives $G(R) (\varphi_B \cup \psi_B) \sqsubseteq (\varphi_A \cup \psi_A) F(R)$
6 product	$R \times S \triangleq \times$	$\varphi \times \psi$, survives; a TENSOR, not a categorical product; SYMMETRIC (5.2a) $(G(R) \times G'(R)) (\varphi_B \times \psi_B) \sqsubseteq (\varphi_A \times \psi_A) (F(R) \times F'(R))$
7 coproduct	$R + S$	$\varphi + \psi$, survives; a BIPRODUCT $(G(R) + G'(R)) (\varphi_B + \psi_B) \sqsubseteq (\varphi_A + \psi_A) (F(R) + F'(R))$
8 meet	$R \cap S$	$\varphi \cap \psi$ componentwise, fails (5.7.1a)
9 converse	R°	φ° , fails, oplax: $\varphi_A^\circ G(R) \sqsubseteq F(R) \varphi_B^\circ$
10 zero object	z with $1 \ z = 0$	the constant relator at such a z , initial and terminal

§5.7.3. RES is a special case of LaT

(5.7.3a)



$G \triangleq \text{const } A, F \triangleq \text{const } B$ — the relators $X \rightarrow A$ and $X \rightarrow B$, each sending every arrow to $1 \triangleq \text{lax}$

with $\mathcal{C}$ two objects and one non-identity arrow $X \rightarrow Y$, a LaT $\text{const } A \Rightarrow \text{const } B$ is the pair $\varphi_X, \varphi_Y : A \rightarrow B$ with $\varphi_Y \sqsubseteq \varphi_X$ — the datum $R \sqsubseteq S$

not every LaT is constant: $\exists : \mathbf{P} \Rightarrow \text{Id}$ (5.4d.1) $\pi_2 : (A \times -) \Rightarrow \text{Id}$ (5.6.1c) $\triangleleft : \text{Id} \Rightarrow \Delta$ (0.4a)