

Relation Algebra

§16. Greedy Algorithms



§16.1. Theory

(16.1a)

h, T, R, H, M as in $\triangleq H$; additionally Q a **connected** preorder on the sets $\frac{T^\circ}{\Xi}$ returns, so that $\frac{T^\circ}{\Xi} \text{ est}(Q)$ is entire.

$$\text{PreservesRecip}(F) \wedge \text{Map}(h) \wedge \text{MonotonicAlg}(h, R) \wedge RR \subseteq R \wedge QF(H)h \subseteq F(H)hR \Rightarrow (\mu X : \frac{T^\circ}{\Xi} \text{ est}(Q)F(X)h) \subseteq \frac{H}{\Xi} \text{ est}(R) \quad (16.1b)$$

the same optimum reached by keeping ONE decomposition at each step, so that no set is ever carried and the recursion runs on values alone



the problem to be solved, $H \triangleq (T)^\circ (h) \triangleq H, Q$

Proposition 10.1 at $T=[V_1, V_2]$, $h=[U_1, U_2]$, $Q=Q_1+Q_2$, $V_2V_1^\circ=\emptyset$

§16.2. The detab-entab problem

(16.2a)

$\text{detab} \triangleq ([\text{nil}, \text{expand}]) : \text{String} \rightarrow \text{String}$ over snoc-lists, $\alpha \triangleq [\text{nil}, \text{snoc}]$, $H = \text{detab}^\circ$; $\text{expand } (xs, a) = (a = \text{TB} \rightarrow \text{fill } xs, xs \# [a])$, $\text{fill } xs = xs \# \text{blanks } (n - (\text{col } xs) \bmod n)$.

$\text{col} \triangleq ([\text{zero}, \text{count}])$, $\text{count } (c, a) = (a = \text{NL} \rightarrow 0, c+1)$; TB the tab, BL the blank, NL the newline, tab stops every n columns.


$R \triangleq \text{length} \leq \text{length}^\circ$, U the preorder with $a \cup b \iff a = \text{TB} \vee a = b$, $V \triangleq \text{prefix}^\circ n (\text{fill } \text{fill}^\circ)$, $Q \triangleq 1 + (V \times U)$.

$\text{unfill } xs$ the shortest prefix of xs with $\text{fill } (\text{unfill } xs) = \text{fill } xs$; tbc the trailing blank count, $\text{triple} \triangleq \{\text{unfill } \text{entab}, \{\text{tbc}, \text{col}\}\}$.

$$0 < n \wedge \text{blank} \neq \text{nl} \Rightarrow (\mu X : \frac{[\text{nil}, \text{expand}]^\circ}{3} \text{est}(Q)F(X)[\text{nil}, \text{snoc}]) \sqsubseteq \frac{\text{detab}^\circ}{3} \text{est}(R), \text{ entab} = \text{triple} \quad (16.2b)$$


$$\text{assocl } \pi_1 (1 \times \text{blanks}) \text{ cat}$$

the shortest input **detab** expands to the given output is one pass along that output, holding each blank back and cashing the held blanks in for a tab wherever the column reaches a tab stop



$$[\text{Char}] \xrightarrow{\frac{1}{3}} E(\text{detab}^\circ) E[\text{Char}] \xrightarrow{\text{est}(R)} [\text{Char}]$$

the specification $\triangleq \text{detab}; \text{detab} \text{ entab} = 1$ and nothing shorter does



at $Q \triangleq 1 + (V \times U)$: one character of input is decided at each step. $F(T, R) \alpha \sqsubseteq \alpha R$. **detab** prefix $\sqsubseteq R^\circ$ **detab** is FALSE, — at $n=8$, **detab** $[a, b, c, d, e, \text{TB}] = [a, b, c, d, e, \text{BL}, \text{BL}, \text{BL}]$, whose prefix $[a, b, c, d, e, \text{BL}, \text{BL}]$ is longer than any input giving it, and **detab** $V^\circ \sqsubseteq R^\circ$ **detab** holds. $\text{expand } V^\circ \sqsubseteq \text{expand } U (\pi_1 V^\circ)$ — shortening the output either leaves the last step alone or discards it

$$[\text{Char}] \xrightarrow{\frac{1}{3}} E[\text{Char}] \xrightarrow{E(\text{expand}^\circ)} E([\text{Char}] \times \text{Char}) \xrightarrow{\text{est}(V \times U)} X \xrightarrow{\text{snoc}} [\text{Char}]$$

Proposition 10.1: **nil** and **expand** have disjoint ranges. The greedy step is to emit a tab whenever a tab is legal, consuming all the blanks back to the previous tab stop

$$= \text{entab } xs = \text{entab } (\text{unfill } xs) \# \text{blanks } (\text{tbc } xs) \quad (10.1)$$

what makes $\text{triple} \triangleq (\text{unfill } \text{entab}, \langle \text{tbc}, \text{col} \rangle)$ a snoc-list reduce: the output splits at the last tab stop

$$= \text{triple} = ([\text{base}, \text{op}]), \text{ entab} = \text{triple} \text{ assocl } \pi_1 (1 \times \text{blanks}) \text{ cat}$$

$$\text{base returns } ([], (0, 0)), \text{ op } ((xs, (t, c)), a) =$$

$$(a = \text{BL} \wedge (c+1) \bmod n \neq 0 \rightarrow (xs, (t+1, c+1)), a = \text{BL} \rightarrow (xs \# [\text{TB}], (0, c+1)),$$

$$a = \text{NL} \rightarrow (xs \# \text{blanks } t \# [\text{NL}], (0, 0)), (xs \# \text{blanks } t \# [a], (0, c+1)))$$

the program: one pass carrying the column and the count of pending blanks

§16.3. The minimum tardiness problem

(16.3a)

$FX = 1 + (X \times \text{Job})$, $\alpha \triangleq [\text{nil}, \text{snoc}]$ on schedules, $\beta \triangleq [\text{nil}, \text{snag}]$ on bags, snag putting a job into a bag; $\text{bagify} \triangleq \langle \beta \rangle : [\text{Job}] \rightarrow \text{Bag Job}$, $H = \text{bagify}^\circ$.

$\text{ct}, \text{dt}, \text{wt} : \text{Job} \rightarrow \text{Real}$ the completion, due and weighting quantities of a job; $\text{penalty}(xs, j) = (\text{sum}(\text{list}(\text{ct}(xs)) + \text{ct}(j) - \text{dt}(j)) \times \text{wt}(j))$.

$\text{cost} \triangleq \frac{\text{prefix}}{3} P(\alpha^\circ [\text{zero}, \text{penalty}]) \text{est}(\geq)$, $\text{cost } [] = 0$, $\text{cost } (xs \# [j]) = \text{bmax}(\text{cost } xs, \text{penalty}(xs, j))$, $R \triangleq \text{cost} \leq \text{cost}^\circ$.


$\text{perm} \triangleq \text{bagify} \text{ bagify}^\circ = ([\text{nil}, \text{add}])$, $\text{add}(xs, j) = ys \# [j] \# zs$ for some $xs = ys \# zs$.

$k \triangleq [\text{zero}, \text{assocr } (1 \times ((\text{bagify}^\circ \times 1) \text{ penalty})) \text{ bmax}]$, $f \triangleq [\text{zero}, (\text{bagify}^\circ \times 1) \text{ penalty}]$, $Q \triangleq f \leq f^\circ$, $Q' \triangleq (\text{bagify}^\circ \times 1) \text{ penalty} \leq \text{penalty}^\circ (\text{bagify} \times 1)$.

$$(\forall j. 0 \leq \text{ct}(j)) \wedge (\forall j. 0 \leq \text{wt}(j)) \Rightarrow (\mu X : \frac{[\text{nil}, \text{snag}]^\circ}{3} \text{est}(Q) F(X) [\text{nil}, \text{snoc}] \sqsubseteq \frac{\text{bagify}^\circ}{3} \text{est}(R) , \quad (16.3b)$$


schedule = (null → nil, pick (schedule × 1) snoc)

an ordering of the given bag with least maximum penalty is got by taking a job of least penalty out of the bag, putting it last, and scheduling what is left the same way



$$\text{bag}(\text{Job}) \xrightarrow{\frac{1}{3}} E(\text{bag}(\text{Job})) \xrightarrow{E(\text{bagify}^\circ)} E[\text{Job}] \xrightarrow{\text{est}(R)} [\text{Job}]$$

the specification $\triangleq \text{bagify}$



No greedy **reduce** exists — one would also solve every prefix of the input, and the best schedule of a prefix need not extend to a best schedule of the whole

$$\text{bag}(\text{Job}) \xrightarrow{\frac{1}{3}} E(\text{bag}(\text{Job})) \xrightarrow{E(\text{snag}^\circ)} E(\text{bag}(\text{Job}) \times \text{Job}) \xrightarrow{\text{est}(Q')} X \xrightarrow{\text{snoc}} [\text{Job}]$$

Proposition 10.1: **nil** and **snag** have disjoint ranges

$$\text{pick} \sqsubseteq \frac{\text{snag}^\circ}{3} \text{est}(Q'), \text{ a partial function, quadratic in the number of jobs}$$

§16.4. The TeX problem

(16.4a)

$\text{intern} \triangleq \text{val round} : \text{Decimal} \rightarrow [0, 2^{16})$, $\text{val} \triangleq ([\text{zero}, \text{shift}])$, $\text{shift}(d, r) = (d + r) / 10$, round r rounds $2^{16}r$ to the nearest integer: $\text{round } r = n \iff 2n - 1 < 2^{17}r < 2n + 1$.


$\text{interval } n = ((2n - 1) / 2^{17}, (2n + 1) / 2^{17})$, $r \text{ inrange } (a, b) \iff a < r < b$, $\text{round}^\circ = \text{interval inrange}$, $R \triangleq \text{length} \leq \text{length}^\circ$.

Interval all pairs (a, b) , $\text{Legal}(a, b) \iff 0 < b < 1$ and $a < b$ (10.9), preserved by $[\text{arb}, \text{step}]$ — arb_legal , step_legal
 $[\text{arb}, \text{step}] : 1 + (\text{Digit} \times \text{Interval}) \rightarrow \text{Interval}$, $\text{step}(d, (a, b)) = ((d + a) / 10, (d + b) / 10)$.

$\text{FX} = 1 + (\text{Digit} \times X)$, $\alpha \triangleq [\text{nil}, \text{cons}]$, $H \triangleq ([\text{arb}, \text{step}])^\circ$, $! : \text{Digit} \times \text{Interval} \rightarrow 1$, $Q \triangleq (l^\circ !^\circ r) \cup 1$ l, r are coprod-laws's injections into $\text{FX} = 1 + (\text{Digit} \times X)$, so $l : 1 \rightarrow F(\text{Interval})$ and $r : \text{Digit} \times \text{Interval} \rightarrow F(\text{Interval})$, $w \triangleq 2^{17}$.

$$\text{interval}(\mu X : \frac{[\text{arb}, \text{step}]^\circ}{3} \text{est}(Q)F(X)\alpha) \sqsubseteq \frac{\text{intern}^\circ}{3} \text{est}(R) , \quad \text{extern}(n)=f(2n-1, 2n+1) \quad (16.4b)$$

a shortest decimal whose internal representation is the given multiple of 2^{-16} is got by emitting the one digit the interval of admissible reals allows, until that interval contains zero and the empty decimal will do



$$\begin{aligned}
 & [0, 2^{16}) \xrightarrow{\frac{1}{3}} E([0, 2^{16})) \xrightarrow{E(\text{intern}^\circ)} E[\text{Digit}] \xrightarrow{\text{est}(R)} [\text{Digit}] && \text{the specification } \triangleq \text{intern} \\
 = & [0, 2^{16}) \xrightarrow{\text{interval}} \text{Interval} \xrightarrow{\frac{1}{3}} E(\text{Interval}) \xrightarrow{E(\text{inrange val}^\circ)} E[\text{Digit}] \xrightarrow{\text{est}(R)} [\text{Digit}] \\
 & \text{round}^\circ \text{ is not a map, but interval is, so it comes out of the transpose} \\
 = & [0, 2^{16}) \xrightarrow{\text{interval}} \text{Interval} \xrightarrow{\frac{1}{3}} E(\text{Interval}) \xrightarrow{E(H)} E[\text{Digit}] \xrightarrow{\text{est}(R)} [\text{Digit}] \\
 & \text{fusion: val inrange}^\circ = ([\text{arb}, \text{step}])^\circ \text{ — the converse of val, cut down to intervals, is a reduce on cons-lists} \\
 \sqsubseteq & [0, 2^{16}) \xrightarrow{\text{interval}} \text{Interval} \xrightarrow{\frac{1}{3}} E(\text{Interval}) \xrightarrow{E([\text{arb}, \text{step}]^\circ)} EF(\text{Interval}) \xrightarrow{\text{est}(Q)} F(H) \xrightarrow{\alpha} [\text{Digit}] \\
 & \frac{[\text{arb}, \text{step}]^\circ}{3} \text{ returns at most two elements — stop, or take one more digit — and ! nil} \sqsubseteq \text{cons } R^\circ \text{ makes it stop whenever stopping is legal}
 \end{aligned}$$

$$= \text{extern} = \text{interval } f, \quad f(a, b) = (a < 0 \rightarrow [], [d] \# f(10a - d, 10b - d))$$

the program, with d the digit above

$$= \text{extern}(n) = f(2n-1, 2n+1), \quad f(p, q) = (p \leq 0 \rightarrow [], [d] \# f(10p - w \cdot d, 10q - w \cdot d))$$

$d = (10q) \text{ div } w$: the same in integer arithmetic only, as chapter 3 required of `intern` — every interval reached is $(p/w, q/w)$

§17. Appendix

§17.1. $P(S) \text{ est}(R) = (\exists S) n(\epsilon \setminus (SR^\circ))$

(17.1a)

(p, q) tabulates $W \triangleq (\exists S) n(\epsilon \setminus (SR^\circ))$, and $y \triangleq \frac{(p \exists S) n(qR)}{\exists}$, a map.

(17.1b)

$$\begin{array}{c} \boxed{\begin{array}{c} W \\ P(S) \text{ est}(R) \end{array}} \xleftarrow{p^\circ q = W, 1 \sqsubseteq yy^\circ} \begin{array}{c} \boxed{\begin{array}{c} p^\circ y \\ P(S) \end{array}} \\ \boxed{\begin{array}{c} y^\circ q \\ \text{est}(R) \end{array}} \end{array}$$

(17.1c)

$$\begin{array}{c} \boxed{\begin{array}{c} y^\circ q \\ \exists \end{array}} \xleftarrow{f^\circ \cdot \neg f^\circ, \cdot \exists \neg \exists} \boxed{\begin{array}{c} q \\ (p \exists S) n(qR) \end{array}} \xleftarrow{\Delta \neg n, f^\circ \cdot \neg f^\circ} \begin{array}{c} \boxed{\begin{array}{c} p^\circ q \\ \exists S \end{array}} \\ \boxed{\begin{array}{c} 1 \\ R \end{array}} \end{array}$$

R reflexive

$$\boxed{\begin{array}{c} \epsilon y^\circ q \\ R^\circ \end{array}} \xleftarrow{\cdot \exists \neg \exists, \circ, \text{meet}} \boxed{\begin{array}{c} R^\circ q^\circ q \\ R^\circ \end{array}} \quad \text{q simple}$$

(17.1d)

$$\begin{array}{c} \boxed{\begin{array}{c} p^\circ y \\ P(S) \end{array}} \xleftarrow{\Delta \neg n, \cdot T \neg / T, \circ, f^\circ \cdot \neg f^\circ} \begin{array}{c} \boxed{\begin{array}{c} p^\circ y \exists \\ \exists S \end{array}} \\ \boxed{\begin{array}{c} p \exists \\ y \exists S^\circ \end{array}} \end{array}$$

$$\boxed{\begin{array}{c} p^\circ y \exists \\ \exists S \end{array}} \xleftarrow{\cdot \exists \neg \exists, \text{meet}} \boxed{\begin{array}{c} p^\circ p \exists S \\ \exists S \end{array}} \quad \text{p simple}$$

$$\boxed{\begin{array}{c} p \exists \\ y \exists S^\circ \end{array}} \xleftarrow{\text{modular law}} \boxed{\begin{array}{c} p \exists \\ (p \exists) n(qRS^\circ) \end{array}} \xleftarrow{1 \sqsubseteq qq^\circ} \boxed{\begin{array}{c} q^\circ p \exists \\ RS^\circ \end{array}} \xleftarrow{\circ, T \cdot \neg T \setminus} \boxed{\begin{array}{c} W \\ \epsilon \setminus (SR^\circ) \end{array}} \quad \text{W's right half}$$

§17.2. $P(\text{est}(R)) \text{ est}(R) = P(\text{dom}(\text{est}(R))) \text{ union est}(R)$

(17.2a)

$$\begin{array}{c} \boxed{P(\text{est}(R)) \text{ est}(R)} \xrightarrow{\text{dom}(\text{est}(R)) \text{ est}(R) = \text{est}(R)} \boxed{P(\text{dom}(\text{est}(R)) \text{ est}(R)) \text{ est}(R)} \\ \xrightarrow{P \text{ a relator}} \boxed{P(\text{dom}(\text{est}(R))) P(\text{est}(R)) \text{ est}(R)} \xrightarrow{\text{est-711}} \boxed{P(\text{dom}(\text{est}(R))) \text{ union est}(R)} \end{array}$$

(17.2b)

$$\begin{array}{c} \boxed{\begin{array}{c} P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ P(\text{est}(R)) \text{ est}(R) \end{array}} \xleftarrow{(17.1b) \text{ at } S := \text{est}(R), \Delta \neg n, T \cdot \neg T \setminus} \boxed{\begin{array}{c} P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ \exists \text{est}(R) \end{array}} \\ \boxed{\begin{array}{c} \epsilon P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ \text{est}(R) R^\circ \end{array}} \xleftarrow{\text{count of } T \cdot \neg T \setminus} \boxed{\begin{array}{c} \exists (\exists n(\epsilon \setminus (\epsilon \setminus (SR^\circ)))) \\ \exists (\exists n(\epsilon \setminus (SR^\circ))) \end{array}} \\ \xleftarrow{\text{modular law}} \boxed{\begin{array}{c} (\exists \exists) n(\epsilon \setminus (\epsilon \setminus (SR^\circ))) \\ \exists \text{est}(R) \end{array}} \xleftarrow{T(UnV) \sqsubseteq TUnTV, \cdot \exists \neg \exists, \text{est-73}} \boxed{\begin{array}{c} P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ \exists \text{est}(R) \end{array}} \\ \xleftarrow{P(\text{dom}(\text{est}(R))) \sqsubseteq 1} \boxed{\begin{array}{c} \text{union est}(R) \\ \exists \text{est}(R) \end{array}} \\ \xleftarrow{\epsilon \text{ lax natural}} \boxed{\begin{array}{c} \epsilon P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ \text{est}(R) R^\circ \end{array}} \xleftarrow{\text{dom}(\text{est}(R)) \epsilon \text{ union est}(R)} \boxed{\begin{array}{c} \text{dom}(\text{est}(R)) \epsilon \text{ union est}(R) \\ \text{est}(R) R^\circ \end{array}} \\ \xleftarrow{\text{est}(R) \epsilon \text{est}(R)} \boxed{\begin{array}{c} \text{est}(R) \text{ est}(R)^\circ \epsilon \text{ union est}(R) \\ \text{est}(R) R^\circ \end{array}} \xleftarrow{\text{dom}(\text{est}(R)) \sqsubseteq \text{est}(R) \text{ est}(R)^\circ} \boxed{\begin{array}{c} \text{est}(R) \epsilon \text{est}(R) \\ \text{est}(R) R^\circ \end{array}} \quad \text{UP of est}$$

§17.3. Shortest paths on a cylinder, on lists

(17.3a)

definition	type	note
$F(A, X) = A + A \times X$	$\mathcal{A} \rightarrow \mathcal{A}$	The base functor: one square of the new column, alone or in front of the path so far.
$L = \text{list}^+$	$\mathcal{A} \rightarrow \mathcal{A}$	A path is a non-empty list of squares, one per column crossed.
α the initial algebra	$F(\text{dCL}(L, E)) \rightarrow \text{dCL}(L, E)$	$\alpha(5) = [5]$ and $\alpha(1, [5]) = [1, 5]$: a path is started by one square, or extended by one.
N the n-tuple relator	$\mathcal{A} \rightarrow \mathcal{A}$	One component per row of a column, so the fold carries n answers at once.
$R \triangleq \text{sum} \leq \text{sum}^\circ$	$[\text{Nat}] \rightarrow [\text{Nat}]$	The cost of a path, which the cheapest minimises.
setify	$A[n] \rightarrow EA$	$\text{setify}(1, 2, 3, 4) = \{1, 2, 3, 4\}$ — which row a component came from is forgotten.
moves	$A[n] \rightarrow E(A[n])$	$\text{moves}(x) = \{\text{up}(x), x, \text{down}(x)\}$ — rotated up, unrotated, rotated down.
trans	$E(A[n]) \rightarrow (E A)[n]$	$\text{trans}\{(a, b, c), (x, y, z)\} = (\{a, x\}, \{b, y\}, \{c, z\})$ — component k of the result is the set of the k-th components.
zip	$F(a[n]) \rightarrow (F a)[n]$	$\text{zip}((1, 2, 3, 4), (\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\})) = ((1, \{[5]\}), (2, \{[6]\}), (3, \{[7]\}), (4, \{[8]\}))$.
$\text{cp} \triangleq \frac{F(1, \exists)}{\exists}$	$F(EA) \rightarrow EFA$	$\text{cp}(1, \{[5], [6], [8]\}) = \{(1, [5]), (1, [6]), (1, [8])\}$.
$\text{gen} \triangleq F(1, \text{moves trans } N(\text{union})) \text{zip } N(\text{cp}(F(A, -), T)E(\alpha))$	$F(NA, N(E(TA))) \rightarrow N(E(TA))$	$\text{gen}((1, 2, 3, 4), (\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\}))$ is worked out in (17.3.1b).
$\text{paths} \triangleq (\text{gen}) \text{setify union}$	$T(NA) \rightarrow E(TA)$	$\text{paths}[(1, 2, 3, 4), (5, 6, 7, 8)]$ is the union of (17.3.1b)'s four sets: 12 paths, 3 from each entry row.
the specification $\text{paths est}(R)$	$[\text{Nat}[n]] \rightarrow [\text{Nat}]$	A cheapest path from the entry side to the exit side.

§17.3.1. $N(E(L A))$

$$\begin{array}{l} 1 \quad 5 \\ 2 \quad 6 \\ 3 \quad 7 \\ 4 \quad 8 \end{array} \quad [(1, 2, 3, 4), (5, 6, 7, 8)] : L \ N \ \text{Nat}$$

(17.3.1a)

$$(\text{gen})[(5, 6, 7, 8)] = (\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\})$$

(17.3.1b)

$$\begin{aligned} \text{gen}((1, 2, 3, 4), (\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\})) = \\ (\{ [1, 5], [1, 6], [1, 8] \}, \\ \{ [2, 5], [2, 6], [2, 7] \}, \\ \{ [3, 6], [3, 7], [3, 8] \}, \\ \{ [4, 5], [4, 7], [4, 8] \}) \end{aligned}$$


(17.3.1c)

N	the 4-tuple, one component per row of the entry column
E	the set of paths that can start in that row
L	a path is the non-empty list of the squares it crosses


row 1's set has no $[1,7]$ — from row 1 only rows 4, 1 and 2 are reachable — and it does have $[1,8]$, because the cylinder glues the bottom row to the top. That is **moves trans N(union)** of $\triangleq$ **paths**: component k collects the paths of rows $k-1, k, k+1$.

§17.3.2. The cross product $cp \triangleq \frac{F(1, \exists)}{\exists}$


(17.3.2a)



cp on all of $F(A, E(L A))$



the $A \times -$ summand: $\exists$ picks one path, $1\% \exists$ collects the results



the A summand, no E to distribute: $1\% \exists$ alone, $a \mapsto \{a\}$

(17.3.2b)


type	note
$F(A, -) = A + A \times -$	one wire carries the whole functor, both summands with it; the $\exists$ under it is $F(1, \exists)$
$\exists : E(L A) \rightarrow L A$	the $\exists$ inside $F(1, \exists)$: a set of paths, one of them — $L A$ is one path, A one square
$F(1, \exists) = 1 + 1 \times \exists$	$: A + A \times E(L A) \rightarrow A + A \times L A$, the relator acting on each summand
$\exists : E(A + A \times L A) \rightarrow A + A \times L A$	the $\exists$ under the bar: a set of cells, one of them — a different $\exists$
$cp : A + A \times E(L A) \rightarrow E(A + A \times L A)$	$\frac{R}{\exists}$ turns $R : X \rightarrow Y$ into $X \rightarrow E Y$
$cp = \frac{1}{\exists} E(F(1, \exists))$	what the picture draws: the unit makes the outer E , leaving one $\exists$ — pow-laws

(17.3.2c)

$w = (1, \{[5], [6], [8]\}) : A \times E(L A)$ the right summand
 $1 \times \exists : A \times E(L A) \rightarrow A \times L A$ 1 keeps the square 1, $\exists$ picks one path
 $w \mapsto (1, [5]), (1, [6]), (1, [8]) : A \times L A$ one output per path in w
 $cp(w) = \{(1, [5]), (1, [6]), (1, [8])\} : E(A + A \times L A)$ the three of them, collected

§17.3.3. $gen = F(1, \text{moves trans } N(\text{union})) \text{ zip } N(cp P(\alpha))$

(17.3.3a)




(17.3.3b)

$u = ((1,2,3,4), (\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\}))$: $F(N A, N(E(L A)))$
 $F(1, \text{moves trans } N(\text{union}))$: 1 keeps the column, the path SETS move
 $\text{moves}(\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\})$
 $= (\{[6]\}, \{[7]\}, \{[8]\}, \{[5]\}),$
 $(\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\}),$
 $(\{[8]\}, \{[5]\}, \{[6]\}, \{[7]\})$: $E(N(E(L A)))$ down, unmoved, up
 $\text{trans}(\text{that})$
 $= (\{[6]\}, \{[5]\}, \{[8]\}, \{[7]\}, \{[6]\}, \{[5]\}),$
 $\{[8]\}, \{[7]\}, \{[6]\}, \{[5]\}, \{[8]\}, \{[7]\})$: $N(E(E(L A)))$ row k gets rows k-1, k, k+1
 $N(\text{union})(\text{that})$
 $= (\{[5], [6], [8]\}, \{[5], [6], [7]\},$
 $\{[6], [7], [8]\}, \{[5], [7], [8]\})$: $N(E(L A))$ every path into row k, none dropped
 $\text{zip}(\text{that})$
 $= ((1, \{[5], [6], [8]\}), (2, \{[5], [6], [7]\}),$
 $(3, \{[6], [7], [8]\}), (4, \{[5], [7], [8]\}))$: $N(F(A, E(L A)))$ each row: its square, and the paths it may be put in front of
 $N(\text{cp } P(\alpha))(\text{that})$
 $= (\{[1,5], [1,6], [1,8]\}, \{[2,5], [2,6], [2,7]\},$
 $\{[3,6], [3,7], [3,8]\}, \{[4,5], [4,7], [4,8]\})$: $N(E(L A))$ cp pairs the square with each path, α prefixes it: $\alpha(1, [5]) = [1,5]$

§17.3.4. gen is an F-algebra; $(\text{gen}): \alpha(\text{gen}) = F(1, (\text{gen}))\text{gen}$

(17.3.4a)



α puts the column back on the list, then the fold reads all of it

the fold reads the rest under F , then one gen puts the column in front

(17.3.4b)

$$xs = [(1,2,3,4), (5,6,7,8)] = \alpha((1,2,3,4), [(5,6,7,8)]) : L(N A)$$

$$(\text{gen})(xs) = \text{gen}((1,2,3,4), (\text{gen})[(5,6,7,8)])$$

both halves are in (17.3.1b): the tail folds to $(\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\})$, one path per row and each of them one square long, and gen on it is (17.3.3b)'s walk.

§17.3.5. $Q : F(N A, N(L A)) \rightarrow N(L A)$, an $F(N A, -)$ -algebra: $Q = F(1, \text{moves trans } N(\text{est}(R))) \text{ zip } N(\alpha)$

(17.3.5a)

$Q : F(N A, N(L A)) \rightarrow N(L A)$: the fold's algebra: a new column, one path per row
 $u = ((1,2,3,4), ([5], [6], [7], [8]))$: $F(N A, N(L A))$
 $F(1, \text{moves trans } N(\text{est}(R)))$: 1 keeps the column, the paths move
 $\text{moves}([5], [6], [7], [8])$
 $= ([6], [7], [8], [5]),$
 $([5], [6], [7], [8]),$
 $([8], [5], [6], [7])$: $E(N(L A))$ down, unmoved, up
 $\text{trans}(\text{that})$
 $= ([6], [5], [8]), ([7], [6], [5]),$
 $[8], [7], [6]), ([5], [8], [7])$: $N(E(L A))$ row k gets rows k-1, k, k+1
 $N(\text{est}(R))(\text{that})$
 $= ([5], [5], [6], [5])$: $N(L A)$ the cheapest into each row – chosen BEFORE the new square
 $\text{zip}(\text{that})$
 $= ((1, [5]), (2, [5]), (3, [6]), (4, [5]))$: $N(F(A, L A))$ each row: its square, and the one predecessor that survived
 $N(\alpha)(\text{that})$
 $= ([1,5], [2,5], [3,6], [4,5])$: $N(L A)$ $\alpha(1, [5]) = [1,5]$

$Q = [N(\text{wrap}), (1 \times \text{moves trans } N(\text{est}(R))) \text{ zip}' N(\text{cons})]$ (17.3.5b)	
at the last column Q starts one path per row, and at each earlier one it puts each square in front of the cheapest of the three kept paths it can step to — the algebra (17.3.5c)'s last step folds	
circuit — the fork is $F(NA, N(LA)) = NA + NA \times N(LA)$	Hinze–Marsden
$F(N A) (N (T A)) \text{---} \boxed{Q} \text{---} N(T A)$	
$FN(T A) \text{---} \boxed{F(1, \text{moves trans } N(\text{est}(R)))} \text{---} FN(T A) \text{---} \boxed{\text{zip}} \text{---} N(F A (T A)) \text{---} \boxed{N(\alpha)} \text{---} N(T A)$ the fusion condition (17.3.5d) read as a definition,	
$FN(N A) (N (T A)) \text{---} \left\{ \begin{array}{l} N(p_{\circ}) \text{---} \boxed{N(\text{wrapA})} \\ q \text{---} \boxed{\text{pm}} \text{---} q \text{---} \boxed{\text{zip}'} \text{---} N(q_{\circ}) \text{---} \boxed{N(\text{consA})} \end{array} \right\} \text{---} N(T A)$ $\text{zip} = 1 + \text{zip}', \alpha = [\text{wrap}, \text{cons}]$	

paths $\text{est}(R) \exists (Q)$ setify $\text{est}(R)$ (17.3.5c)	
shortest paths on a cylinder: the dynamic programming approach to exhaustive search allows a path of least cost to be found in $O(n \times m)$ time; Q is (17.3.5b)'s algebra.	
circuit	Hinze–Marsden
$T(N\ A) \text{---} \boxed{\text{paths}} \text{---} E(T\ A) \text{---} \boxed{\text{est}(R)} \text{---} T\ A$	
$= T(N\ A) \text{---} \boxed{F(N\ A)\ (N\ (E\ (T\ A)))} \text{---} \boxed{\text{gen}} \text{---} NE(T\ A) \text{---} \boxed{\text{setify}} \text{---} EE(T\ A) \text{---} \boxed{\text{union}} \text{---} E(T\ A) \text{---} \boxed{\text{est}(R)} \text{---} T\ A$ $\triangleq \text{paths at paths}$	
$\sqsubseteq T(N\ A) \text{---} \boxed{F(N\ A)\ (N\ (E\ (T\ A)))} \text{---} \boxed{\text{gen}} \text{---} NE(T\ A) \text{---} \boxed{\text{setify}} \text{---} EE(T\ A) \text{---} \boxed{P(\text{est}(R))} \text{---} E(T\ A) \text{---} \boxed{\text{est}(R)} \text{---} T\ A$ $P(\text{est}(R))\ \text{est}(R) \sqsubseteq \text{union}\ \text{est}(R) \text{--- a minimum in each set, then a minimum of those; } R \text{ transitive}$	
$\sqsubseteq T(N\ A) \text{---} \boxed{F(N\ A)\ (N\ (E\ (T\ A)))} \text{---} \boxed{\text{gen}} \text{---} NE(T\ A) \text{---} \boxed{N(\text{est}(R))} \text{---} N(T\ A) \text{---} \boxed{\text{setify}} \text{---} E(T\ A) \text{---} \boxed{\text{est}(R)} \text{---} T\ A$ $\text{setify lax natural}$	
$\sqsubseteq T(N\ A) \text{---} \boxed{F(N\ A)\ (N\ (T\ A))} \text{---} \boxed{Q} \text{---} N(T\ A) \text{---} \boxed{\text{setify}} \text{---} E(T\ A) \text{---} \boxed{\text{est}(R)} \text{---} T\ A$ $\text{fusion at (17.3.5d)}$	

gen $N(\text{est}(R)) \exists F(1, N(\text{est}(R))) Q$ (17.3.5d)	
fusion: the condition for fusion in (17.3.5c)'s last step, used to derive a definition of Q .	
circuit	Hinze–Marsden
$F(N\ A)\ (N\ (E\ (T\ A))) \text{---} \boxed{\text{gen}} \text{---} NE(T\ A) \text{---} \boxed{N(\text{est}(R))} \text{---} N(T\ A)$	
$\sqsubseteq FNE(T\ A) \text{---} \boxed{F(1, N(\text{est}(R)))} \text{---} FN(T\ A) \text{---} \boxed{Q} \text{---} N(T\ A)$ $(7.13), \text{ then zip, trans, moves lax natural}$	

(7.13) is $F(1, \text{est}(R)) \alpha \sqsubseteq \text{cp } P(\alpha)\ \text{est}(R)$, mon-thm71 at the map α with $\frac{F(1, \exists)\alpha}{\exists} = \text{cp } P(\alpha)$: extending every path in a set and then taking a minimum is beaten by extending one minimum. It is the crux here, not the greedy theorem.