


Companion to Bird & de Moor, Algebra of Programming


$$\text{Map}(h) \wedge F(R^\circ) \ h \sqsubseteq h \ R^\circ \wedge R^\circ \ R^\circ \sqsubseteq R^\circ \wedge T^\circ F(H) \ h = H \wedge Q F(H) \ h \sqsubseteq F(H) \ h \ R \Rightarrow H^\circ \frac{T^\circ}{\exists} \text{est}(Q) \quad (10.1c)$$

$$F(\frac{H}{\exists} \text{est}(R)) \ h \sqsubseteq R^\circ$$

H° followed by the greedy body at M is $\sqsubseteq R^\circ$: an answer of the body is never worse than an answer of H to the same input




(a)	$H^\circ \frac{T^\circ}{\exists} \text{est}(Q) F(\frac{H}{\exists} \text{est}(R)) \ h$	$T^\circ F(H) \ h = H$ $\Rightarrow H^\circ = h^\circ F(H^\circ) \ T$
(b)	$= h^\circ F(H^\circ) \ T \frac{T^\circ}{\exists} \text{est}(Q) F(\frac{H}{\exists} \text{est}(R)) \ h$	$T \frac{T^\circ}{\exists} \sqsubseteq \epsilon$
(c)	$\sqsubseteq h^\circ F(H^\circ) \ \epsilon \text{est}(Q) F(\frac{H}{\exists} \text{est}(R)) \ h$	$\epsilon \text{est}(R) \sqsubseteq R^\circ$
(d)	$\sqsubseteq h^\circ F(H^\circ) \ Q^\circ F(\frac{H}{\exists} \text{est}(R)) \ h$	$Q F(H) \ h \sqsubseteq F(H) \ h \ R$ $\Rightarrow h^\circ F(H^\circ) \ Q^\circ \sqsubseteq R^\circ \ h^\circ F(H^\circ)$




(a)	$\sqsubseteq R^\circ \ h^\circ F(H^\circ \frac{H}{\exists} \text{est}(R)) \ h$	$S^\circ \frac{S}{\exists} \text{est}(R) \sqsubseteq R^\circ$
(b)	$\sqsubseteq R^\circ \ h^\circ F(R^\circ) \ h$	$\text{Map}(f)$ $\Rightarrow F(R) \ f \sqsubseteq f \ R \Leftrightarrow f^\circ F(R) \ f \sqsubseteq R$
(c)	$\sqsubseteq R^\circ \ R^\circ$	$R^\circ \ R^\circ \sqsubseteq R^\circ$
(d)	$\sqsubseteq R^\circ$	

$$V_2 \ V_1^\circ = \emptyset \implies \frac{[V_1, V_2]^\circ}{\exists} \text{est}(Q_1 + Q_2) [U_1, U_2] = (\text{ran}(V_1) \ \frac{V_1^\circ}{\exists} \text{est}(Q_1) U_1) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{\exists} \text{est}(Q_2) U_2) \quad \text{Proposition 10.1}$$


when V_1 and V_2 have disjoint ranges, the greedy step over $[V_1, V_2]$ runs the V_1 step on inputs V_1 reaches and the V_2 step on inputs V_2 reaches



(a)	$\frac{[V_1, V_2]^\circ}{\exists} \text{est}(Q_1 + Q_2) [U_1, U_2]$	$\frac{[V_1, V_2]^\circ}{\exists} \text{est}(Q_1 + Q_2) [U_1, U_2] = (\text{ran}(V_1) \ \frac{V_1^\circ}{\exists} \text{est}(Q_1 + Q_2) [U_1, U_2]) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{\exists} \text{est}(Q_1 + Q_2) [U_1, U_2])$
(b)	$= (\text{ran}(V_1) \ \frac{V_1^\circ}{\exists} \text{est}(Q_1 + Q_2) [U_1, U_2]) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{\exists} \text{est}(Q_1 + Q_2) [U_1, U_2])$	$V_2 \ V_1^\circ = \emptyset \implies \text{ran}(V_1) \ \frac{[V_1, V_2]^\circ}{\exists} = \text{ran}(V_1) \ \frac{V_1^\circ}{\exists} P(l)$
(c)	$= (\text{ran}(V_1) \ \frac{V_1^\circ}{\exists} P(l) \text{est}(Q_1 + Q_2) [U_1, U_2]) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{\exists} \text{est}(Q_1 + Q_2) [U_1, U_2])$	$V_2 \ V_1^\circ = \emptyset \implies \text{ran}(V_2) \ \frac{[V_1, V_2]^\circ}{\exists} = \text{ran}(V_2) \ \frac{V_2^\circ}{\exists} P(r)$



(a)	$= (\text{ran}(V_1) \ \frac{V_1^\circ}{\exists} P(l) \text{est}(Q_1 + Q_2) [U_1, U_2]) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{\exists} P(r) \text{est}(Q_1 + Q_2) [U_1, U_2])$	$P(l) \text{est}(Q_1 + Q_2) = \text{est}(Q_1) l$
(b)	$= (\text{ran}(V_1) \ \frac{V_1^\circ}{\exists} \text{est}(Q_1) l [U_1, U_2]) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{\exists} P(r) \text{est}(Q_1 + Q_2) [U_1, U_2])$	$P(r) \text{est}(Q_1 + Q_2) = \text{est}(Q_2) r$



(a)	$= (\text{ran}(V_1) \ \frac{V_1^\circ}{\exists} \text{est}(Q_1) l [U_1, U_2]) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{\exists} \text{est}(Q_2) r [U_1, U_2])$	$l [R, S] = R$
(b)	$= (\text{ran}(V_1) \ \frac{V_1^\circ}{\exists} \text{est}(Q_1) U_1) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{\exists} \text{est}(Q_2) r [U_1, U_2])$	$r [R, S] = S$
(c)	$= (\text{ran}(V_1) \ \frac{V_1^\circ}{\exists} \text{est}(Q_1) U_1) \cup (\text{ran}(V_2) \ \frac{V_2^\circ}{\exists} \text{est}(Q_2) U_2)$	


§10.2. The detab-entab problem

(10.2a)


	name	type	definition	meaning
1	detab	$\text{String} \rightarrow \text{String}$	$([\text{nil}, \text{expandFn } n \text{ tb } \text{nl blank}]) = \text{detab}$	replace every tab by blanks up to the next tab stop; $H = \text{detab}^\circ$
2	expand(x,a)	String	$\text{expand}(x,a) \triangleq \text{if } a = \text{tb} \text{ then fill } x \text{ else snoc } x \text{ a}$	a tab fills the line to the next stop; any other character is appended
3	fill(x)	String	$\text{fill}(x) \triangleq x ++ \text{ blanks } (n - \text{col } x \% n)$	append blanks up to the next multiple of n columns
4	col()	$\text{String} \rightarrow \mathbb{N}$	$\text{col}(\text{wrap}(a)) = 0, \text{col}(\text{snoc}(x_1, a)) = \text{if } a = \text{nl} \text{ then } 0 \text{ else } \text{col } x_1 + 1$	the column the string ends at: characters since the last newline
5	R	$[\text{Char}] \rightarrow [\text{Char}]$	$u R v \iff \text{length}(u) \leq \text{length}(v)$	x is no longer than y
6	U	$\text{Char} \rightarrow \text{Char}$	$a U b \iff a = \text{tb } v \text{ a} = b$	a is the tab or a=b
7	V	$[\text{Char}] \rightarrow [\text{Char}]$	$x V y \iff \text{prefix}(x,y) \wedge \text{fill}(x) = \text{fill}(y)$	a prefix that fills to the same string
8	Q	$F[\text{Char}] \rightarrow F[\text{Char}]$	$Q(l(\text{val}), l(\text{val}_1)) = \text{true}, Q(r(p), r(q)) = V(\pi_1(p), \pi_1(q)) \wedge U(\pi_2(p), \pi_2(q)),$ $(\forall \text{val}. \forall \text{val}'. u = l(\text{val}) \implies v = l(\text{val}')) \implies \text{false}) \wedge (\forall p. \forall q. u = r(p) \implies v = r(q) \implies \text{false}) \implies Q(u,v) = \text{false}$	compare two decompositions by V on the strings and U on the characters
9	unfill()	$\text{String} \rightarrow \text{String}$	$\text{unfill}(\text{wrap}(a)) = [],$ $\text{unfill}(\text{snoc}(x_1, a)) = \text{if } a = \text{blank} \wedge \text{col}(\text{snoc } x_1 \text{ a}) \% n \neq 0 \text{ then } \text{unfill } x_1$ $\text{else } \text{snoc } x_1 \text{ a}$	the shortest prefix that fills to the same string
10	tbc()	$\text{String} \rightarrow \mathbb{N}$	$\text{tbc}(\text{wrap}(a)) = 0, \text{tbc}(\text{snoc}(x_1, a)) = \text{if } a = \text{blank} \wedge \text{col}(\text{snoc } x_1 \text{ a}) \% n \neq 0$ $\text{then } \text{tbc } x_1 + 1 \text{ else } 0$	the number of trailing blanks
11	triple	$[\text{Char}] \rightarrow \text{String} \times (\mathbb{N} \times \mathbb{N})$	$\text{triple} \triangleq (\text{unfill } \text{entab}, \{\text{tbc}, \text{col}\})$	what entab keeps: the unfilled output, its trailing blank count and its column

$$0 < n \wedge \text{blank} \neq \text{nl} \implies \text{expand } V^\circ \sqsubseteq \text{expand } u (\pi_1 V^\circ) \quad (10.2b)$$

shortening the output of one **expand** step to a **V**-smaller string either keeps the whole step (**expand**) or drops its character and shortens its input string ($\pi_1 V^\circ$)



(a)	$\text{expand } V^\circ$	$\text{expand} = (\text{istab } \pi_1 \text{ fill}) \cup (\text{nottab } \text{snoc})$
(b)	$= ((\text{istab } \pi_1 \text{ fill}) \cup (\text{nottab } \text{snoc})) V^\circ$	$(S \cup T) R = (S R) \cup (T R)$
(c)	$= (\text{istab } \pi_1 \text{ fill } V^\circ) \cup (\text{nottab } \text{snoc } V^\circ)$	$0 < n \wedge \text{blank} \neq \text{nl} \implies \text{fill } V^\circ = \text{fill}$




(a)	$= (\text{istab } \pi_1 \text{ fill}) \cup (\text{nottab } \text{snoc } V^\circ)$	$0 < n \wedge \text{blank} \neq \text{nl} \implies \text{snoc } V^\circ \sqsubseteq \text{snoc } u (\pi_1 V^\circ)$
(b)	$\sqsubseteq (\text{istab } \pi_1 \text{ fill}) \cup (\text{nottab } (\text{snoc } u (\pi_1 V^\circ)))$	$(\text{istab } \pi_1 \text{ fill}) \cup (\text{nottab } (\text{snoc } u X)) \sqsubseteq \text{expand } u X$
(c)	$\sqsubseteq \text{expand } u (\pi_1 V^\circ)$	

$$0 < n \wedge \text{blank} \neq n\mathbf{l} \Rightarrow \text{detab } V^\circ \sqsubseteq R^\circ \text{ detab} \quad (10.2c)$$

any V -shortening of **detab**'s output is **detab**'s output on an input no longer than the given one



(a)	$\text{detab } V^\circ$	$\text{detab} = [\text{nil}, \text{snoc}]^\circ F(\text{detab}) [\text{nil}, \text{expand}]$
(b)	$= [\text{nil}, \text{snoc}]^\circ F(\text{detab}) [\text{nil}, \text{expand}] V^\circ$	$[R, S] Z = [R Z, S Z]$
(c)	$= [\text{nil}, \text{snoc}]^\circ F(\text{detab}) [\text{nil } V^\circ, \text{expand } V^\circ]$	$\text{nil } V^\circ = \text{nil}$
(d)	$= [\text{nil}, \text{snoc}]^\circ F(\text{detab}) [\text{nil}, \text{expand } V^\circ]$	$(10.2b) \ 0 < n \wedge \text{blank} \neq n\mathbf{l} \Rightarrow \text{expand } V^\circ \sqsubseteq \text{expand } u (\pi_1 V^\circ)$




(a)	$\sqsubseteq [\text{nil}, \text{snoc}]^\circ F(\text{detab}) [\text{nil}, \text{expand } u (\pi_1 V^\circ)]$	$[\text{nil}, \text{snoc}]^\circ F(\text{detab}) [\text{nil}, \text{expand } u X] = \text{detab } u (\text{snoc}^\circ (\text{detab} \times 1) X)$
(b)	$= \text{detab } u (\text{snoc}^\circ (\text{detab} \times 1) \pi_1 V^\circ)$	$\text{snoc}^\circ (\text{detab} \times 1) \pi_1 Y = \text{snoc}^\circ \pi_1 \text{detab } Y$
(c)	$= \text{detab } u (\text{snoc}^\circ \pi_1 \text{detab } V^\circ)$	$0 < n \wedge \text{blank} \neq n\mathbf{l} \Rightarrow \text{detab } u (\text{snoc}^\circ \pi_1 \text{detab } V^\circ) \sqsubseteq R^\circ \text{detab}$
(d)	$\sqsubseteq R^\circ \text{detab}$	

$$0 < n \wedge \text{blank} \neq n\mathbf{l} \Rightarrow (\mu X : \frac{[\text{nil}, \text{expand}]^\circ}{3} \text{est}(Q) F(X) [\text{nil}, \text{snoc}]) \sqsubseteq \frac{\text{detab}^\circ}{3} \text{est}(R) \quad (10.2d)$$

the shortest input **detab** expands to the given output is one pass along that output, holding each blank back and caching the held blanks in for a tab wherever the column reaches a tab stop


the specification — $\text{detab}(\text{wrap}(a)) = []$, $\text{detab}(\text{snoc}(x_1, a)) = \text{expand}(\text{detab}(x_1), a)$; $\text{detab } \text{entab} = 1$ and nothing shorter does



(a)	$\frac{\text{detab}^\circ}{3} \text{est}(R)$	at $Q \triangleq 1 + (V \times U)$: one character of input is decided at each step. $F(\tau, R) \alpha \alpha R$. $\text{detab } \text{prefix} \sqsubseteq R^\circ \text{detab}$ is FALSE, — at $n=8$, $\text{detab} [a, b, c, d, e, \text{TB}] = [a, b, c, d, e, \text{BL}, \text{BL}, \text{BL}]$, whose prefix $[a, b, c, d, e, \text{BL}, \text{BL}]$ is longer than any input giving it, and $\text{detab } V^\circ \sqsubseteq R^\circ \text{detab}$ holds. $\text{expand } V^\circ \sqsubseteq \text{expand } u (\pi_1 V^\circ)$ — shortening the output either leaves the last step alone or discards it
(b)	$\sqsubseteq \frac{[\text{nil}, \text{expand}]^\circ}{3} \text{est}(Q) F(X) [\text{nil}, \text{snoc}]$	Proposition 10.1: nil and expand have disjoint ranges. The greedy step is to emit a tab whenever a tab is legal, consuming all the blanks back to the previous tab stop
(c)	$\sqsubseteq \frac{\text{expand}^\circ}{3} \text{est}(V \times U) (X \times 1) \text{snoc}$	

$$\text{blank} \neq \text{nl} \wedge \text{tb} \neq \text{nl} \Rightarrow \langle \text{detab}, \text{detab col} \rangle = \langle \llbracket \text{base}, \text{step} \rrbracket \rangle \quad (10.2e)$$

one pass over the input carries the output so far together with its column



(a)	$\langle \text{detab}, \text{detab col} \rangle$	both components are folds over the same input; step on $(\text{detab } x, \text{col}(\text{detab } x))$ is $(\text{detab}, \text{col detab})$ one character further — not a tabulated row
(b)	$= \langle \llbracket \text{base}, \text{step} \rrbracket \rangle$	

Exercise 10.1

- $\text{blank} \neq \text{nl} \wedge \text{tb} \neq \text{nl} \Rightarrow \langle \llbracket \text{base}, \text{step} \rrbracket \rangle (\text{convert}(l), r) \Leftrightarrow r = \text{loop}(\text{step}, (\llbracket \cdot, 0 \rrbracket, l))$
the fold of $\llbracket \text{base}, \text{step} \rrbracket$ over the input converted to a snoc-list is the left loop of **step** from **base**

- $\pi_1(\text{loop}(\text{step}, (\llbracket \cdot, c_0 \rrbracket, x))) = \text{loop}'((f, g), (c_0, x))$

Exercise 10.1: when **step** only appends $f(c, a)$ to the output and moves the state by g , the output of the loop is $\text{loop}'(f, g)$, which never carries the output

- $a \neq \text{tb} \Rightarrow \text{expand}(y, b) = \text{snoc}(x, a) \Leftrightarrow \text{fill}(y) = \text{snoc}(x, a) \wedge b = \text{tb} \vee (y, b) = (x, a)$
expand produces $x\#a$ from a tab after a string that fills to it, or from x and a

(10.2f)

- $0 < n \wedge \text{blank} \neq \text{nl} \Rightarrow (\exists y. \text{fill}(y) = \text{snoc}(x, a)) \Leftrightarrow a = \text{blank} \wedge \text{col}(\text{snoc } x \text{ } a) \% n = 0$
a string is a **fill** exactly when it ends in a blank on a tab stop

(10.1)

- $\text{entab}(x) = \text{entab}(\text{unfill } x) ++ \text{blanks}(\text{tbc } x)$

(10.1): the output of **entab** is that of **unfill** x followed by the trailing blanks held back

- $\text{tbc}[] = 0$


the empty string has no trailing blanks

- $\text{tbc}(\text{snoc}(x, a)) = \text{if } a = \text{blank} \wedge \text{col}(\text{snoc } x \text{ } a) \% n \neq 0 \text{ then } \text{tbc } x + 1 \text{ else } 0$

a blank off a tab stop adds one held blank; anything else releases them all

$$\text{blank} \neq \text{nl} \Rightarrow \langle \text{tbc}, \text{col} \rangle = \langle \llbracket \text{base}, \text{op} \rrbracket \rangle \quad (10.2g)$$


one pass counts the held blanks and the column together



(a)	$\langle \text{tbc}, \text{col} \rangle$	op on $(\text{tbc } x, \text{col } x)$ is (tbc, col) one character further, from the equations for tbc — (10.1) $\text{entab}(x) = \text{entab}(\text{unfill } x) ++ \text{blanks}(\text{tbc } x)$, $\text{tbc}[] = 0$, $\text{tbc}(\text{snoc}(x, a)) = \text{if } a = \text{blank} \wedge \text{col}(\text{snoc } x \text{ } a) \% n \neq 0 \text{ then } \text{tbc } x + 1 \text{ else } 0$
(b)	$= \langle \llbracket \text{base}, \text{op} \rrbracket \rangle$	

$$\text{blank} \neq \text{nl} \Rightarrow \text{triple} = \langle \llbracket \text{base}, \text{op} \rrbracket \rangle \quad (10.2h)$$

one pass carries the output up to the held blanks, their count and the column



(a)	triple	op on $\text{triple } x$ is triple one character further: (10.1) where the blanks are released — (10.1) $\text{entab}(x) = \text{entab}(\text{unfill } x) ++ \text{blanks}(\text{tbc } x)$, $\text{tbc}[] = 0$, $\text{tbc}(\text{snoc}(x, a)) = \text{if } a = \text{blank} \wedge \text{col}(\text{snoc } x \text{ } a) \% n \neq 0 \text{ then } \text{tbc } x + 1 \text{ else } 0$
(b)	$= \langle \llbracket \text{base}, \text{op} \rrbracket \rangle$	

$\text{entab} = \text{triple assocl } \pi_1 (1 \times \text{blanks}) \text{ cat}$ (10.2i)		
entab is triple with the held blanks appended to its output		
list Char (a) list Char (b)		
(a)	entab	(10.1) — (10.1) $\text{entab}(x) = \text{entab} (\text{unfill } x) ++ \text{blanks } (\text{tbc } x), \text{tbc}[] = 0, \text{tbc}(\text{snoc}(x,a)) = \text{if } a = \text{blank} \wedge \text{col}(\text{snoc } x \ a) \% n \neq 0 \text{ then } \text{tbc } x + 1 \text{ else } 0$
(b)	$= \text{triple assocl } \pi_1 (1 \times \text{blanks}) \text{ cat}$	


§10.3. The minimum tardiness problem

				(10.3a)
	name	type	definition	meaning
1	bag(Job)	$\mathcal{A}$	$\text{bag}(\text{Job}) \triangleq \text{Quotient}(\text{perm})$	a bag of jobs: a list up to reordering
2	snag(p)	bag(Job)	$\text{snag}(p) \triangleq \pi_2 p :: \pi_1 p$	put one job into a bag
3	bagify	[Job]→bag(Job)	$\langle [\text{nil}, \text{snag}] \rangle = \text{bagify}$	forget the order of a schedule; $H = \text{bagify}^\circ$ lists every schedule of a bag
4	penalty(s,j)	Int	$\text{penalty}(s,j) \triangleq (\text{sum}(\text{list}(\text{ct})) \ s + \text{ct } j - \text{dt } j) \text{wt}(j)$	how late job j finishes after the schedule xs, times its weight
5	cost	[Job] → Int	$\text{cost}(\text{wrap}(a)) = 0, \text{cost}(\text{snoc}(x_1,a)) = \text{bmax}(\text{cost}(x_1), \text{penalty}(x_1,a))$	the largest penalty of any job in the schedule
6	bmax(a,b)	Int	$\text{bmax}(a,b) \triangleq \text{if } a \leq b \text{ then } b \text{ else } a$	the larger of two numbers
7	R	[Job]→[Job]	$R = \text{cost} \leq \text{cost}^\circ$	xs costs no more than ys
8	add	[Job]×Job→[Job]	$p \text{ add } w \iff \text{add}(\pi_2(p), w)$	insert j anywhere in xs; $\text{perm} \triangleq \text{bagify}$ $\text{bagify}^\circ = \langle [\text{nil}, \text{add}] \rangle$
9	k	F(Int×bag(Job))→Int	$k \triangleq [\text{nil}, \text{kStep}]$	the cost of a schedule from the cost of its front and the penalty of its last job
10	f	F bag(Job) → Int	$f(l(\text{val})) = 0, f(r(p)) = (\text{bagify}^\circ \times 1) \text{penalty}(p)$	the penalty of the last job, read off the bag before it
11	Q	F(bag(Job))→F(bag(Job))	$u \ Q \ v \iff f_u \leq f_v$	the last job of u gets no larger penalty than that of v
12	Q'	bag(Job)×Job→bag(Job)×Job	$p \ Q' \ q \iff (\text{bagify}^\circ \times 1) \text{penalty}(p) \leq (\text{bagify}^\circ \times 1) \text{penalty}(q)$	Q on the snag summand: compare the penalties of the two last jobs
13	g	F[Job]→Int	$g \triangleq [\text{zero}, \text{penalty}], g(s,c) \iff c = \text{gs}$	(10.5): the penalty of the last job, zero for the empty schedule
14	m	F[Job]→Int	$m \triangleq [\text{zero}, \pi_1 \text{cost}]$	(10.6), B&dM's h renamed as in $(\forall j. 0 \leq \text{ct}(j)) \wedge (\forall j. 0 \leq \text{wt}(j))$ $\implies (\mu X : \frac{[\text{nil}, \text{snag}]^\circ}{3} \text{est}(Q) \text{F}(X) [\text{nil}, \text{snoc}]) \sqsubseteq \frac{\text{bagify}^\circ}{3}$ $\text{est}(R) : \text{the cost of the schedule before the last job}$

$$[\text{nil}, \text{snoc}] \text{ cost} = F(\langle \text{cost}, \text{bagify} \rangle) k$$

(10.3b)

the cost of a schedule is got from the cost and the bag of the schedule before its last job, and that job




(a)	$[\text{nil}, \text{snoc}] \text{ cost}$	$\text{cost}(xs \# [j]) = \text{bmax}(\text{cost } xs, \text{penalty}(xs, j))$, and $\text{penalty}(xs, j)$ reads only the bag of xs — $\text{penalty}(s, j) \triangleq (\text{sum}(\text{list}(\text{ct})) s + \text{ct } j - \text{dt } j) \text{wt}(j)$
(b)	$= F(\langle \text{cost}, \text{bagify} \rangle) k$	

$$F(\geq \times 1) k \sqsubseteq k \geq$$

Exercise 10.5

a larger cost before the last job gives a larger cost after it




(a)	$F(\geq \times 1) k$	bmax is monotone in its first argument
(b)	$\sqsubseteq k \geq$	

$$F(R^\circ \cap (\text{bagify } \text{bagify}^\circ)) [\text{nil}, \text{snoc}] \sqsubseteq [\text{nil}, \text{snoc}] R^\circ$$

(10.2)

improving the schedule before the last job, among schedules of the same bag, improves the whole schedule




(a)	$F(R^\circ \cap (\text{bagify } \text{bagify}^\circ)) [\text{nil}, \text{snoc}]$	Proposition 9.3 at $S \triangleq \text{bagify}$, $\leq \triangleq$ — Proposition 9.3 $\text{Map}(\text{cost}) \wedge \text{Simple}(S) \wedge R = \text{cost} \leq \text{cost}^\circ \wedge h$ $\text{cost} = F(\langle \text{cost}, S \rangle) k \wedge F(\leq \times 1) k \sqsubseteq k \leq$ $\implies F(R \cap (S \circ^\circ)) h \sqsubseteq h R$, with $[\text{nil}, \text{snoc}] \text{ cost}$ $= F(\langle \text{cost}, \text{bagify} \rangle) k$ and Exercise 10.5 $F(\geq \times 1) k$ $\sqsubseteq k \geq$
(b)	$\sqsubseteq [\text{nil}, \text{snoc}] R^\circ$	

$$[\text{nil}, \text{snoc}] \text{ cost} = \langle g, m \rangle \text{ bmax}$$

(10.4)


the cost of a schedule is the larger of the penalty of its last job and the cost of the schedule before it



(a)	$[\text{nil}, \text{snoc}] \text{ cost}$	$\text{cost}(xs \# [j]) = \text{bmax}(\text{cost } xs, \text{penalty}(xs, j))$ and bmax commutes — $\text{cost}(\text{wrap}(a)) = 0$, $\text{cost}(\text{snoc}(x_1, a)) =$ $\text{bmax}(\text{cost}(x_1), \text{penalty}(x_1, a))$
(b)	$= \langle g, m \rangle \text{ bmax}$	

$$(\forall j. 0 \leq ct(j)) \wedge (\forall j. 0 \leq wt(j)) \Rightarrow \text{add} \sqsubseteq \pi_1 R \quad (10.7)$$


putting one more job anywhere into a schedule never lowers its cost



(a)	add	induction on where j goes: every job after it starts later, so no penalty falls
(b)	$\sqsubseteq \pi_1 R$	

$$[\text{nil}, \text{snag}] \text{ bagify}^\circ = [\text{nil}, (\text{bagify}^\circ \times 1) \text{ add}] \quad (10.8)$$


the orderings of a bag with one more job are the orderings of the bag with that job put in anywhere



(a)	$[\text{nil}, \text{snag}] \text{ bagify}^\circ$	take out the last copy of j and put it back — $p \text{ add } w \Leftrightarrow \text{add}(\pi_2(p), w)$
(b)	$= [\text{nil}, (\text{bagify}^\circ \times 1) \text{ add}]$	

$$(\forall j. 0 \leq ct(j)) \wedge (\forall j. 0 \leq wt(j)) \Rightarrow [\text{nil}, \text{snag}] \text{ bagify}^\circ \sqsubseteq F(\text{bagify}^\circ) m \leq \text{cost}^\circ \quad (10.3c)$$


an ordering of a bag with one more job costs at least the cost of the ordering of the bag before its last job




(a)	$[\text{nil}, \text{snag}] \text{ bagify}^\circ$	(10.8) $[\text{nil}, \text{snag}] \text{ bagify}^\circ = [\text{nil}, (\text{bagify}^\circ \times 1) \text{ add}]$
(b)	$= [\text{nil}, (\text{bagify}^\circ \times 1) \text{ add}]$	(10.7) $(\forall j. 0 \leq ct(j)) \wedge (\forall j. 0 \leq wt(j)) \Rightarrow \text{add} \sqsubseteq \pi_1 R$
(c)	$\sqsubseteq [\text{nil}, (\text{bagify}^\circ \times 1) \pi_1 R]$	$[\text{nil}, \pi_1 R] \sqsubseteq [\text{zero}, \pi_1 \text{ cost}] \leq \text{cost}^\circ$
(d)	$\sqsubseteq F(\text{bagify}^\circ) [\text{zero}, \pi_1 \text{ cost}] \leq \text{cost}^\circ$	
(e)	$= F(\text{bagify}^\circ) m \leq \text{cost}^\circ$	

$$(\forall j. 0 \leq ct(j)) \wedge (\forall j. 0 \leq wt(j)) \Rightarrow (Q \cap ([nil, snag] [nil, snag]^\circ)) F(bagify^\circ) [nil, snoc] \sqsubseteq F(bagify^\circ) [nil, snoc] R \quad (10.3)$$

a schedule of a bag ending in a job of least penalty is no worse than any schedule of the same bag




(a)	$(Q \cap ([nil, snag] [nil, snag]^\circ)) F(bagify^\circ) [nil, snoc]$	$(S \cap T) R \sqsubseteq (S R) \cap (T R)$
(b)	$\sqsubseteq (Q F(bagify^\circ) [nil, snoc]) \cap ([nil, snag] [nil, snag]^\circ F(bagify^\circ) [nil, snoc])$	$[nil, snag]^\circ F(bagify^\circ) [nil, snoc] = bagify^\circ$
(c)	$= (Q F(bagify^\circ) [nil, snoc]) \cap ([nil, snag] bagify^\circ)$	$(10.3c) (\forall j. 0 \leq ct(j)) \wedge (\forall j. 0 \leq wt(j))$ $\Rightarrow [nil, snag] bagify^\circ \sqsubseteq F(bagify^\circ) m \leq cost^\circ$
(d)	$\sqsubseteq (Q F(bagify^\circ) [nil, snoc]) \cap (F(bagify^\circ) m \leq cost^\circ)$	$(Q F(bagify^\circ) [nil, snoc]) \cap (F(bagify^\circ) M) \sqsubseteq F(bagify^\circ) ((F(bagify^\circ) Q F(bagify^\circ)) \cap (M [nil, snoc]^\circ)) [nil, snoc]$




(a)	$\sqsubseteq F(bagify^\circ) ((F(bagify^\circ) Q F(bagify^\circ)) \cap (m \leq cost^\circ [nil, snoc]^\circ)) [nil, snoc]$	$F(bagify^\circ) Q F(bagify^\circ) = g \leq g^\circ$
(b)	$= F(bagify^\circ) ((g \leq g^\circ) \cap (m \leq cost^\circ [nil, snoc]^\circ)) [nil, snoc]$	$(R S)^\circ = S^\circ R^\circ$
(c)	$= F(bagify^\circ) ((g \leq g^\circ) \cap (m \leq ([nil, snoc] cost)^\circ)) [nil, snoc]$	$\langle X, Y \rangle \langle R, S \rangle^\circ = (X R^\circ) \cap (Y S^\circ)$
(d)	$= F(bagify^\circ) \langle g \leq, m \leq \rangle \langle g, [nil, snoc] cost \rangle^\circ [nil, snoc]$	$(10.3d) \langle g \leq, m \leq \rangle \langle g, [nil, snoc] cost \rangle^\circ$ $[nil, snoc] \sqsubseteq [nil, snoc] R$
(e)	$\sqsubseteq F(bagify^\circ) [nil, snoc] R$	

$$\langle g \leq, m \leq \rangle \langle g, [nil, snoc] cost \rangle^\circ [nil, snoc] \sqsubseteq [nil, snoc] R \quad (10.3d)$$

a job whose penalty is at most the last penalty, put after a schedule costing at most the cost before the last job, gives a schedule costing at most the whole



(a)	$\langle g \leq, m \leq \rangle \langle g, [nil, snoc] cost \rangle^\circ [nil, snoc]$	$1 \sqsubseteq cost cost^\circ$
(b)	$\sqsubseteq \langle g \leq, m \leq \rangle \langle g, [nil, snoc] cost \rangle^\circ [nil, snoc] cost cost^\circ$	$[nil, snoc] cost = \langle g, [nil, snoc] cost \rangle bmax$
(c)	$= \langle g \leq, m \leq \rangle \langle g, [nil, snoc] cost \rangle^\circ \langle g, [nil, snoc] cost \rangle bmax cost^\circ$	$\langle g, [nil, snoc] cost \rangle^\circ \langle g, [nil, snoc] cost \rangle \sqsubseteq 1$




(a)	$\sqsubseteq \langle g \leq, m \leq \rangle bmax cost^\circ$	$\langle g \leq, m \leq \rangle bmax \sqsubseteq \langle g, m \rangle bmax \leq$
(b)	$\sqsubseteq \langle g, m \rangle bmax \leq cost^\circ$	$(10.4) [nil, snoc] cost = \langle g, m \rangle bmax$
(c)	$= [nil, snoc] cost \leq cost^\circ$	$R = cost \leq cost^\circ$
(d)	$= [nil, snoc] R$	

$$\text{schedule} = (\text{null} \rightarrow \text{nil}, \text{pick}(\text{schedule} \times 1) \text{ snoc}), (\forall j. 0 \leq \text{ct}(j)) \wedge (\forall j. 0 \leq \text{wt}(j)) \Rightarrow (\mu X : \quad (10.3e)$$

$$\frac{[\text{nil}, \text{snag}]^\circ}{3} \text{ est}(Q) F(X) [\text{nil}, \text{snoc}] \sqsubseteq \frac{\text{bagify}^\circ}{3} \text{ est}(R)$$

an ordering of the given bag with least maximum penalty is got by taking a job of least penalty out of the bag, putting it last, and scheduling what is left the same way

the specification — $\text{bagify} \triangleq \text{bagify}$



(a)	$\frac{\text{bagify}^\circ}{3} \text{ est}(R)$	No greedy reduce exists — one would also solve every prefix of the input, and the best schedule of a prefix need not extend to a best schedule of the whole
(b)	$\sqsubseteq \frac{[\text{nil}, \text{snag}]^\circ}{3} \text{ est}(Q) F(X) [\text{nil}, \text{snoc}]$	Proposition 10.1: nil and snag have disjoint ranges
(c)	$\sqsubseteq \frac{\text{snag}^\circ}{3} \text{ est}(Q') (X \times 1) \text{ snoc}$	$\text{pick} \sqsubseteq \frac{\text{snag}^\circ}{3} \text{ est}(Q')$, a partial function, quadratic in the number of jobs
(d)	$\sqsubseteq \text{pick} (X \times 1) \text{ snoc}$	

§10.4. The TeX problem


(10.4a)

	name	type	definition	meaning
1	intern	$[\text{Digit}] \rightarrow [0, 2^{16})$	$\text{intern} \triangleq \text{val round}$	the value of a decimal fraction, rounded to 16 binary places
2	val	$[\text{Digit}] \rightarrow \text{Real}$	$\text{val} \triangleq ([\text{zero}, \text{shift}])$	the real number a digit list denotes after the decimal point
3	shift	$\text{Digit} \times \text{Real} \rightarrow \text{Real}$	$\text{shift} \triangleq \text{shift}, \text{shift}((d, r)) \triangleq (d + r)/10$	put digit d in front of r : $(d+r)/10$
4	round	$\text{Real} \rightarrow [0, 2^{16})$	$r \text{ round } n \iff (2n-1)/w < r \wedge r < (2n+1)/w$, $\text{round}^\circ = \text{interval inrange}$	the nearest integer to $2^{16}r$: $\text{round } r=n$ iff $2n-1 < 2^{17}r < 2n+1$
5	interval	$[0, 2^{16}) \rightarrow \text{Interval}$	$\text{interval}(n) \triangleq ((2n-1)/w, (2n+1)/w)$	the open interval $((2n-1)/2^{17}, (2n+1)/2^{17})$ of the reals that round to n
6	inrange	$\text{Interval} \rightarrow \text{Real}$	$p \text{ inrange } r \iff \text{lo } p < r \wedge r < \text{hi } p$	r lies strictly inside (a, b)
7	Legal(p)	Prop	$\text{Legal}(p) \triangleq 0 < \text{hi } p \wedge \text{hi } p < 1 \wedge \text{lo } p < \text{hi } p$	(10.9): $0 < b < 1$ and $a < b$; Interval is the legal pairs
8	arb	$1 \rightarrow \text{Interval}$	$x \text{ arb } p \iff \text{lo } p < 0 \wedge 0 < \text{hi } p$	any legal interval
9	step	$\text{Digit} \times \text{Interval} \rightarrow \text{Interval}$	$\text{step}(p) \triangleq (\text{shift}((\pi_1 p, \text{lo } (\pi_2 p))), \text{shift}((\pi_1 p, \text{hi } (\pi_2 p))))$	put digit d in front of both ends: $((d+a)/10, (d+b)/10)$
10	H	$\text{Interval} \rightarrow [\text{Digit}]$	$H \triangleq ([\text{arb}, \text{step}])^\circ$	every digit list whose interval the given one is
11	R	$[\text{Digit}] \rightarrow [\text{Digit}]$	$x R y \iff \text{length}(x) \leq \text{length}(y)$	x is no longer than y
12	l	$1 \rightarrow \text{FX}$	$l \triangleq \text{inl}$	the injection of the empty case into $\text{FX} = 1 + (\text{Digit} \times X)$
13	r	$\text{Digit} \times X \rightarrow \text{FX}$	$r \triangleq \text{inr}$	the injection of the digit case into $\text{FX} = 1 + (\text{Digit} \times X)$
14	!	$\text{Digit} \times \text{Interval} \rightarrow 1$	$! \triangleq \rightarrow ()$	the one map to 1
15	Q	$F(\text{Interval}) \rightarrow F(\text{Interval})$	$Q \triangleq (l^\circ !^\circ r) \cup 1$, $Q(u, z) \iff (\exists t. u = l(t) \wedge (\exists s. z = r(s))) \vee u = z$	u is the empty case and z a digit case, or $u=z$
16	w	Int	$w \triangleq 131072$	2^{17}

$$([arb, step]) \sqsubseteq val \text{ inrange}^\circ$$

(10.4b)

every interval got by folding a decimal's digits with $[arb, step]$ has the decimal's value strictly inside it




(a)	$F(\text{inrange}^\circ) [arb, step]$	$F(S) [T, U] = [T, (1 \times S) U]$
(b)	$= [arb, (1 \times \text{inrange}^\circ) step]$	$(1 \times \text{inrange}^\circ) step \sqsubseteq \text{shift inrange}^\circ$
(c)	$\sqsubseteq [arb, \text{shift inrange}^\circ]$	$\text{zero inrange}^\circ = arb$
(d)	$= [\text{zero inrange}^\circ, \text{shift inrange}^\circ]$	$[R, S] Z = [R Z, S Z]$
(e)	$= [\text{zero}, \text{shift}] \text{ inrange}^\circ$	


$$Q F(X) \alpha \sqsubseteq F(X) \alpha R$$

(10.4c)


choosing by Q among the one-step decompositions before building with $F(X)$ and α yields only decimals that are R -related to one built without choosing



(a)	$Q F(X) \alpha$	$Q \triangleq (\text{!}^\circ \text{!}^\circ r) \cup 1$
(b)	$\triangleq ((\text{!}^\circ \text{!}^\circ r) \cup 1) F(X) \alpha$	$(S \cup T) R = (S R) \cup (T R)$
(c)	$= (\text{!}^\circ \text{!}^\circ r F(X) \alpha) \cup (F(X) \alpha)$	$1 \sqsubseteq R$




(a)	$\sqsubseteq (\text{!}^\circ \text{!}^\circ r F(X) \alpha) \cup (F(X) \alpha R)$	$r F(X) \alpha \sqsubseteq \text{!}^\circ \text{!}^\circ R$
(b)	$\sqsubseteq (\text{!}^\circ \text{!}^\circ \text{!}^\circ \text{!}^\circ R) \cup (F(X) \alpha R)$	$\text{!}^\circ \text{!}^\circ \sqsubseteq 1$
(c)	$\sqsubseteq (\text{!}^\circ \text{!}^\circ R) \cup (F(X) \alpha R)$	$\text{!}^\circ F(X) = \text{!}^\circ$



(a)	$= (\text{!}^\circ \text{!}^\circ F(X) \alpha R) \cup (F(X) \alpha R)$	$\text{!}^\circ \text{!}^\circ \sqsubseteq 1$
(b)	$\sqsubseteq (F(X) \alpha R) \cup (F(X) \alpha R)$	$R \cup R = R$
(c)	$= F(X) \alpha R$	

$$\text{extern}(n)=f(2n-1,2n+1), \quad \text{interval} \left(\mu X : \frac{[\text{arb}, \text{step}]^\circ}{3} \text{est}(Q) F(X) \alpha \right) \sqsubseteq \frac{\text{intern}^\circ}{3} \text{est}(R) \quad (10.4d)$$

a shortest decimal whose internal representation is the given multiple of 2^{-16} is got by emitting the one digit the interval of admissible reals allows, until that interval contains zero and the empty decimal will do



(a)	$\text{interval} \left(\mu X : \frac{[\text{arb}, \text{step}]^\circ}{3} \text{est}(Q) F(X) \alpha \right)$	$\frac{[\text{arb}, \text{step}]^\circ}{3} \text{est}(Q) F\left(\frac{H}{3} \text{est}(R)\right) \alpha \sqsubseteq \frac{H}{3} \text{est}(R)$
(b)	$\sqsubseteq \text{interval} \frac{H}{3} \text{est}(R)$	$\frac{H}{3} \text{est}(R) \sqsubseteq \frac{\text{inrange val}^\circ}{3} \text{est}(R)$
(c)	$\sqsubseteq \text{interval} \frac{\text{inrange val}^\circ}{3} \text{est}(R)$	$\frac{\text{intern}^\circ}{3} = \text{interval} \frac{\text{inrange val}^\circ}{3}$
(d)	$= \frac{\text{intern}^\circ}{3} \text{est}(R)$	

- $f(a,b) = x \iff a < 0 \wedge x = [] \vee \neg a < 0 \wedge (\exists d. \exists y. d = [10b] \wedge f(10a-d, 10b-d) = y \wedge x = [d] \# y)$
 f gives (a,b) the empty decimal when $a < 0$, and otherwise the digit $d = [10b]$ followed by what it gives $(10a-d, 10b-d)$

(10.4e)

- $(\mu X : \frac{[\text{arb}, \text{step}]^\circ}{3} \text{est}(Q) F(X) \alpha) = f$
the least solution of the greedy recursion of $\text{interval} \left(\mu X : \frac{[\text{arb}, \text{step}]^\circ}{3} \text{est}(Q) F(X) \alpha \right) \sqsubseteq \frac{\text{intern}^\circ}{3} \text{est}(R)$ is f
- $d = [10b] \wedge e = [10b] \implies d = e$
Exercise 10.12: two digits d and e with $0 < 10b-d < 1$ and $0 < 10b-e < 1$ are equal
- $(0 < q-p \wedge q < w) \wedge p < 0 \implies f(p,q) = []$
the program's first clause: $f(p,q) = []$ when $p < 0$
- $(0 < q-p \wedge q < w) \wedge \neg p < 0 \wedge d = 10q/w \implies f(p,q) = [d] \# f(10p-wd, 10q-wd)$
the program's second clause: with $d = (10q) \text{ div } w$, $f(p,q)$ is d followed by $f(10p-w \cdot d, 10q-w \cdot d)$
- $(0 < q-p \wedge q < w) \wedge \text{Reach}(p,q) \implies f(p/w, q/w) = f(p,q)$
on the pairs the program reaches from $\text{interval } n$, the integer f at (p,q) is a value of the rational f at $(p/w, q/w)$
- $\text{interval } f = \text{extern}$
 f after interval is the program extern , whose value at n is $f(2n-1, 2n+1)$

§10. Appendix

§10.1. $P(S) \text{ est}(R) = (\exists S) n(\in \setminus (SR^\circ))$

(10.1a)

(p, q) tabulates $W \triangleq (\exists S) n(\in \setminus (SR^\circ))$, and $y \triangleq \frac{(p \exists S) n(qR)}{\exists}$, a map.

(10.1b)

$$\begin{array}{c} \boxed{\begin{array}{c} W \\ P(S) \text{ est}(R) \end{array}} \xleftarrow{p^\circ q = W, 1 \sqsubseteq yy^\circ} \begin{array}{c} \boxed{\begin{array}{c} p^\circ y \\ P(S) \end{array}} \\ \boxed{\begin{array}{c} y^\circ q \\ \text{est}(R) \end{array}} \end{array}$$

(10.1c)

$$\begin{array}{c} \boxed{\begin{array}{c} y^\circ q \\ \exists \end{array}} \xleftarrow{f^\circ \cdot \neg f^\circ, \cdot \exists \neg \exists} \boxed{\begin{array}{c} q \\ (p \exists S) n(qR) \end{array}} \xleftarrow{\Delta \neg n, f^\circ \cdot \neg f^\circ} \begin{array}{c} \boxed{\begin{array}{c} p^\circ q \\ \exists S \end{array}} \\ \boxed{\begin{array}{c} 1 \\ R \end{array}} \end{array}$$

R reflexive

$$\boxed{\begin{array}{c} \epsilon y^\circ q \\ R^\circ \end{array}} \xleftarrow{\cdot \exists \neg \exists, \circ, \text{meet}} \boxed{\begin{array}{c} R^\circ q^\circ q \\ R^\circ \end{array}} \quad \text{q simple}$$

(10.1d)

$$\begin{array}{c} \boxed{\begin{array}{c} p^\circ y \\ P(S) \end{array}} \xleftarrow{\Delta \neg n, \cdot T \neg / T, \circ, f^\circ \cdot \neg f^\circ} \begin{array}{c} \boxed{\begin{array}{c} p^\circ y \exists \\ \exists S \end{array}} \\ \boxed{\begin{array}{c} p \exists \\ y \exists S^\circ \end{array}} \end{array}$$

$$\boxed{\begin{array}{c} p^\circ y \exists \\ \exists S \end{array}} \xleftarrow{\cdot \exists \neg \exists, \text{meet}} \boxed{\begin{array}{c} p^\circ p \exists S \\ \exists S \end{array}} \quad \text{p simple}$$

$$\boxed{\begin{array}{c} p \exists \\ y \exists S^\circ \end{array}} \xleftarrow{\text{modular law}} \boxed{\begin{array}{c} p \exists \\ (p \exists) n(qRS^\circ) \end{array}} \xleftarrow{1 \sqsubseteq qq^\circ} \boxed{\begin{array}{c} q^\circ p \exists \\ RS^\circ \end{array}} \xleftarrow{\circ, T \cdot \neg T \setminus} \boxed{\begin{array}{c} W \\ \in \setminus (SR^\circ) \end{array}} \quad \text{W's right half}$$

§10.2. $P(\text{est}(R)) \text{ est}(R) = P(\text{dom}(\text{est}(R))) \text{ union est}(R)$

(10.2a)

$$\boxed{P(\text{est}(R)) \text{ est}(R)} \xrightarrow{\text{dom}(\text{est}(R)) \text{ est}(R) = \text{est}(R)} \boxed{P(\text{dom}(\text{est}(R)) \text{ est}(R)) \text{ est}(R)}$$

$$\begin{array}{c} \xrightarrow{\text{P a relator}} \boxed{P(\text{dom}(\text{est}(R))) P(\text{est}(R)) \text{ est}(R)} \xrightarrow{(7.11) R \sqsubseteq R \sqsubseteq R} \boxed{P(\text{dom}(\text{est}(R))) \text{ union est}(R)} \\ \Rightarrow P(\text{est}(R)) \text{ est}(R) \sqsubseteq \text{union est}(R) \end{array}$$

(10.2b)

$$\begin{array}{c} \boxed{\begin{array}{c} P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ P(\text{est}(R)) \text{ est}(R) \end{array}} \xleftarrow{1 \sqsubseteq R} \boxed{\begin{array}{c} P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ \exists \text{est}(R) \end{array}} \\ \Rightarrow P(S) \text{ est}(R) = (\exists S) n(\in \setminus (SR^\circ)) \text{ at } S := \text{est}(R), \\ \Delta \neg n, T \cdot \neg T \setminus \end{array}$$

$$\boxed{\begin{array}{c} \epsilon P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ \text{est}(R) R^\circ \end{array}} \xleftarrow{\text{count of } T \cdot \neg T \setminus} \boxed{\begin{array}{c} \exists (\exists n(\in \setminus (\in \setminus (R^\circ)))) \\ \exists (\exists n(\in \setminus (R^\circ))) \end{array}}$$

$$\boxed{\begin{array}{c} P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ \exists \text{est}(R) \end{array}} \xleftarrow{P(\text{dom}(\text{est}(R))) \sqsubseteq 1} \boxed{\begin{array}{c} \text{union est}(R) \\ \exists \text{est}(R) \end{array}}$$

$$\begin{array}{c} \xleftarrow{T(UnV) \sqsubseteq TUnTV, \cdot \exists \neg \exists, (7.3) \text{ union } (\in \setminus R) = \in \setminus (\in \setminus R)} \boxed{\begin{array}{c} (\exists \exists) n(\in \setminus (\in \setminus (R^\circ))) \\ \exists \text{est}(R) \end{array}} \xleftarrow{\text{modular law}} \boxed{\begin{array}{c} \exists (\exists n(\in \setminus (\in \setminus (R^\circ)))) \\ \exists (\exists n(\in \setminus (R^\circ))) \end{array}} \end{array}$$

count of $T \cdot \neg T \setminus$

$$\boxed{\begin{array}{c} \epsilon P(\text{dom}(\text{est}(R))) \text{ union est}(R) \\ \text{est}(R) R^\circ \end{array}} \xleftarrow{\in \text{ lax natural}} \boxed{\begin{array}{c} \text{dom}(\text{est}(R)) \epsilon \text{union est}(R) \\ \text{est}(R) R^\circ \end{array}}$$

$$\begin{array}{c} \xleftarrow{\text{dom}(\text{est}(R)) \sqsubseteq \text{est}(R) \text{ est}(R)^\circ} \boxed{\begin{array}{c} \text{est}(R) \text{ est}(R)^\circ \epsilon \text{union est}(R) \\ \text{est}(R) R^\circ \end{array}} \xleftarrow{\text{est}(R)^\circ \sqsubseteq \in, \in \epsilon \text{union} \sqsubseteq \in} \boxed{\begin{array}{c} \text{est}(R) \epsilon \text{est}(R) \\ \text{est}(R) R^\circ \end{array}} \end{array}$$

UP of est

§10.3. Shortest paths on a cylinder, on lists

(10.3a)

	definition	type	note
1	$F(A,X)=A+A \times X$	$\mathcal{A} \rightarrow \mathcal{A}$	The base functor: one square of the new column, alone or in front of the path so far.
2	$L = \text{list}^+$	$\mathcal{A} \rightarrow \mathcal{A}$	A path is a non-empty list of squares, one per column crossed.
3	α the initial algebra	$F(\mu F) \rightarrow \mu F$	$\alpha(5)=[5]$ and $\alpha(1,[5])=[1,5]$: a path is started by one square, or extended by one.
4	N the n-tuple relator	$\mathcal{A} \rightarrow \mathcal{A}$	One component per row of a column, so the fold carries n answers at once.
5	$R \triangleq \text{sum} \leq \text{sum}^\circ$	$[\mathbb{N}] \rightarrow [\mathbb{N}]$	The cost of a path, which the cheapest minimises.
6	setify	$A[n] \rightarrow PA$	$\text{setify}(1,2,3,4)=\{1,2,3,4\}$ — which row a component came from is forgotten.
7	moves	$A[n] \rightarrow P(A[n])$	$\text{moves}(x)=\{\text{up}(x), x, \text{down}(x)\}$ — rotated up, unrotated, rotated down.
8	trans	$P(A[n]) \rightarrow (P A)[n]$	$\text{trans}\{(a,b,c), (x,y,z)\} = (\{a,x\}, \{b,y\}, \{c,z\})$ — component k of the result is the set of the k-th components.
9	zip	$F(a[n]) \rightarrow (F a)[n]$	$\text{zip}((1,2,3,4), (\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\})) = ((1, \{[5]\}), (2, \{[6]\}), (3, \{[7]\}), (4, \{[8]\}))$.
10	$\text{cp} \triangleq \frac{F(1, \exists)}{\exists}$	$F(PA) \rightarrow PFA$	$\text{cp}(1, \{[5], [6], [8]\}) = \{(1, [5]), (1, [6]), (1, [8])\}$.
11	$\text{gen} \triangleq F(1, \text{moves trans } N(\text{union}))$ $\text{zip } N(\text{cp}(F(A, -)) E(\alpha))$	$F(NA, N(P(TA))) \rightarrow N(P(TA))$	$\text{gen}((1,2,3,4), (\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\}))$ is worked out in $\text{gen} \triangleq F(1, \text{moves trans } N(\text{union})) \text{zip } N(\text{cp}(F(A, -)) E(\alpha))$.
12	$\text{paths} \triangleq \langle \text{gen} \rangle \text{setify union}$	$T(NA) \rightarrow P(TA)$	$\text{paths}[(1,2,3,4), (5,6,7,8)]$ is the union of $\text{gen} \triangleq F(1, \text{moves trans } N(\text{union})) \text{zip } N(\text{cp}(F(A, -)) E(\alpha))$'s four sets: 12 paths, 3 from each entry row.
13	the specification $\text{paths est}(R)$	$[\mathbb{N}[n]] \rightarrow [\mathbb{N}]$	A cheapest path from the entry side to the exit side.

§10.3.1. $N(E(L A))$

(10.3.1a)

```

1  5
2  6      [(1,2,3,4), (5,6,7,8)] : L N Nat
3  7
4  8

```

(10.3.1b)

```

gen  $\triangleq$  F(1, moves trans N(union)) zip N(cp(F(A, -)) E( $\alpha$ ))

⟨gen⟩[(5,6,7,8)] = ({[5]}, {[6]}, {[7]}, {[8]})

gen((1,2,3,4), ({[5]}, {[6]}, {[7]}, {[8]})) =
( {[1,5], [1,6], [1,8]},
  {[2,5], [2,6], [2,7]},
  {[3,6], [3,7], [3,8]},
  {[4,5], [4,7], [4,8]} )

```




(10.3.1c)

1	N	the 4-tuple, one component per row of the entry column
2	E	the set of paths that can start in that row
3	L	a path is the non-empty list of the squares it crosses

row 1's set has no $[1,7]$ — from row 1 only rows 4, 1 and 2 are reachable — and it does have $[1,8]$, because the cylinder glues the bottom row to the top. That is $\text{moves trans } N(\text{union})$ of $\text{gen} \triangleq F(1, \text{moves trans } N(\text{union})) \text{ zip } N(\text{cp}(F(A, -)) E(\alpha))$: component k collects the paths of rows $k-1, k, k+1$.

§10.3.2. The cross product $\text{cp} \triangleq \frac{F(1, \exists)}{\exists}$

(10.3.2a)

cp on all of $F(A, E(L A))$ the $A \times -$ summand: $\exists$ picks one path, $1\% \exists$ collects the resultsthe A summand, no E to distribute: $1\% \exists$ alone, $a \mapsto \{a\}$

(10.3.2b)


type	note
1 $F(A, -) = A + A \times -$	one wire carries the whole functor, both summands with it; the $\exists$ under it is $F(1, \exists)$
2 $\exists : E(L A) \rightarrow L A$	the $\exists$ inside $F(1, \exists)$: a set of paths, one of them — $L A$ is one path, A one square
3 $F(1, \exists) = 1 + 1 \times \exists$	$: A + A \times E(L A) \rightarrow A + A \times L A$, the relator acting on each summand
4 $\exists : E(A + A \times L A) \rightarrow A + A \times L A$	the $\exists$ under the bar: a set of cells, one of them — a different $\exists$
5 $\text{cp} : A + A \times E(L A) \rightarrow E(A + A \times L A)$	$\frac{R}{\exists}$ turns $R : X \rightarrow Y$ into $X \rightarrow E Y$
6 $\text{cp} = \frac{1}{\exists} E(F(1, \exists))$	what the picture draws: the unit makes the outer E , leaving one $\exists$ — $\frac{R}{\exists} = \frac{1}{\exists} E(R)$

(10.3.2c)

$w = (1, \{[5], [6], [8]\})$: $A \times E(L A)$ the right summand
 $1 \times \exists : A \times E(L A) \rightarrow A \times L A$ 1 keeps the square 1, $\exists$ picks one path
 $w \mapsto (1, [5]), (1, [6]), (1, [8])$: $A \times L A$ one output per path in w
 $\text{cp}(w) = \{(1, [5]), (1, [6]), (1, [8])\}$: $E(A + A \times L A)$ the three of them, collected

§10.3.3. $\text{gen} = F(1, \text{moves trans } N(\text{union})) \text{ zip } N(\text{cp } P(\alpha))$

(10.3.3a)




(10.3.3b)

$\text{gen} \triangleq F(1, \text{moves trans } N(\text{union})) \text{ zip } N(\text{cp}(F(A, -)) E(\alpha))$
 $u = ((1, 2, 3, 4), (\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\})) : F(N A, N(E(L A)))$
 $F(1, \text{moves trans } N(\text{union}))$ 1 keeps the column, the path SETS move
 $\text{moves}(\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\})$
 $= (\{[6]\}, \{[7]\}, \{[8]\}, \{[5]\}),$
 $(\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\}),$
 $(\{[8]\}, \{[5]\}, \{[6]\}, \{[7]\})$: $E(N(E(L A)))$ down, unmoved, up
 $\text{trans}(\text{that})$
 $= (\{[6]\}, \{[5]\}, \{[8]\}, \{[7]\}, \{[6]\}, \{[5]\}),$
 $\{[8]\}, \{[7]\}, \{[6]\}, \{[5]\}, \{[8]\}, \{[7]\})$: $N(E(E(L A)))$ row k gets rows k-1, k, k+1
 $N(\text{union})(\text{that})$
 $= (\{[5], [6], [8]\}, \{[5], [6], [7]\},$
 $\{[6], [7], [8]\}, \{[5], [7], [8]\})$: $N(E(L A))$ every path into row k, none dropped
 $\text{zip}(\text{that})$
 $= ((1, \{[5], [6], [8]\}), (2, \{[5], [6], [7]\}),$
 $(3, \{[6], [7], [8]\}), (4, \{[5], [7], [8]\}))$: $N(F(A, E(L A)))$ each row: its square, and the paths it may be put in front of
 $N(\text{cp } P(\alpha))(\text{that})$
 $= (\{[1, 5], [1, 6], [1, 8]\}, \{[2, 5], [2, 6], [2, 7]\},$
 $\{[3, 6], [3, 7], [3, 8]\}, \{[4, 5], [4, 7], [4, 8]\})$: $N(E(L A))$ cp pairs the square with each path, α prefixes it: $\alpha(1, [5]) = [1, 5]$

§10.3.4. gen is an F-algebra; $(\text{gen}) : \alpha(\text{gen}) = F(1, (\text{gen})) \text{gen}$

(10.3.4a)

 α puts the column back on the list, then the fold reads all of itthe fold reads the rest under F , then one gen puts the column in front

$$xs = [(1, 2, 3, 4), (5, 6, 7, 8)] = \alpha((1, 2, 3, 4), [(5, 6, 7, 8)]) : L(N A) \quad (10.3.4b)$$

$$(\text{gen})(xs) = \text{gen}((1, 2, 3, 4), (\text{gen})[(5, 6, 7, 8)])$$

both halves are in $\text{gen} \triangleq F(1, \text{moves trans } N(\text{union})) \text{ zip } N(\text{cp}(F(A, -)) E(\alpha))$: the tail folds to $(\{[5]\}, \{[6]\}, \{[7]\}, \{[8]\})$, one path per row and each of them one square long, and gen on it is $\text{gen} \triangleq F(1, \text{moves trans } N(\text{union})) \text{ zip } N(\text{cp}(F(A, -)) E(\alpha))$'s walk.

§10.3.5. $Q : F(N A, N(L A)) \rightarrow N(L A)$, an $F(N A, -)$ -algebra: $Q = F(1, \text{moves trans } N(\text{est}(R))) \text{ zip } N(\alpha)$

(10.3.5a)

$$Q : F(N A, N(L A)) \rightarrow N(L A)$$

the fold's algebra: a new column, one path per row

$u = ((1, 2, 3, 4), ([5], [6], [7], [8])) : F(N A, N(L A))$
 $F(1, \text{moves trans } N(\text{est}(R)))$ 1 keeps the column, the paths move
 $\text{moves}([5], [6], [7], [8])$
 $= ([6], [7], [8], [5]),$
 $([5], [6], [7], [8]),$
 $([8], [5], [6], [7])$: $E(N(L A))$ down, unmoved, up
 $\text{trans}(\text{that})$
 $= ([6], [5], [8]), ([7], [6], [5]),$
 $[8], [7], [6]), ([5], [8], [7])$: $N(E(L A))$ row k gets rows k-1, k, k+1
 $N(\text{est}(R))(\text{that})$
 $= ([5], [5], [6], [5])$: $N(L A)$ the cheapest into each row – chosen BEFORE the new square
 $\text{zip}(\text{that})$
 $= ((1, [5]), (2, [5]), (3, [6]), (4, [5]))$: $N(F(A, L A))$ each row: its square, and the one predecessor that survived
 $N(\alpha)(\text{that})$
 $= ([1, 5], [2, 5], [3, 6], [4, 5])$: $N(L A)$ $\alpha(1, [5]) = [1, 5]$

$$F(1, \text{moves trans } N(\text{est}(R))) = [l, pm \ r] \wedge zip = [N(inl), zip' \ N(inr)] \wedge inl \ \alpha = wrap \wedge inr \ \alpha \quad (10.3.5b)$$

$$= cons \Rightarrow Q = [N(wrap), pm \ zip' \ N(cons)]$$

at the last column Q starts one path per row, and at each earlier one it puts each square in front of the cheapest of the three kept paths it can step to — the algebra $R \ R \sqsubseteq R \wedge F(1, N(\text{est}(R))) \ Q \sqsubseteq \text{gen } N(\text{est}(R))$


$\Rightarrow (Q)$ setify $\text{est}(R) \sqsubseteq \text{paths } \text{est}(R)$'s last step folds

circuit — the fork is $F(NA, N(LA)) = NA + NA \times N(LA)$

Hinze–Marsden

$$F(N \ A) \ (N \ (T \ A)) \rightarrow \boxed{Q} \rightarrow N(T \ A)$$


$$F(N \ A, N \ (T \ A))$$



$$FN(T \ A) \rightarrow \boxed{F(1, \text{moves trans } N(\text{est}(R)))} \rightarrow FN(T \ A) \rightarrow \boxed{zip} \rightarrow N(F \ A \ (T \ A)) \rightarrow \boxed{N(\alpha)} \rightarrow N(T \ A)$$

= the fusion condition $R \ R \sqsubseteq R \wedge F(1, \text{est}(R)) \ \alpha \sqsubseteq cp(F(A, -)) \ E(\alpha) \ \text{est}(R)$
 $\Rightarrow F(1, N(\text{est}(R))) \ Q \sqsubseteq \text{gen } N(\text{est}(R))$ read as a definition,

$$F(NA, -) \ N \ TA$$



$$F(N \ A) \ (N \ (T \ A)) \rightarrow \left(\begin{array}{l} \boxed{l} \rightarrow N(p_0) \rightarrow \boxed{N(wrap)} \\ \boxed{r} \rightarrow q \rightarrow \boxed{pm} \rightarrow q \rightarrow \boxed{zip'} \rightarrow N(q_0) \rightarrow \boxed{N(cons)} \end{array} \right) \rightarrow N(T \ A)$$

$$zip = 1 + zip', \ \alpha = [wrap, cons]$$


$$R \subseteq R \wedge F(1, N(\text{est}(R))) \quad Q \subseteq \text{gen } N(\text{est}(R)) \Rightarrow \langle Q \rangle \text{ setify } \text{est}(R) \subseteq \text{paths } \text{est}(R) \quad (10.3.5c)$$

shortest paths on a cylinder: the dynamic programming approach to exhaustive search allows a path of least cost to be found in $O(n \times m)$ time; Q is $F(1, \text{moves trans } N(\text{est}(R))) = [l, pm \ r] \wedge \text{zip} = [N(\text{inl}), \text{zip}' \ N(\text{inr})] \wedge \text{inl } \alpha = \text{wrap} \wedge \text{inr } \alpha = \text{cons}$
 $\Rightarrow Q = [N(\text{wrap}), pm \ \text{zip}' \ N(\text{cons})]$'s algebra.

circuit


Hinze–Marsden

$$T(N \ A) \text{---} \boxed{\text{paths}} \text{---} P(T \ A) \text{---} \boxed{\text{est}(R)} \text{---} T \ A$$




$$= T(N \ A) \text{---} \boxed{[F(N \ A) \ (N \ (P \ (T \ A)))]} \text{---} \boxed{\text{gen}} \text{---} NP(T \ A) \text{---} \boxed{\text{setify}} \text{---} PP(T \ A) \text{---} \boxed{\text{union}} \text{---} P(T \ A) \text{---} \boxed{\text{est}(R)} \text{---} T \ A$$

$\text{paths} \triangleq \langle \text{gen} \rangle \text{ setify union at paths}$




$$\sqsubseteq T(N \ A) \text{---} \boxed{[F(N \ A) \ (N \ (P \ (T \ A)))]} \text{---} \boxed{\text{gen}} \text{---} NP(T \ A) \text{---} \boxed{\text{setify}} \text{---} PP(T \ A) \text{---} \boxed{P(\text{est}(R))} \text{---} P(T \ A) \text{---} \boxed{\text{est}(R)} \text{---} T \ A$$

$P(\text{est}(R)) \text{ est}(R) \sqsubseteq \text{union } \text{est}(R)$ — a minimum in each set, then a minimum of those; R transitive




$$\sqsubseteq T(N \ A) \text{---} \boxed{[F(N \ A) \ (N \ (P \ (T \ A)))]} \text{---} \boxed{\text{gen}} \text{---} NP(T \ A) \text{---} \boxed{N(\text{est}(R))} \text{---} N(T \ A) \text{---} \boxed{\text{setify}} \text{---} P(T \ A) \text{---} \boxed{\text{est}(R)} \text{---} T \ A$$

$\text{setify lax natural}$



$$\sqsubseteq T(N \ A) \text{---} \boxed{[F(N \ A) \ (N \ (T \ A))]} \text{---} \boxed{Q} \text{---} N(T \ A) \text{---} \boxed{\text{setify}} \text{---} P(T \ A) \text{---} \boxed{\text{est}(R)} \text{---} T \ A$$

fusion at $R \subseteq R \wedge F(1, \text{est}(R)) \quad \alpha \sqsubseteq \text{cp}(F(A, -)) \ E(\alpha) \ \text{est}(R)$
 $\Rightarrow F(1, N(\text{est}(R))) \quad Q \subseteq \text{gen } N(\text{est}(R))$



$R \subseteq R \wedge F(1, \text{est}(R)) \alpha \subseteq \text{cp}(F(A, -)) E(\alpha) \text{ est}(R) \Rightarrow F(1, N(\text{est}(R))) Q \subseteq \text{gen } N(\text{est}(R))$ (10.3.5d) fusion: the condition for fusion in $R \subseteq R \wedge F(1, N(\text{est}(R))) Q \subseteq \text{gen } N(\text{est}(R))$ $\Rightarrow (Q) \text{ setify } \text{est}(R) \subseteq \text{paths } \text{est}(R)$'s last step, used to derive a definition of Q .	
circuit	Hinze–Marsden
$F(N \ A) \ (N \ (P \ (T \ A))) \text{---} \boxed{\text{gen}} \text{---} NP(T \ A) \text{---} \boxed{N(\text{est}(R))} \text{---} N(T \ A)$	
$\sqsubseteq FNP(T \ A) \text{---} \boxed{F(1, N(\text{est}(R)))} \text{---} FN(T \ A) \text{---} \boxed{Q} \text{---} N(T \ A)$ (7.13), then zip, trans, moves lax natural	

(7.13) is $F(1, \text{est}(R)) \alpha \subseteq \text{cp } P(\alpha) \text{ est}(R)$, Theorem 7.1 $\text{Map}(f) \wedge \text{PreservesRecip}(F) \wedge R^\circ \subseteq \text{est}(R)$
 $\Rightarrow f^\circ F(R) f \subseteq R \iff F(\text{est}(R)) f \subseteq \frac{F(\exists) f}{\exists} \text{est}(R)$ at the map α with $\frac{F(1, \exists) \alpha}{\exists} = \text{cp } P(\alpha)$: extending every path in a set and then taking a minimum is beaten by extending one minimum. It is the crux here, not the greedy theorem.