The allegory axioms, and what defines them in the Frobenius calculus

Freyd, Categories, Allegories §2.11–§2.12 against FunctorialSemanticsForRelationalTheories.pdf Def. 4.1 and Lean Freyd/S2_10.lean:Allegory, diag/CB.lean:CartBicat, diag/CB_Allegory.lean:allegoryOfCartBicat

Where the pictures come from. Every picture of a theorem is emitted by `./scripts/diag-export` from that theorem's Lean type. Nothing about a statement is redrawn here. The two figures marked transcribed are the exception: they copy the paper's definition pictures, and a definition has no statement to export.

Equation numbering. The numbers below are the 2017 preprint's. The CartBicat fields no longer encode them — a field is named for what it says, `+Tbd↑` rather than `Lemg_37` — and each docstring carries the preprint's number. The repo's copy of the paper is now the published version — Fong, Spivak, Math. Z. (2019?) — whose numbering differs: (37)↔(40), (38)↔(41), (39)↔(42), (40)↔(43), (33)↔(35), (34)↔(36), (35)↔(37), (36)↔(38), (42)↔(44), (43)↔(45), and Example 2.16(?)↔Lemma 4.1 at (?)–(?)(?). Renumbering the Lean fields to match the published numbers has not been made.

Freyd's allegory takes reciprocation ° and intersection ∩ as primitive operations and lists eight equational axioms on them. A cartesian bicategory of relations takes neither: every object carries one special Frobenius structure Δ, !, ∇, ¡, the hom-order ≤ is primitive, and both ° and ∩ are defined from the four dots. Every one of Freyd's eight axioms is then a theorem. The table below is that dictionary, row per row, axiom.

Reading the pictures: left to right is composition — the book's diagrammatic order, so R ; S means first R then S — vertical stacking is ⊗, a bare wire is an identity, a white box is a relation, and a box that is mirrored and turned is its converse — the mirroring is the paper's own notation, the text is this note's, because at a glance a chamber is easy to read a single letter and still be read, a whole term like R ; S as a box. A meet or a converse as a labelled box, so a composite under a ° spans the run; and any wiring with no shape is one of the Frobenius... [illegible] converse R°, Freyd's symbol, where the paper writes R†: these terms and ... [illegible] the reader to match them. The Lean sources write R° as ... §7 is a worked example in other models — see the caution at the end of ... Rel(Set) throughout there is no clash for context against.

In a chain like the tabled at the head of each row is the relation that ... Counting them is the point of drawing a proof at all — a chain of one bookkeeping.

1. Definition 4.1, as the paper states it

`FunctorialSemanticsForRelationalTheories.pdf` Def. 4.1, pp. 20–21 — transcribed here as `class CartBicat` (diag/CB.lean): field for field. Every picture below is reported from the Lean field, so the class and the definition cannot drift apart.

A **cartesian bicategory of relations** is a poset-enriched category, symmetric monoidal, satisfying four conditions. Clauses 1 and 2 are the two structures every object carries; clause 3 makes each the other's adjoint and imposes Frobenius; clause 4 is the only condition on arrows.

The tower this note describes is STRICT: the two bracketings of a threefold product are the same object, so no associator or unitor appears in any diagram. The coherence arrows in any diagrams are identities.

1.1. Every object carries a cocommutative comonoid (Δ, !)



(8) **coassociativity** `CartBicat.Δ_assoc` — copying twice on the left is copying twice on the right.	
(9) **cocommutativity** `CartBicat.Δ_comm` — the two copies are interchangeable.	
(10) **counit** `CartBicat.Δ_counit` — copy, then discard one copy, and nothing has happened.	

1.2. Every object carries a commutative monoid (∇, ¡)



(5) **associativity** `CartBicat.∇_assoc`	
(6) **commutativity** `CartBicat.∇_comm`	
(7) **unit** `CartBicat.∇_unit` — create, then merge, and nothing has happened.	

1.3. Five inequalities relate the monoid and the comonoid



(37) ∇ ; Δ ≤ 1 — half of Δ ⊣ ∇

(38) 1 ≤ Δ ; ∇ — the other half

(39) ! ; ¡ ≤ 1 — half of ! ⊣ ¡

(40) 1 ≤ ¡ ; ! — the other half

(41) the **Frobenius law** `CartBicat.Δ_frob_left` and its mirror `frob_right`. This is the one equation of the group; the other four are inequalities

1.4. Every arrow is a lax comonoid homomorphism



(42) R ; Δ ≤ Δ ; (R ⊗ R) `CartBicat.lax_Δ` — running R once and copying is below copying and running it on both strands.

(43) R ; ! ≤ ! `CartBicat.lax_!`

The paper derives it — "one direction is given by (38) and the other is proved as follows" — and so does the Lean: `Freyd.Diag.CartBicat.special`; its picture is in the correspondence table below.

Read as inequalities in one direction only: an arrow that satisfied these as EQUATIONS would be a map, and the whole point is that a general relation is not.

2. Freyd's primitives are the Frobenius calculus's definitions

Freyd's primitive, and what defines it here	picture
reciprocation R° — §2.11, primitive. Here conv R := bend ((R ⊗ 1) ; cap) (diag/CB_Ideal.lean:223), not a generator. Input arrives bottom left and runs under R, the cap ⊤ ; ¡ Δ at the top left creates a pair, one strand feeding R and the other leaving as the output at the top right; the cup ∇ ; ! at the bottom right combines R's output against the input. So the cup and the cap that reverse the arrow — a wire merely bumped up over the box and back down is an isotopy of R itself	 transcribed
intersection R ∩ S — §2.11, primitive. Here the convolution meet R ⋒ S := Δ ; (R ⊗ S) ; ∇ (diag/CB_Derived.lean:24). Copy the input, run both, merge: the merge forces the two results to agree.	 transcribed
containment R ⊆ S — §2.11, defined as R ∩ S = R. Here the poset-enrichment ≤ is primitive and ⊆ is derived, so the two directions are swapped. `meet_eq_left_of_le` `CB_Derived.lean:196` is the bridge.	no picture — this is the 2-cell itself
maximal morphism ⊤ — §2.15, and Freyd needs a unitary allegory for it. Here ⊤ := ! ; ¡ (diag/CB_Derived.lean:28) is free in Def. 4.1. `Freyd.Diag.≤_top`	

3. The eight axioms, and what proves each

Freyd's axiom, and what supplies it	picture, from the Lean statement
(R°)° = R — §2.11 `recip_recip`: here the **snake**, from (41). `Freyd.Diag.CartBicat.conv_conv`. Straightening down and back up straightens out, so bending twice is the identity.	
(R S)° = S° R° — §2.11 `recip_comp`: here mirroring the picture. `Freyd.Diag.CartBicat.conv_comp`. Reflecting a chain left to right reverses it: each box comes back upright.	
(R ∩ S)° = R° ∩ S° — §2.11 `recip_inter`: the same mirroring. `Freyd.Diag.conv_inter_Δ.sus(?)` acts each other's mirror image, so mirroring commutes with the meet, and only turns the boxes over.	
R ∩ R = R — §2.11 `inter_idem`: here (42) and the **special law**. `Freyd.Diag.meet_idem`. Copy, run R twice, merge — (42) pushes R back through the copy and Δ ; ∇ = 1 closes the bubble.	
R ∩ S = S ∩ R — §2.11 `inter_comm`: here cocommutativity (9) and commutativity (6). `Freyd.Diag.meet_comm`. The two strands of Δ are interchangeable, and so are those of ∇.	
R ∩ (S ∩ T) = (R ∩ S) ∩ T — §2.11 `inter_assoc`: here coassociativity (8) and associativity (5). `Freyd.Diag.meet_assoc`. Either bracketing builds the same three-way copy-tree, and finally the same merge tree.	
semi-distributivity R (S ∩ T) ⊆ R S ∩ R T — §2.11 `semidistrib`, §2.11b: here (42), the lax **comonoid inequation**. `Freyd.Diag.semidistrib_of_lax`. Running R and then copying is below copying and then running R on each strand.	
modular identity R S ∩ T ⊆ R (S ∩ R° T) — §2.112 `modular`, the axiom **Freyd adjoins**: "we adjoin one more containment, the law of modularity". Here (41) and (42), derived. `Freyd.Diag.modular_of_frobenius`. Using the dropped prefix Δ ; (R ⊗ T) off both sides and what is left is `≈(?)slide_conv.diag(CB.lean:560)` — a box slides through a wedge at the cost of its converse on the other strand.	

4. The other direction: what each law of Def. 4.1 says in Freyd's language

picture, from the Lean statement	Def. 4.1	Freyd
	(37) ∇ ; Δ ≤ 1 — half of Δ ⊣ ∇	Δ° Δ ⊆ 1 — **Δ is simple** §2.13. Not used anywhere in the bridge; see below.
	(38) 1 ≤ Δ ; ∇ — the other half	1 ⊆ Δ Δ° — **Δ is entire** §2.13; with (37), Δ ; ∇ = 1 is a map.
	(39) ! ; ¡ ≤ 1 — half of ! ⊣ ¡	!° ! ⊆ 1 — **! is simple** §2.13. Also unused in the bridge.
	(40) 1 ≤ ¡ ; ! — the other half	1 ⊆ ! !° — **! is entire** §2.13; with (39), ! is a map; this is what makes ⊤ = ! ; ¡ contain every arrow.
	(41) the **Frobenius law** — Δ and ∇ as an axiom; ∇ ⊗ 1 ; Δ drawn as stacked merging associator Δ ; ? drawn as ... box, having the two merging; the shape in this tensor.	**the modular identity** §2.112 — and reciprocation §2.11 itself: the cup and cap only satisfy the snake because of (41).
	the **special law** not an axiom: derived from (38) and (40). `Freyd.Diag.CartBicat.special`	Δ ; Δ° = 1 — **Δ is injective**, so by §2.141 of ⊤ = g ; Δ it is a mono itself.
	(42) **every arrow is a lax comonoid homomorphism**	§2.136's R (S ∩ T) ⊆ R S ∩ R T, read at S := 1(?), T := 1(?) — Freyd's semi-distributivity, product-free.
	(43) **every arrow is lax for the counit**	§2.13's step `!_p,a` is maximal in (a, ?), hence R ; ! ⊆ ! — that is, R ⊆ ⊤.

5. Unitary and pre-tabular — the two conditions the converse needs

Freyd's §2.11 axioms are all a cartesian bicategory has to give back. Going the other way needs two further conditions, and they are exactly the two the repo has not formalised. Both definitions below are from `Freyd/S2_10.lean`, which follows the book.

condition	Freyd	what the Frobenius side already has
unitary §2.15	**I** is a partial unit when 1 is the largest endomorphism on **I**, so `PartialUnit I := ∀ R : I → I, R ⊆ 1` (S 539); it is a **unit** when moreover every object carries an entire arrow to it, `IsUnit I := PartialUnit I ∧ ∀ a, ∃ R : a → I, Entire R` (S 545). `UnitaryAllegory` is an allegory with one (S 548). The point of a unit is that it makes ⊤_(a, b) exist: ⊤ = p_R° — the tops of the tails, which is where §2.15(?)'s "R ⊆ p_R° p_R" — the topos(?)'s (15) — comes from.	**already there, for free.** The monoidal unit I is the unit object: `!_a : a → I` is the only arrow every object carries to an entire arrow to it — (40) says so — and ⊤ ; ! : I → I is maximal by `Freyd.Diag.≤_top`. So every object is (?) unitary in the sense of entirely assumed. Not recorded in Lean: `allegoryOfCartBicat` is a plain `Allegory`, and no `UnitaryAllegory` instance is derived from `CartBicat`.
pre-tabular §2.145	f, g **tabulate** R when both are maps, R = f° g, and f f° ∩ g g° = 1 (Tabulates, `(S2.?) R` in §2.143(?)); R is **tabular** when some pair tabulates it (S 596). An allegory is `PreTabularAllegory` when every morphism is contained in a tabular one (S 556). §2.145: "If R is unitary then pre-tabularity is equivalent to existence of tabulations for maximal morphisms."	**this is what ⊗ is.** Given unitarity, the condition reduces to tabulating each ⊤_(a, b), and a tabulation of ⊤_(a, b) is precisely an object a ⊗ b with its two projections — the monoidal product Def. 4.1 assumes. That is the whole content of the allegory fact (the product the tabulations and ⊗ the relation (?)... not done (?), Map(R) a CartBicat.UnitaryPretabular. `diag/PLAN.md` holds it back for want of a consumer.

Together they are the whole gap: **cartesian bicategory of relations = unitary pre-tabular allegory** (`DiagrammaticAlgebraOfFirstOrderLogic` §10, after Carboni–Walters). One half of that equivalence is `allegoryOfCartBicat` and is a theorem in this repo, the other half is the converse.

6. Union and residual — what phases 8 and 9 add

∪ and / are past where Freyd's §2.11 list stops, and past what Def. 4.1 can build: they need a **biproduct** on top of the Frobenius structure (`diag/Tape.lean`, after `TapeDiagrams.pdf`) and a **second composition** with linear adjoints (`diag/FO.lean`, after `DiagrammaticAlgebraOfFirstOrderLogic.pdf`). Both are built, so both are drawn.

The two notations say which layer a picture is in. A union is a **tape**: the rounded wrapper is the second monoidal product, and its ∪-open and close a branch a particle takes exactly one of — the same S... A residual is a **dashed** box, because it is not a composite of the generators at all: its one box would claim it was drawable in the first.

statement	picture, from the Lean statement
R ⊆ R ∪ S — §2.2 in Freyd; here `Freyd.Diag.≤union_left`. `union R S := (R, S) ; (1, 1)diag/Tape.lean:84`: offer both branches, then forget which was taken.	
R ∪ S = S ∪ R — `Freyd.Diag.Biprod.union_comm`. The two branches of a tape are interchangeable, exactly as the two strands of Δ are.	
⊥ ∪ R = R — `Freyd.Diag.Biprod.bot_union`. `bot a b := toZero a ; (toZero b)°:diag/Tape.lean:87`, routed through the zero object: the emptier has no shape for it, so it prints as a box.	
R (S ∪ T) = R S ∪ R T — `Freyd.Diag.Biprod.comp_union_composition`. Composition distributes over union, which is done easily: Freyd §2.2's distributivity; the reason a distributive allegory is more than an allegory.	
(R ∪ S)° = R° ∪ S° — `Freyd.Diag.Biprod.conv_union`. The converse is a ⊤-functor for the tape layer too.	
R ∩ (S ∪ T) = (R ∩ S) ∪ (R ∩ T) — `Freyd.Diag.FbCBRig.meet_union_distrib`, the law that makes `distributiveAllegoryOfFbCb` a **distributive** allegory (§2.2). The one picture with both layers in it: meets inside a tape on the left, tapes around meets on the right.	
(R / S) S ⊆ R — `Freyd.Diag.ClosedLinearBicat.residual_comp_ev`: the residual is a lower bound of the arrows it is the greatest of. Freyd's §2.31 division R / S is the largest T with T S ⊆ R, and `le_residual_iff(diag/FO.lean:178)` is that Galois statement in full. The dashed frame marks the box as belonging to the second composition.	

7. Lemma 4.2 (ii) — the proof, not just the statement

`diag-export --proof` draws a **proof**. A `calc` chain elaborates to nested `Trans.trans`, so its intermediate terms are all recoverable from the proof term: flatten the spine, infer the type of each leaf, draw canvas per step. Nothing is replayed and nothing is retold — the nine rows below are the chain the Lean kernel accepted for `Freyd.Diag.CartBicat.conv_comp`, which is Lemma 4.2 (ii), (R ; S)° = S° R°.

7.1. First: why these pictures are not pictures of a converse

They are not, and that is the thing to get straight before reading them. The converse is drawn with **two bends** — a cup on the way in and a cap on the way out; each a pair of dots, ⊤ ; Δ and ∇ ; ! — which is the left-hand figure below. The proof never draws that, `conv_unique` says an arrow is the **converse** of R as soon as it satisfies the equation with UNBENT arrows a ⊗ b → 1: two strands in on the left, one cup on the right. Bending is a bijection (`bend_unbend`, `unbend_bend`), so settling the unbent equation settles the bent one, and the cup never has to be drawn at all.

That is why every row below has a cap and no cup, and why an arrow sitting on the **bottom** strand, mirrored.

the converse, as the paper draws it	the one rule the proof uses, twice
	
R° = R with both wires bent round: in at the bottom left, up at the top left, out at the bottom right. The box is rotated(?) 180°, as `FunctorialSemanticsForRelationalTheories.pdf` draws it — a cup = ? then Δ, a cap = ∇ then !. **Transcribed**: being a definition has no statement to export.	`conv_slide (diag/CB.lean:440)`: a box on the BOTTOM strand facing a cap is the same as its mirror image on the TOP strand. The boxed is what turns R into R°. This is the entire content of Lemma 4.2 (ii); the other moves are bookkeeping.

7.2. The chain

One row per term, not per step: the right-hand side of every step is the picture twice. The relation that reaches it is in the columns the Lean has it; in the picture the text is what the Lean it is (?). What is shown is the relation to the next picture, and the rule, in which the...

Lemma 4.2 (ii). (R ; S)° ; 1 ⊗ S° ; R° = S° R°

term	picture	the rule that reaches it
1 ⊗ (S° ; R°) ; cap		**the start.** What `conv_unique` asks about: S° R°, both boxes on the bottom strand, mirrored — that is what "these are converses" looks like once bending is done.
= 1 ⊗ S° ; (1 ⊗ R° ; cap)		**1.** ⊗ is a functor, so doing two things to one strand in sequence is the sequence of doing them. Same picture.
= 1 ⊗ S° ; (R ⊗ 1 ; cap)		**2. conv_slide.** R° the box nearest the cap, so it slides round the bend: it comes out upright on the top strand, and comes back upright as R. The first of the two steps that move pictures.
= (1 ⊗ S° ; R ⊗ 1) ; cap		**3.** `simp only [Cat.assoc]`. Re-bracketing — visible in the picture. Same picture.
= R ⊗ S° ; cap		**4. tensorHom_split** — interchange. R is on the top strand and S° on the bottom, so running them in sequence and running them side by side are the same thing. Moves as a re-spacing only because the equation lays a diagram out left to right.
= (R ⊗ 1 ; 1 ⊗ S°) ; cap		**5. tensorHom_split** — the same law, back the other way. Split them apart again, R first, S° is now the box facing the cap. Which is what the next step needs.
= R ⊗ 1 ; (1 ⊗ S° ; cap)		**6.** `simp only [Cat.assoc]`. Re-bracketing. Same picture.
= R ⊗ 1 ; (S ⊗ 1 ; cap)		**7. conv_slide**, on S this time. S° slides round the bend and comes up upright on the top strand. Both boxes now top, both upright — the R and the S are in last step.
= R ⊗ 1 ; S ⊗ 1 ; cap		**8.** `simp only [Cat.assoc]`. Re-bracketing. Same picture.
= (R ; S) ⊗ 1 ; cap		**9. tensorHom_comp** — interchange once more; the two top-strand boxes merge into one. This is R ; S unbent; so `conv_unique` reads off S° R° = (R ; S)°. Done.

What the chain measures. Read down the pictures and most consecutive pairs are the same drawing — steps 1, 3, 6, 8 and 9 change nothing at all, and step 4 changes only spacing — while the term column beside them changes every time. Those are associativity, the unit laws and interchange — precisely the laws a string diagram absorbs into its geometry rather than as rewrites. Seven of these nine Lean steps are bureaucracy the picture language does not charge for. The two that move are the two applications of `conv_slide`.

How this differs from the paper's proof. The paper proves (ii) by adding in the middle of R ; S and reading the two halves off as S° and R° — three pictures, all bent. The Lean never draws a bend, because the ... the kernel accepted. Drawn by hand it would have shown the paper's form and two unrolled(?) ... the paper's proof.

8. The modular law — proved, drawn

Freyd **adjoins** the modular identity: an allegory has no structure to derive it from, so it is an axiom. Here it is a theorem, and this is the proof — `Freyd.Diag.modular_of_frobenius (diag/CB_Allegory.lean:48)`, four steps, of which exactly one step is an inequality.

the modular law, R S ∩ T ⊆ R (S ∩ R° T)

term	picture	the rule that reaches it
(R ; S) ∩ T		**the start.** R ; S next, a copy and a merge with T above and T below.
= (Δ ; (R ; T)) ; (S ⊗ 1) ; ∇		**Factor out the shared prefix.** (R ⊗ S) ∩ T = Δ ; ((R ⊗ 1) ; (S ⊗ 1)), so the copying(?) before the S can be set aside. An equality, not drawn or see.
≤ (Δ ; (R ⊗ T)) ; (S ⊗ R° ; T) ; 1		**≈V_slide_conv**, under the shared prefix. R slides through the wedge and comes up as its converse R° on the other strand. The only inequality in the proof — everything else is reshaping.
= R ; (S ∩ (R° ; T))		**Fold the prefix back in.** R ⊗ (T S°) is ≤ R (? ; (1 ⊗ S°)), and the copy-merge pair reads as a meet again. (R ; ? S°), which is the goal.

8.1. Inside the middle step

`≈V_slide_conv (diag/CB.lean:560)` is where the mathematics is, and it mentions only S — R and T sit untouched in the prefix throughout. Beyond its own it says one arrow slides through a merge:



S occurs once on the left and twice on the right. That is the tell: the lax copy inequation (42) is the only law in Def. 4.1 that may duplicate a box, so it has to be the inequality step — and its own chain shows it is the only one.

≈V_slide_conv(S ; ∇ ; ∇) ≤ (1 ⊗ S°) ; (S ⊗ 1) ; ∇ ; S

term	picture	the rule that reaches it
S ⊗ 1 ; ∇		**the start.** (S ⊗ 1) ; ∇
= (S ; (Δ ⊗ 1 ; 1 ⊗ cap)) ⊗ ...		**≈V_of_cap.** backwards. The merge unfolds into a copy and a cap, which puts the box where the copy can reach it. (Δ ⊗ 1) ; (1 ⊗ cap) is this.
≤ (Δ ; S ⊗ S) ⊗ 1 ; 1 ⊗ cap		**(42), lax_Δ.** S ; Δ becomes Δ ; (S ⊗ S): the box is copied. The whole of the content of the law, in one step.
= 1 ⊗ S° ; (S ; S) ; ∇ ; S		**≈cap_Tm_sv and conv_slide.** Reshaping back into the cap-to-converse S° — the converse is its own(?) as Lemma 4.2 (ii) in one go.

What the associators cost. ≈V and ρ have no shape in this calculus and the picture it is made of associators chain(?). Ours are the rows the stating pictures are looser than the string ... The honest price of drawing a proof in a strict(?) calculus and a hand-drawn version would have hidden them by working up to ...

9. Maps: SV, TOT, INJ, SUR, and the two inequations that always hold

Def. 4.1 states a law 4 every arrow is lax for Δ and for ! — two arrowed inequations. (42) and (43), both one direction. Equip(?) with the same statements about ∇ and ?. Those four hold for every arrow for the boxes. Give(?) ... for equipped with(?)... holding is a property of the arrow: that is (SV), (TOT), (INJ), (SUR).

	holds for every arrow	
copy Δ	(42) R ; Δ ≤ Δ ; (R ⊗ R)	
discard !	(43) R ; ! ≤ !	
merge ∇	(SV) ∇ ; R ≤ (R ⊗ R) ; ∇ — the converse of (42), not in the Lean; see below.	
unit ¡	(TOT) ¡ ; R ≥ ¡ — the converse of (43). Not in the Lean.	

The paper gives the same four properties a second time, as ADJOINT forms against `diag/CB_Derived.lean`, which are defined in the style(?) of the definitions, and there they are forward(?) to the definitions in §2.13, via a diagram map and a book map are one predicate with no translation(?).

property	adjoint form, the definition	Freyd
single valued `Freyd.Diag.SingleValued`		`Simple R := R° R ⊆ 1` §2.13
total `Freyd.Diag.Total`		`Entire R := 1 ⊆ R R°` §2.13. With simple: R is a map
injective `Freyd.Diag.Injective`		R R° ⊆ 1 §2.141 a ?-1 simple, so a ... map or a monic map
surjective `Freyd.Diag.Surjective`		1 ⊆ R° R §2.141 "g is a cover if 1 ⊆ g° g"; that is, R° is entire

Why (44) and (45) have no picture. Both are converse-images, so they exist, and it does not. Getting it means Δ° = ∇, ¡° = ? and Lemma 4.2 (iii) (if g ⊣ g° ... ⊣ g* ... f* ...) — and the first of those is a clause of Carboni & Walters' definition that Def. 4.1 as printed omits (`diag/CB.lean`'s header records this). It is the same gap that leaves Lemma 4.2 (iii) unproved in the Lean. The ones for the four reverses opposite are unaffected: they are recorded under the paper's own labels in `diag/CB_Derived.lean`, and nothing downstream depends on Lemma 4.4.

10. Entireness of an intersection

Two definitions from Freyd, both used below. The **domain** of R is the part of the identity where R is defined, `Dom R := 1 ∩ R R°` (§2.122). R is **entire** if it is defined everywhere, `Dom R = 1`, equivalently `1 ⊆ R R°` (§2.13) — which is the `Total` above, the picture opposite (47). The question this section answers is when an **intersection** is entire. Freyd's §2.124, "a lemma we will use repeatedly", and B&dM 4.18.



`Freyd.Diag.entire_inter_iff(AOP/A4_2.lean:213)` : Total (R ∩ S) ↔ 1 ≤ R ; S°

Read the two sides: on the left R ∩ S is named **twice**, on the right R and S are named **once each** and the meet is gone. That is the whole point of the lemma — the meet is needed to **ask** whether something is entire, but not to **answer** it — and it is why every later use quotes the right-hand form. Freyd gives it at §2.147, where W ∩ S ... chapter 4.

`Freyd.Diag.entire_inter_iff (AOP/A4_2.lean:187)` is the same statement for a bare allegory, and the proof is `Iff.rfl` on top of `dom_inter:in Freyd/S2_10.lean` the order X ⊆ Y is **defined** as X ∩ Y = X, so `"Dom(R ∩ S) = 1"` and `"1 ⊆ R S°"` are the same proposition once §2.124 is known. Here is a primitive, so both directions are real steps — and they cost very different amounts.

→ given R ∩ S entire, ... show ...

term	picture	the rule that reaches it
1		**the start**: 1.
≤ R ∩ S ; (R ∩ S)°		**the hypothesis**: R ∩ S entire. By the definition of entire, 1 ≤ P P° at P = R ∩ S.
≤ R ; S°		**monotonicity, twice:** R ∩ S ≤ R, and R ∩ S ≤ S under °. No modular law.

← given ... show R ∩ S entire, ...

term	picture	the rule that reaches it
1		**the start**: 1.
≤ 1 ∩ (R ; S°)		**the hypothesis**, met with 1. `meet_glb`: 1 ≤ 1 and 1 ≤ R S°, so 1 is below the meet of the two.
= 1 ∩ (R ∩ (1 ; S ; R))°		**§2.124:** 1 ∩ R S° ≤ R ∩ (S ∩ R) (S ; R°) ... S° relates R to a ... S. when some R has R and ... S. The 1 ... R's... reads "some S has R and S and S°" — ... Two arrows become one: the S is the step, and it is what made "some" that is the ... Freyd writes the right-hand Dom (S ∩ R). Dom R is the same for S ∩ R °R(?), so it is the same term, not an extra operation.
≤ R ∩ S ; (R ∩ S)°		**X ≤ X ∩ Y, Y, thus** S ∩ R = R ∩ S. The outer fork drops away, leaving the goal.

What the asymmetry says. One direction is two applications of monotonicity; the other needs §2.124, which needs the modular law, the lax copy inequation (42) — §7 above. So the picture chain locates the cost exactly: the third row of the second table is the whole strength of the allegory, and everything else is bookkeeping.

11. Anything missing?

Nothing on Freyd's side. All eight §2.11 axioms are theorems of Def. 4.1, and `allegoryOfCartBicat (diag/CB_Allegory.lean:85)` discharges the `Allegory` structure field for field. The modular identity — which Freyd has to **adjoin** — is derived.

Three things are stated but unproved. `≈Δ∘!=` and `≈!∘!=` are not drawn at all, and Lemma 4.4, that the comonoid and adjoint forms of the maps agree in Def. 4.1 are related — no Frobenius structure at a composite object — which is why `diag/CB.lean`'s header records them.

Two laws on the Frobenius side do no work here. Gripping the repo, `≈Rδu1` and `≈!∘!=` are never used by (see below); (44) and (45) are not stated as map conditions in `diag/CB.lean`. All three trace to one omission: the Δ° = ∇ instance dropped by uniqueness of adjoints — which is what they make to Δ ⊣ ∇ and ! ⊣ ¡ by uniqueness of adjoints — not to prove any property of the calculus.

The bridge runs one way only. For the reason the sections above gave: a cartesian bicategory carries a symmetric monoidal ⊗ and ⊤, but an allegory carries neither — see §5.

What Def. 4.1 alone does not reach. Everything past Freyd's §2.11 needs structure beyond it, and none of it is missing — it is built, but from different theories. `ant from FunctorialSemanticsForRelationalTheories.pdf`:

construct	Freyd	diagrammatic counterpart	status
tabulation	§2.14	none	no counterpart. Def. 4.1 gives no splitting of coreflexives; ⊤ = ! ; ¡ is all the tabular content there is.
∪, ⊥	§2.2, distributive allegory	a **biproduct** on top of Def. 4.1 — `diag/Tape.lean`, `TapeDiagrams.pdf`; union, bot, and distributive-ness; Biproduct from Def. 4.1 above	built, and drawn above. Not derivable from Def. 4.1 alone — the paper's calculus has no union at all.
R / S, complement	§2.31, division allegory	`LinearBicat, ClosedLinearBicat`; `diag/FO.lean`, `DiagrammaticAlgebraOfFirstOrderLogic.pdf`	built, and drawn above. The `Rel(Set)` instance (`diag/RelSetFO.lean`) needs `Classical.choice`.
∧, power transpose	§2.4	none	open. No calculus in this tower has a power object.
Allegory ⟹ CB	§2.15 + §2.165	`cartBicatOfUnitaryPretabular`	deliberately not built.

One caution on §7–§9 of the paper, which look like they extend to this table and the axioms. The paper says it is of ⊤ — only the black fragment ⊤, not ... while "⊥ is not the empty relation and ∪ is not the union".

12. The shunting rule

A map may be moved from one side of a containment to the other, becoming its converse on the way. This is Freyd §2.11 ... program calculation — B&dM 4.19 and 4.20 — and it is what makes program derivation rules all the same. A map is nothing more than this...

`AOP/A4_2.lean` has both rules for a bare `Allegory` (`map_shunt_right`, `map_shunt_left`), which is where every downstream derivation calls them. They are also proved in the diagrammatic layer (`shunt_right`, `map_shunt_left` in `diag/CB_Derived.lean` — same statements, so that there is a Lean diagram for the exporter to draw. Nothing new is assumed: both are proved from the Map alone.

Each rule is an ↔, so each its two pictures. The statement of a branch is its hypothesis: it is the hypothesis: the left is the goal and it reaches, both — with the map on the other side, and it is the chain proves.

The proofs, drawn. Each is two chains, and `diag-export --proof` reaches inside its `Iff.intro` into its two branches rather than drawing the longer one and dropping the other. The point to watch is ...

shunting right, → given ... show ...

term	picture	the rule that reaches it
R		**the start**: R.
≤ R ; (f ; f°)		**Total f.** 1 ≤ f f°, so f f° may be inserted after R. This is the only thing the forward direction needs.
≤ S ; f°		Re-bracket to (R f) f° and apply the hypothesis R f ≤ S. R ≤ S f°, which is the goal.

shunting right, ← given ... show ...

term	picture	the rule that reaches it
R ; f		**the start**: R f.
≤ S ; f° ; f		Apply the hypothesis R ≤ S f° on the left. The two f's become adjacent.
≤ S		**SingleValued** f° f ≤ 1, so the pair cancels and S is left. The only thing the backward direction costs.

Shunting left is the mirror, and its two chains come out the same shape — Total f on the way out, SingleValued f on the way back — with the map on the other side of R and the converse on the left, with the rule on its opposite branch.

shunting left, → given ... show ...

term	picture	the rule that reaches it
R		**the start**: R.
≤ f ; (f° ; R)		**Total f.** f f° is inserted before R.
≤ f ; S		Re-bracket to f (f° R) and apply the hypothesis f° R ≤ S.

shunting left, ← given ... show ...

term	picture	the rule that reaches it
f° ; R		**the start**: f° R.
≤ (f° ; f) ; S		Apply the hypothesis on the right. The two f's become adjacent.
≤ S		**SingleValued**. The pair cancels.

What the rule reaches for. An adjunction. R f ≤ S ⟺ R ≤ S f°: says f ↔ f° in the poset-enriched category, and the residual is the same for f as the other(?). Lemma 4.2 (iii) — if a single right adjoint, and it be f°, that is proved there. Lemma 4.4 (Freyd.Diag.lemma_4_4_total) turns it round: any map need not be ... shunting argument — an arrow with a right adjoint in an allegory is a map. It is precisely the ... is the diagram that shows the total Lemma 4.4, and this is the shape of every ...

Where it is spent. In this repo, `map_shunt_right/map_shunt_left` are called by the relational-product universal property (`AOP/A4_2.lean:37`) and the coproduct characterisation (`AOP/A5_13.lean:67`) and by the catamorphism work (`AOP/A6_?:lean:?`). It is the single most used lemma of the AOP layer.

13. Division and negation — the cheatsheet's §8, drawn

`ADP/Cheatsheet.typ`'s eighth section is B&dM §4.4–4.5: the two divisions, the symmetric division, negation, and Schröder's rule. Every one of its laws that has a picture in some repo statement — it has one(?) are Freyd §2.31(?)... and Schröder in AOP/A4_5.lean — so every one of them has a Lean statement, and the pictures say so.

What the notation says here. Dashed means "not a composite of the generators" at the other end. The `ClosedLinearBicat` (`diag/FO.lean`) – ... are what a **division** or a **boolean** allegory adds to Allegory, Chap. 2.3. The same type for a second ... box, and the diagrams... and De Morgan's are the only ones on the diagram ... so the boxes cannot be paired ... and the converse is the one ... dashed not-box.

The same caveat as §6. A dashed box hides an interior; a dashed box is not a tape inside a frame, it is not a wire. It says it is in the second composition, as a whole ... the box picture ... can(?) be ... the converse.

The residual drawn earlier is the first of / ↔ ... it is (R ; S) ≤ T ... same Galois statement ... T ≤ R / S ... and in a model carrying both divisions the adjunctions(?) hold. This is why the table ... and the residual ... each divisions both ...

statement	picture, from the Lean statement
Division is the right adjoint of composition: T ⊆ R/S ⟺ T S ⊆ R — `Freyd.Division.div_le_iff (Freyd/S2_30.lean:54)`. The cheatsheet's first, reversed, since Freyd divides on the right. Also `le_div_iff` is the Freyd adjoint of (?) ... `DivisionAllegory` with no definition.	
T ⊆ S\R ⟺ S T ⊆ R — `Freyd.Diag.le_leftDiv_iff (?)`, the mirror. `S\R = (R°/S°)°` (S 301); so the left division is computed(?) by the converse and is not a second primitive.	
cancel: (R/S) S ⊆ R — `Freyd.Diag.leftDiv_cancel(?)`, field: `DivisionAllegory.div_cancel_le (?)`; and ... (S\R) ⊆ R — `Freyd.Diag.leftDiv_comp_le (S 317)`. The value of each adjunction, the counits(?) ... with the division boxes ... inside.	
associate: R/(S T) = (R/S)/T ... — `Freyd.Diag.div_comp_assoc (S 183)`; and (S T)\R = T\(S\R) — `Freyd.Diag.leftDiv_comp(?)`. Both are shown as ... a curved(?) dashed box ... a single box ... one ... a division of an ... and the right-hand side ... to ... of the other ...	
maps: If (R/S) = (f R)/(S) ... `Freyd.Diag.div_map (S 366)`; and R/(f° S) = ... (f° = ...) — `Freyd.Diag.div_comp_map (S 376)`. Both are the shunting rule of the previous section, and here ... and the proofs are the ... a map ...	
symmetric division: (R/S) ∩ (S/R)° = S/R ... — `Freyd.Diag.symDiv_recip (Freyd/S2_30.lean:?)`, and (R/?)(S/?) ... — `Freyd.Diag.symDiv_comp (?)` R/S ∩ (S\R)° — the cheatsheet's one(?) ... convex(?) and symmetric, and in ... a converse ... of ...	

Negation is a note on R — ¬R = R ⇒ ⊥ (AOP/A4_5.lean:44) on top of a Boolean allegory, with the formulas Heyting algebra ... R ∩ ... R = R in an extra axiom, `BooleanAllegory` (AOP/A4_5.lean:123). The two rows marked boolean depend on that ... of the dividing ... the Boolean ... under the Frobenius side ... drawing the ... below.

statement	picture, from the Lean statement
R ⊆ S ⟺ R ∩ ¬S = ⊥ — `Freyd.Diag.le_iff_inter_neg_zero (AOP/A4_5.lean:144)`. The bridge every Schröder proof crosses: containment turned into a disjointness, which the complement can state. `inter_zero` can make a statement.	
R ∩ ¬R = ⊥ — `Freyd.Diag.inter_neg_zero (?)`. The one pair of a meet of a relation with its own complement ...	
De Morgan: ¬(R ∪ S) = ¬R ∩ ¬S — `Freyd.Diag.neg_union (S 158)`. The union is under the ¬ on the left, as it prints; the meet is on the right is drawn. Two equations, both conventions.	
De Morgan, other side (boolean): ¬(R ∩ S) = ¬R ∪ ¬S — `Freyd.Diag.neg_inter (S 131)`. Needs ¬¬R = R, `neg_union` ... ¬R ∪ ... the ... and ...	
excluded middle (boolean): R ∪ ¬R = ⊤ — `Freyd.Diag.union_neg_top (S 139)`. `boolean_iff(?)` ... is the ... to the ... whose ... of the fixed ...	
Schröder: R S ⊆ T ⟺ R° ¬T ⊆ ¬S ⟺ ¬T S° ⊆ ¬R — `Freyd.Diag.schroeder_left(?)`, and R S ⊆ T ⟺ ¬T S° ⊆ ¬R — `Freyd.Diag.schroeder_right (?)`. The workhorse of §4.5, and the picture that is ... the same two boxes with one ... converse, ... each of ... the ... and ...	
division through negation: ¬(R)/Y = ¬(R Y°) — `Freyd.Diag.neg_div_neg(?)` ... (R/S) ... ¬(¬R) ... `Freyd.Diag.div_eq_neg_comp (S 293)`, which ... to ... The cheatsheet's "¬ eq" ... a division ... here, ... the ... form ... of ...	
lexical order: R ; S ⊆ R ∩ ... — `Freyd.Diag.topHom_trans(?) (AOP/A4_7_Galois.lean:304)`. R ; S ⊆ R ∩ (R° ; S) ... — compare ... lexicographic ... the ...	
Galois, point-free: f(x) ≤ y ⟺ x ≤ g(y) is R f° = g S — `Freyd.Diag.RelSet.galoisPF (AOP/A4_7_Galois.lean:36)`, `Freyd.Diag.galois_iff (?)` (S 188), proving it equivalent to the pointwise form. ... the `Rel(Set)` ... a ... and ... the picture is the point-free form's whole advantage — two composites, no quantifiers.	

The measurement. Twenty-one pictures, and a generator appears in the in one spot in the `le_iff_inter_neg_zero`, `inter_neg_zero` and `neg_union_neg_top`. The other sisters are boxes threaded on a wire. That composition are what the Frobenius generators build, and §4.4–4.5 is a pass at the Frobenius calculus.

What is not drawn, and why. The cheatsheet's pointwise row, x ⊆ (T\U) ..., x ↦ ..., x ⊆ y ..., quantifies over elements — definition to depict in point-free ... and there is the same ... The other ones are plain ... `zipPF`, ... the second ...