

<sup>1</sup>  
<sup>2</sup> Susceptible host dynamics explain pathogen resilience  
<sup>3</sup>

<sup>4</sup> Sang Woo Park, . . . , Sarah Cobey

## <sup>5</sup> **Abstract**

<sup>6</sup> Major priority for epidemiological research in the time of anthropogenic change is  
<sup>7</sup> understanding how infectious disease dynamics respond to perturbations. Interven-  
<sup>8</sup> tions to slow the spread of COVID-19 significantly disrupted the transmission of  
<sup>9</sup> other human pathogens, providing unique opportunities to learn about pathogen  
<sup>10</sup> characteristics from spatiotemporal variation in re-emergence patterns. As inter-  
<sup>11</sup> ventions lifted, a key question of whether and when respiratory pathogens would  
<sup>12</sup> eventually return to their pre-pandemic dynamics remains to be answered. To ad-  
<sup>13</sup> dress this gap, we develop a framework for estimating pathogen resilience based on  
<sup>14</sup> how fast epidemic patterns return to their pre-pandemic, endemic dynamics. Our  
<sup>15</sup> analysis reveals a possibility that some pathogens may have settled to endemic cycles  
<sup>16</sup> that are different from their pre-pandemic patterns. Finally, we show that the re-  
<sup>17</sup>plenishment rate of the susceptible pool is a key determinant of pathogen resilience.  
<sup>18</sup> Our framework offers a novel perspective to characterizing the dynamics of endemic  
<sup>19</sup> pathogens and their responses to COVID-19 interventions.

Understanding how ecological systems respond to perturbations is a fundamental challenge in predicting species persistence and extinction [1, 2, 3]. These responses can be characterized in terms of resilience, which often measures how fast a system returns to its stable, reference state following a perturbation [4, 5, 6, 7]. Both theoretical and empirical efforts to quantify resilience of ecological systems have provided key insights for understanding the dynamics of complex systems and linking these findings to actionable strategies for species conservation [8]. However, despite rich literature on ecological resilience, there have been limited applications to measuring the resilience of host-pathogen systems, especially for human pathogens.

Non-pharmaceutical interventions (NPIs) to slow the spread of COVID-19 disrupted the transmission of other human pathogens, providing large-scale natural experiments for understanding how various host-pathogen systems respond to perturbations [9, 10, 11, 12]. In particular, as interventions lifted, large heterogeneities in outbreak dynamics were observed across different pathogens in different countries (Figure 1), likely reflecting differences in NPI patterns, pathogen characteristics, immigration/importation from other countries, and pre-pandemic pathogen dynamics [13]. Even though more than four years have already passed since the emergence of COVID-19, current circulation patterns for many respiratory pathogens appear to be different from their pre-pandemic, seasonal patterns, especially in Hong Kong and Korea: some pathogens, such as human metapneumovirus and bocavirus in Korea, are circulating at lower levels, whereas other pathogens, such as RSV in Korea, seem to exhibit different seasonality (Figure 1). These observations pose two fundamental questions for current and future infectious disease dynamics: (1) can we learn about underlying pathogen characteristics, such as their transmissibility or duration of immunity, from re-emergence patterns? and (2) can we predict whether and when other respiratory pathogens will eventually return to their pre-pandemic dynamics?

To address this question, we propose a framework for characterizing the resilience of a host-pathogen system based on how fast the system recovers from perturbation. We begin by laying out a few representative scenarios that capture the potential impact of COVID-19 interventions on endemic pathogen dynamics and illustrating how resilience can be measured by comparing the pre- and post-pandemic dynamics of susceptible and infected hosts. In practice, information on susceptible hosts are often unavailable, and traditional methods for reconstructing the dynamics of susceptible hosts require long-term endemic time series [14, 15], which cannot be applied due to disruptions in epidemic patterns caused by COVID-19 interventions. Instead, we utilize Takens' embedding theorem to reconstruct empirical attractors from data and further measure the distance from this empirical attractor [16]. This reconstruction allows us to characterize the rate at which this distance decreases over time, which correspond to pathogen resilience. We apply this framework to analyzing pathogen surveillance data for a wide array of respiratory and non-respiratory pathogens from Canada, Hong Kong, Korea, and US. Finally, we show that susceptible host dynamics are a key determinants of pathogen resilience. Our study offers unique insights into understanding pathogen re-emergence patterns following COVID-19 interventions.




Figure 1: **Observed heterogeneity in responses to COVID-19 pandemic across respiratory pathogens and norovirus in (A) Canada, (B) Hong Kong, (C) Korea, and (D) US.** Red points and lines represent data before 2020. Blue points and lines represent data since 2020. Gray lines and shaded regions represent the mean seasonal patterns and corresponding 95% confidence intervals based on the observed outbreak patterns before 2020.

## 63 Conceptual introduction to pathogen resilience

64 In classical ecological literature, resilience of an ecological system is measured by  
 65 the rate at which the system returns to its reference state following a perturbation  
 66 [4, 5, 6, 7]. This rate corresponds to the largest real part of the eigenvalues of the  
 67 linearized system near equilibrium—here, we refer to this value as the *intrinsic* re-  
 68 silience of the system, which represents the expected rate of return from perturbated  
 69 states. However, respiratory pathogens often exhibit seasonal variation in transmis-  
 70 sion, meaning that the intrinsic resilience of a host-pathogen system varies across  
 71 season. Nonetheless, we can still measure the *empirical* resilience of a host-pathogen  
 72 system by looking at how fast the system returns to the pre-pandemic, endemic  
 73 dynamics after interventions are lifted.

74 As an example, consider an intervention that reduce transmission by 50% for 6  
 75 months starting in 2020, which causes epidemic patterns to deviate from its original  
 76 stable annual cycle for a short period of time and eventually come back (Figure 2A).  
 77 To measure the empirical resilience of this system, we first need to be able to measure

78 the distance from its pre-pandemic attractor. There are many different ways we can  
79 measure the distance from attractor, but for illustrative purposes, we choose one of  
80 the most parsimonious approach: that is, we look at how the susceptible (S) and  
81 infected (I) populations change over time and measure the distance on the SI phase  
82 plane (Figure 2B). In this simple case, the locally estimated scatterplot smoothing  
83 (LOESS) fit indicates that the distance from attractor decreases linearly on average  
84 (Figure 2C). Furthermore, the overall rate of return matches the intrinsic resilience  
85 of the seasonally unforced system (Figure 2C).

86 Alternatively, NPIs can permanently change our behavior and have persisting  
87 impact on the pathogen dynamics; as an example, we consider a scenario in which a  
88 10% reduction in transmission persists even after the NPIs are lifted (Figure 2D–F).  
89 In such cases, we cannot know whether the pathogen will return to its original cycle  
90 or a different cycle until many years have passed after the NPIs are lifted, meaning  
91 that we cannot measure the distance against the new attractor that the system will  
92 eventually approach. Nonetheless, we can still measure the distance against the orig-  
93 inal, pre-pandemic attractor and ask how the distance changes over time (Figure 2E).  
94 The LOESS fit suggests that the distance from the attractor will initially decrease  
95 exponentially on average (equivalently, linearly on a log scale) and eventually plateau  
96 (Figure 2F). Here, a permanent 10% reduction in transmission rate slows down the  
97 system, which causes the distance from the attractor to decrease at a slower rate  
98 (Figure 2F) than it would have otherwise in the absence of permanent transmission  
99 reduction (Figure 2C). This example shows that resilience is not necessarily an in-  
100 trinsic property of a specific pathogen. Instead, pathogen resilience is a property of  
101 a specific attractor that a host-pathogen system approaches, which depends on both  
102 pathogen and host characteristics.

103 Finally, transient phenomena can also complicate the picture (Figure 2G–I). For  
104 example, a stage-structured model for RSV initially exhibits a stable annual cycle,  
105 but perturbations from NPIs cause the epidemic to exhibit biennial cycles (Figure  
106 2G). Despite this biennial cycle, we see that the system eventually approaches the  
107 original pre-pandemic attractor (Figure 2H), suggesting that this biennial cycle is a  
108 transient phenomenon. The LOESS fit indicates that the distance from the attractor  
109 will initially decrease exponentially at a rate that is consistent with the intrinsic  
110 resilience of the seasonally unforced system, but the rate of decrease slows down  
111 as the epidemic exhibits a biennial cycle (Figure 2I). In classical ecological theory,  
112 this behavior is also referred to as a ghost attractor, which causes long transient  
113 dynamics and slow transitions [17]. As we show in Supplementary Figure S1, strong  
114 seasonal forcing in transmission can also lead to transient phenomena for a simple  
115 SIRS model, causing a slowing down of the system.

116 In Supplementary Materials, we also explore measuring the resilience of a two-  
117 strain host-pathogen system: when the dynamics two strains (or two pathogens) are  
118 coupled through cross immunity, we would expect the entire system to be character-  
119 ized by a single resilience value (rather than having two separate resilience for each  
120 strain). Simulations from a simple two-strain system illustrate that separate anal-




Figure 2: **Conceptual framework for measuring pathogen resilience following NPIs across different scenarios.** (A, D, G) Simulated epidemic trajectories across various models. Red and blue solid lines represent epidemic dynamics before and after interventions are introduced, respectively. Red dashed lines represent counterfactual epidemic dynamics in the absence of interventions. Gray regions indicate the duration of interventions. (B, E, H) Phase plane representation of the corresponding model. Red and blue solid lines represent epidemic trajectories on an SI phase plane before and after interventions are introduced, respectively. (C, F, I) Changes in logged distance from attractor over time. Blue lines represent the logged distance from attractor. Orange lines represent the locally estimated scatterplot smoothing (LOESS) fits to the logged distance from attractor. Dotted lines are superimposed as a comparison to have the same slope as the intrinsic resilience of the seasonally unforced system.

yses of individual strain dynamics (e.g., RSV A vs B) and a joint analysis of total

122 infections (e.g., total RSV infections) yield identical resilience estimates, confirming  
123 our expectation (Supplementary Figure S2, 3). Analogous to a single system, strong  
124 seasonal forcing in transmission can cause the system to slow down through transient  
125 phenomena (Supplementary Figure S4).

126 These observations indicate three possibilities. First, we can directly estimate  
127 the empirical resilience of a host-pathogen system by looking at how fast the system  
128 approaches a pre-pandemic attractor, provided that we can measure the distance  
129 from attractor. The empirical approach to estimating pathogen resilience is partic-  
130 ularly convenient because it does not require us to know the true underlying model.  
131 As we show in Supplementary Figure S5, estimating the intrinsic resilience from fit-  
132 ting standard compartmental models can lead to biased estimates, especially under  
133 model misspecification. Second, resilience estimates allow us to make phenomenolog-  
134 ical predictions about the dynamics of a host-pathogen system following a perturba-  
135 tion: assuming that the distance from the attractor will decrease exponentially over  
136 time, we can obtain a ballpark estimate for when the system will reach an attractor.  
137 Finally, deviation from an exponential decrease in the distance from attractor can  
138 provide information about whether the system has reached an alternative attractor,  
139 or a ghost attractor, that is different from the original, pre-pandemic attractor. These  
140 alternative attractors may reflect continued perturbations from permanent changes  
141 in transmission patterns as well as changes in immune landscapes.

## 142 Inferring pathogen resilience from real data

143 Based on these observations, we now set out to infer pathogen resilience from real  
144 data. Here, we briefly lay out our approach to estimating pathogen resilience from  
145 real data (Figure 3). We then test this approach against simulations and apply it to  
146 real data.


147 So far, we focused on simple examples that assume a constant transmission re-  
148 duction. However, in practice, the impact of NPIs on pathogen transmission is  
149 likely more complex (Figure 3A), reflecting introduction and relaxation of various  
150 intervention strategies. These complexities can lead to longer delays between the  
151 introduction of NPIs and pathogen re-emergence as well as temporal variation in  
152 outbreak sizes (Figure 3B): in this example, continued transmission reduction from  
153 NPIs limits the size of the first outbreak in 2021 following the emergence, allowing  
154 for a larger outbreak in 2022 when NPIs are further relaxed.

155 Previously, we relied on the dynamics of susceptible and infected hosts to compute  
156 the distance from attractor (Figure 2), but information on susceptible hosts are  
157 often not available in practice. In addition, uncertainties in case counts due to  
158 observation error as well as the possibility of complex, multiannual attractor adds  
159 challenges to measuring the distance from attractor. To address these challenges, we  
160 first reconstruct an empirical attractor by utilizing Takens' theorem, which states  
161 that an attractor of a nonlinear multidimensional system can be mapped onto a

162 delayed embedding [16]. Here, we use delayed copies of logged values of pre-pandemic  
 163 cases  $C(t)$  (Figure 3C) to reconstruct the attractor:

$$\langle \log(C(t) + 1), \log(C(t - \tau) + 1), \dots, \log(C(t - (M - 1)\tau) + 1) \rangle, \quad (1)$$

164 where the delay  $\tau$  and embedding dimension  $M$  are determined based on autocorrelations  
 165 and false nearest neighbors, respectively [18, 19]. We then apply the same delay  
 166 and embedding dimensions to the entire time series to determine the position on a  
 167 multi-dimensional state space (Figure 3D), which allows us to measure the nearest  
 168 neighbor distance between the current state of the system and the empirical attractor  
 169 (Figure 3E). In principle, we can quantify how fast this distance decreases by fitting  
 170 a linear regression on a log scale, where the slope of the linear regression corresponds  
 171 to pathogen resilience. As we show in Supplementary Figure S6, overall temporal  
 172 variations in the distance from attractor, especially the observed rate of decrease,  
 173 appear robust to choices about embedding delays and dimensions; we note that using  
 174 longer delays and higher dimensions tend to smooth out temporal variations in the  
 175 distance from attractor.



**Figure 3: A schematic diagram explaining how pathogen resilience can be inferred from real data.** (A) A realistic example of a synthetic NPI, represented by a relative reduction in transmission. (B) The impact of the synthetic NPI on epidemic dynamics simulated using a stochastic SIRS model. (C) Generating delayed copies of the logged time series allows us to obtain an embedding. (D) Two dimensional representation of an embedding. (E) Delayed embedding allows us to calculate the nearest neighbor distance from the empirical attractor, which is determined based on the pre-pandemic time series. This distance time series can be used to infer pathogen resilience by fitting a linear regression after choosing an appropriate window for regression.

176 Complex changes in the distance from attractor suggest that estimating pathogen  
 177 resilience from linear regression will likely be sensitive to our choice of fitting windows  
 178 for the regression. In Supplementary Materials, we explore an automated window  
 179 selection criteria for linear regression and test it against randomized, stochastic sim-  
 180 ulations across a wide range of realistic NPI shapes. We find that resilience estimates

181 based on the automated window selection criteria are moderately correlated (0.54)  
182 with the intrinsic resilience of the post-NPI attractor (Supplementary Figure S7). In  
183 contrast, a naive approach that uses the entire time series, starting from the peak  
184 distance, only gives a correlation of 0.21 and consistently underestimates the intrinsic  
185 resilience (Supplementary Figure S7).

186 Now, we apply this approach to pathogen surveillance data presented in Figure  
187 1. For each time series, we apply Takens' theorem independently to reconstruct the  
188 empirical attractor and obtain the corresponding time series of distance from attrac-  
189 tors (Supplementary Figure S8 for the distance time series and linear regression fits).  
190 Then, we use the automated window selection criteria to fit a linear regression and  
191 estimate the empirical resilience for each pathogen in each country. For most res-  
192 piratory pathogens, resilience estimates fall between 0.4/year and 1.8/year (Figure  
193 4A), with the exception of Rhinovirus in the US (0.066/year; 95% CI: 0.018/year–  
194 0.113/year) and Bocavirus in Korea (0.087/year; 95% CI: 0.023/year–0.151/year).  
195 Excluding these exceptions, the mean resilience of common respiratory pathogens is  
196 0.974/year (95% CI: 0.784/year–1.16/year). As a reference, this is  $\approx$  7 times higher  
197 than the intrinsic resilience of pre-vaccination measles dynamics ( $\approx$  0.13/year). Fi-  
198 nally, resilience estimates for norovirus appears to be comparable to the intrinsic  
199 resilience of measles: 0.119/year (95%CI: 0.004/year–0.233/year) for Korea and  
200 0.385/year (95% CI: 0.167/year–0.603/year). A simple ANOVA shows that there  
201 are significant differences in resilience estimates across countries ( $p < 0.036$ ) and  
202 pathogens ( $p < 0.030$ ).

203 Using resilience estimates, we now predict when each pathogen will return to  
204 their original pre-pandemic cycles. Specifically, we extend our linear regression fits  
205 to distance-from-attractor time series and ask when the predicted regression line  
206 will cross a threshold value, which we set to a mean of pre-pandemic distances. We  
207 predict that a return to pre-pandemic cycles would be imminent for most pathogens  
208 (Figure 4B; see Supplementary Figure S9 for a zoomed out version). In addition, we  
209 also predict that many pathogens should have already returned to their pre-pandemic  
210 dynamics by the end of 2024; but these predictions contradict some of the observed  
211 pathogen dynaics. For example, we predict that both human metapneumovirus and  
212 RSV in Korea should have returned to their attractors by now, but the magnitude  
213 and timing of recent epidemics are different from pre-pandemic patterns (Figure 1).  
214 These observations suggest the possibility that some common respiratory pathogens  
215 may have converged to different attractors.

216 In Supplementary Materials, we also consider using a lower threshold for the false  
217 nearest neighbor approach when determining the embedding dimension; this gives  
218 a higher embedding dimension. As explained earlier (Supplementary Figure S6),  
219 this gives a smoother distance-from-attractor time series (compare Supplementary  
220 Figure S10 with S8); this also requires us to use longer time series, which prevents  
221 us from estimating resilience for some pathogens. Overall, resulting estimates of  
222 pathogen resilience with higher embedding dimensions still fall between 0.3/year  
223 and 2.1/year for the most part (Supplementary Figure S11). A direct comparison




Figure 4: **Summary of resilience estimates.** (A) Estimated pathogen resilience. The gray horizontal line represents the intrinsic resilience of pre-vaccination measles dynamics. (B) Predicted timing of when each pathogen will return to their pre-pandemic cycles. The dashed line in panel B indicates the end of 2024 (current observation time). Error bars represent 95% confidence intervals.

<sup>224</sup> between two approaches (i.e., original estimate vs using higher dimensions) shows a  
<sup>225</sup> strong consistency in resilience estimates (Supplementary Figure S12).

## <sup>226</sup> Susceptible host dynamics explain variation in pathogen <sup>227</sup> resilience

<sup>228</sup> So far, we focused on quantifying pathogen resilience from the observed patterns of  
<sup>229</sup> pathogen re-emergence following COVID-19 interventions. But what factors deter-  
<sup>230</sup> mine how resilient a host-pathogen system is? Here, we use a standard Susceptible-  
<sup>231</sup> Infected-Recovered-Susceptible (SIRS) model to show that susceptible host dynamics

232 are the key determinants of pathogen resilience. To do so, we vary the basic reproduction  
 233 number  $\mathcal{R}_0$ , which represents the average number of secondary infections caused  
 234 by a newly infected individual in a fully susceptible population, and the duration of  
 235 immunity and compute intrinsic resilience for each parameter.




Figure 5: **Linking pathogen resilience to epidemiological parameters and susceptible host dynamics.** (A) The heat map represents intrinsic resilience as a function of the basic reproduction number  $\mathcal{R}_0$  and the duration of immunity. (B) The heat map represents per-capita susceptible replenishment rate as a function of the basic reproduction number  $\mathcal{R}_0$  and the duration of immunity. The standard SIRS model is used to compute intrinsic resilience and per-capita susceptible replenishment rate. Lines correspond to a set of parameters that are consistent with mean resilience estimates for each pathogen. Pathogens are ranked based on their mean resilience estimates, averaged across different countries.

236 We find an increase in  $\mathcal{R}_0$  and a decrease in duration of immunity correspond  
 237 to an increase in pathogen resilience (Figure 5A). These variations can be under-  
 238 stood in terms of the susceptible host dynamics, where faster per-capita susceptible  
 239 replenishment rate causes the system to be more resilient (Figure 5B). This rate can  
 240 be expressed as a ratio between absolute rate at which new susceptibles enter the  
 241 population and the equilibrium number of susceptible individuals in the population,  
 242  $\bar{S}$ . Therefore, both higher  $\mathcal{R}_0$  and shorter duration of immunity can drive faster  
 243 per-capita susceptible replenishment rate (Figure 5B), especially because higher  $\mathcal{R}_0$   
 244 leads to lower  $\bar{S}$ .

245 Finally, we can now rank different pathogens based on the average values of em-  
 246 pirical resilience, which allows us to determine a set of parameters that are consistent  
 247 with the estimated resilience (Figure 5A). Across all pathogens we consider, except  
 248 for bocavirus and norovirus, we estimate that the average duration of immunity is  
 249 likely to be short ( $< 6$  years) across a plausible range of  $\mathcal{R}_0$ . These rankings further

allow us to map each pathogen onto a set of parameters that are consistent with its empirical resilience (Figure 5A) and obtain a plausible range of susceptible replenishment rates for each pathogen (Figure 5B). However, we note that there is no one-to-one correspondance between susceptible replenishment rates and pathogen resilience, leading to a wide uncertainty in the estimates for susceptible replenishment rates (Figure 5B).

## Discussion

The COVID-19 interventions have caused major disruptions to circulation patterns of both respiratory and non-respiratory pathogens, adding challenges to predicting their future dynamics [9, 10, 11, 12]. On the other hand, these interventions offer large-scale natural experiments for understanding how different pathogens respond to perturbations. In this study, we show that pathogen re-emergence patterns following COVID-19 interventions can be characterized through the lens of ecological resilience. Traditionally, ecological resilience measures how fast a system returns to a reference state following a perturbation. In the context of respiratory pathogens, resilience measures how fast epidemics return to their endemic cycles after interventions are lifted.

We use an attractor reconstruction approach to quantify how distance from attractor changes over time for each pathogen [16]. We show that the resilience of a host-pathogen system can be estimated by fitting a linear regression to a logged distance-from-attractor time series. Overall, we estimate that the resilience for most common respiratory pathogens ranges between 0.4/year and 1.8/year, which is 3–14 times more resilient than prevaccination measles, indicating potential challenges in controlling common respiratory pathogens.

Our framework allows us to make phenomenological predictions about when each pathogen will return to their endemic cycles. The ability to predict future epidemic patterns from resilience estimates offers a new paradigm for epidemic forecasting. While this approach cannot predict the exact timing of outbreaks or epidemic patterns, it is nonetheless useful for predicting when epidemics will settle down to regular cycles after a large perturbation, such as COVID-19 interventions.

Our analyses suggest a possibility that several pathogens may have converged to different endemic cycles compared to their pre-pandemic epidemic patterns. Key examples include human metapnuemovirus, RSV, and bocavirus in Korea as well as RSV in Hong Kong. These changes may reflect changes in surveillance or actual shift in the dynamics, caused by permanent changes in behavior or population-level immunity. While it seems unlikely that permanent changes in behavior would only affect a few pathogens and not others, we cannot rule out this possibility given heterogeneity in the age of infection across different respiratory pathogens [SWP: CITE]. A shift in population-level immunity is plausible, as the emergence of SARS-CoV-2 and extinction of influenza B/Yamagata likely caused major changes in immune landscapes;

290 interactions among co-circulating pathogens, such as cross immunity between RSV  
291 and HMPV [20], may have also contributed to changes in population-level immu-  
292 nity. However, we currently do not know how immunity, or lack thereof, from these  
293 pathogens would affect infection from other pathogens. Future studies should use  
294 detailed mechanistic models, coupled with behavioral and immunological data, to  
295 test these hypotheses and better understand post-pandemic dynamics of endemic  
296 pathogens.

297 We show that susceptible host dynamics shape pathogen resilience, where faster  
298 replenishment of the susceptible population causes the pathogen to be more resilient.  
299 For simplicity, we focus on waning immunity and birth as a main driver of the suscep-  
300 tible host dynamics but other mechanisms can also contribute to the replenishment  
301 of the susceptible population. In particular, pathogen evolution, especially the emer-  
302 gence of antigenically novel strains, can cause effective waning of immunity in the  
303 population; therefore, we hypothesize that faster rates of antigenic evolution can also  
304 cause a pathogen to be more resilient. Future studies should explore the relationship  
305 between the rate of evolution and resilience for antigenically evolving pathogens.

306 Quantifying pathogen resilience also offers novel approaches to validating population-  
307 level epidemiological models. So far, the majority of model validation in epidemiology  
308 is based on the ability of a model to reproduce the observed epidemic dynamics and  
309 to predict future dynamics [21, 20, 22, 23, 24]. However, there can be plethora of  
310 models that meet these criteria. For example, two major RSV models have been pro-  
311 posed so far to explain biennial epidemic patterns: (1) a stage- and age-structured  
312 model that allows for disease severity to vary with number of past infections and  
313 age of infection [22] and (2) a pathogen-interaction model that accounts for cross  
314 immunity between RSV and human metapneumovirus [20]. Since both models can  
315 accurately reproduce the observed epidemic patterns, standard criteria for model  
316 validation do not allow us to distinguish between these two models from population-  
317 level data alone. Instead, we can measure the empirical resilience of each model  
318 by simulating various perturbations and compare them to estimates of empirical re-  
319 silience from data, using COVID-19 interventions as an opportunity. Future studies  
320 should further investigate using pathogen resilience for validating epidemic models.

321 There are several limitations to our work. First of all, we did not extensively ex-  
322 plore other approaches to reconstructing the attractor. Recent studies showed that  
323 more sophisticated approaches, such as using non-uniform embedding, can provide  
324 more robust reconstruction for noisy data [19]. In the context of causal inference,  
325 choices about embedding can have major impact on the resulting inference [25]. Our  
326 resilience estimates are likely overly confident given a lack of uncertainties in at-  
327 tractr reconstruction as well as the simplicity of our statistical framework. Short  
328 pre-pandemic time series also contributes to the crudeness of our estimates. Nonethe-  
329 less, as illustrated in our sensitivity analyses (Supplementary Figure S6, S10–S12),  
330 inferences about pathogen resilience appear to be robust to decisions about embed-  
331 ding lags and dimensions—this is because tracking the rate at which the system  
332 approaches the attractor is likely a much simpler problem than making inferences

333 about causal directionality. Our qualitative prediction that common respiratory  
334 pathogens are more resilient than prevaccination measles is also likely to be robust  
335 to these predictions, given how rapid many respiratory pathogens returned to their  
336 original cycles following COVID-19 interventions.

337 Predicting the impact of anthropogenic changes on infectious disease dynamics  
338 is a fundamental aim of infectious disease research in a rapidly changing world.  
339 Our study illustrates that quantifying pathogen resilience can help us understand  
340 how infectious disease pathogens respond to major perturbations caused by public  
341 health interventions. More broadly, a detailed understanding of the determinants of  
342 pathogen resilience may offer unique insights into pathogen persistence and control-  
343 lability.

## 344 Materials and Methods

### 345 Data

346 We gathered time series on respiratory infections from four different countries: Canada,  
347 Hong Kong, Korea, and United States (US). As a reference, we also included time  
348 series data on norovirus infections for available countries—in contrast to respiratory  
349 pathogens, we expect gastrointestinal viruses, such as norovirus, to be less affected  
350 by COVID-19 intervention measures.

351 Weekly time series of respiratory infection cases in Canada come from the Res-  
352piratory Virus Detection Surveillance System, which collect data from select labo-  
353ratories across Canada. We extracted the data from <https://www.canada.ca/en/public-health/services/surveillance/respiratory-virus-detections-canada.html>.

356 Weekly time series of respiratory infection cases in Hong Kong came from the  
357 Centre for Health Protection, Department of Health. We extracted the data from  
358 <https://www.chp.gov.hk/en/statistics/data/10/641/642/2274.html>.

359 Weekly time series of respiratory infection cases in Korea came from Korea Dis-  
360ease Control and Prevention Agency. We extracted the data from <https://dportal.kdca.go.kr/pot/is/st/ari.do>.

362 Finally, weekly time series of respiratory infection cases in the US comes from  
363 the National Respiratory and Enteric Virus Surveillance System.

### 364 Empirical attractor reconstruction

### 365 Linear regression

### 366 Mathematical modeling

367 Throughout the paper, we use a series of mathematical models to illustrate the con-  
368 cept of pathogen resilience and to understand the determinants of pathogen resilience.

369 In general, the intrinsic resilience for a given system is given by the largest real part  
 370 of the eigenvalues of the linearized system at endemic equilibrium. Here, we focus on  
 371 the SIRS model and present the details of other models in Supplementary Materials.  
 372 The SIRS (Susceptible-Infected-Recovered-Susceptible) model is the simplest model  
 373 that allows for waning of immunity, where recovered (immune) individuals are as-  
 374 sumed to become fully susceptible after an average of  $1/\delta$  time period. The dynamics  
 375 of the SIRS model is described by the following set of differential equations:

$$\frac{dS}{dt} = \mu - \beta(t)SI - \mu S + \delta R \quad (2)$$

$$\frac{dI}{dt} = \beta(t)SI - (\gamma + \mu)I \quad (3)$$

$$\frac{dR}{dt} = \gamma I - (\delta + \mu)R \quad (4)$$

(5)

376 where  $\mu$  represents the birth/death rate,  $\beta(t)$  represents the time-varying trans-  
 377 mission rate, and  $\gamma$  represents the recovery rate. The basic reproduction number  
 378  $\mathcal{R}_0 = \beta(t)/(\gamma + \mu)$  is defined as the average number of secondary infections caused  
 379 by a single infected individual in a fully susceptible population and measures the  
 380 intrinsic transmissibility of a pathogen.

381 For this model, the intrinsic resilience corresponds to:

$$-\frac{\text{Re}(\lambda)}{2} = \frac{\delta + \beta I^* + \mu}{2}. \quad (6)$$

382 Here,  $I^*$  represents the prevalence values at endemic equilibrium:

$$I^* = \frac{(\delta + \mu)(\beta - (\gamma + \mu))}{\beta(\delta + \gamma + \mu)}. \quad (7)$$

383 The susceptible replenishment rate is given by:

$$S_{\text{replenish}} = \frac{\mu + \delta R}{S^*}, \quad (8)$$


384 where  $S^* = 1/\mathcal{R}_0$  represents the equilibrium proportion of susceptible individuals.

385 When infection provides life-long immunity ( $\delta \rightarrow 0$ ), the SIRS model becomes  
 386 the SIR model. In this case, the intrinsic resilience is inversely proportional to the


387 In illustrating the impact of

<sup>388</sup> **Supplementary Text**


## Supplementary Figures




**Figure S1: Impact of seasonal transmission on pathogen resilience.** (A, D, G) Simulated epidemic trajectories using the SIRS model without seasonal forcing (A), with seasonal forcing of amplitude of 0.2 (D), and with seasonal forcing of amplitude of 0.2 (G). Red and blue solid lines represent epidemic dynamics before and after interventions are introduced, respectively. Red dashed lines represent counterfactual epidemic dynamics in the absence of interventions. Gray regions indicate the duration of interventions. (B, E, H) Phase plane representation of the corresponding model. Red and blue solid lines represent epidemic trajectories on an SI phase plane before and after interventions are introduced, respectively. (C, F, I) Changes in logged distance from attractor over time. Blue lines represent the logged distance from attractor. Orange lines represent the locally estimated scatterplot smoothing (LOESS) fits to the logged distance from attractor. Dotted lines are superimposed as a comparison to have the same slope as the intrinsic resilience of the seasonally unforced system.




**Figure S2: Conceptual framework for measuring pathogen resilience following NPIs for a multi-strain system without seasonal forcing.** (A, D, G) Simulated epidemic trajectories using a multi-strain system without seasonal forcing. Red and blue solid lines represent epidemic dynamics before and after interventions are introduced, respectively. Red dashed lines represent counterfactual epidemic dynamics in the absence of interventions. Gray regions indicate the duration of interventions. (B, E, H) Phase plane representation of the corresponding model. Red and blue solid lines represent epidemic trajectories on an SI phase plane before and after interventions are introduced, respectively. (C, F, I) Changes in logged distance from attractor over time. Blue lines represent the logged distance from attractor. Orange lines represent the locally estimated scatterplot smoothing (LOESS) fits to the logged distance from attractor. Dotted lines are superimposed as a comparison to have the same slope as the intrinsic resilience of the seasonally unforced system.




**Figure S3: Conceptual framework for measuring pathogen resilience following NPIs for a multi-strain system with seasonal forcing.** (A, D, G) Simulated epidemic trajectories using a multi-strain system with seasonal forcing (amplitude of 0.2). Red and blue solid lines represent epidemic dynamics before and after interventions are introduced, respectively. Red dashed lines represent counterfactual epidemic dynamics in the absence of interventions. Gray regions indicate the duration of interventions. (B, E, H) Phase plane representation of the corresponding model. Red and blue solid lines represent epidemic trajectories on an SI phase plane before and after interventions are introduced, respectively. (C, F, I) Changes in logged distance from attractor over time. Blue lines represent the logged distance from attractor. Orange lines represent the locally estimated scatterplot smoothing (LOESS) fits to the logged distance from attractor. Dotted lines are superimposed as a comparison to have the same slope as the intrinsic resilience of the seasonally unforced system.




**Figure S4: Conceptual framework for measuring pathogen resilience following NPIs for a multi-strain system with strong seasonal forcing.** (A, D, G) Simulated epidemic trajectories using a multi-strain system with seasonal forcing (amplitude of 0.4). Red and blue solid lines represent epidemic dynamics before and after interventions are introduced, respectively. Red dashed lines represent counterfactual epidemic dynamics in the absence of interventions. Gray regions indicate the duration of interventions. (B, E, H) Phase plane representation of the corresponding model. Red and blue solid lines represent epidemic trajectories on an SI phase plane before and after interventions are introduced, respectively. (C, F, I) Changes in logged distance from attractor over time. Blue lines represent the logged distance from attractor. Orange lines represent the locally estimated scatterplot smoothing (LOESS) fits to the logged distance from attractor. Dotted lines are superimposed as a comparison to have the same slope as the intrinsic resilience of the seasonally unforced system.




**Figure S5: Mechanistic model fits to simulated data and inferred intrinsic resilience.** (A, D, G, J) Simulated case time series from corresponding models (red) and SIRS model fits (black). (B, E, H, K) Assumed changes in transmission due to COVID-19 interventions (red) and estimated changes from the SIRS model (black). Solid lines and shaded regions represent fitted posterior median and 95% credible intervals. (C, F, I, L) True intrinsic resilience of the seasonally unforced system (vertical lines) and the posterior distribution of the inferred intrinsic resilience from the SIRS model (density plots).



**Figure S6: Sensitivity of the distance from attractor to choices about embedding lags and dimensions.** Sensitivity analysis for the distance-from-attractor time series shown in Figure 3E in the main text by varying the embedding lag between 10–15 days and embedding dimensions between 2–6 dimensions.



**Figure S7: Impact of fitting window selection on the estimation of empirical resilience.** We simulate 500 epidemics using stochastic SIRS model with randomly drawn parameters and randomly generated intervention impacts. For each simulation, we reconstruct the empirical attractor based on the approach outlined in Figure 3 and estimate the distance from attractor. The naive approach fits a linear regression on a log scale, starting from the maximum distance until the end of the time series. The window-based approach tries to select an appropriate window by smoothing the distance estimates and finding a time period when the smoothed time series crosses pre-determined threshold, relatively to the maximum distance; then, a linear regression is fitted by using the raw distance within this time period.



**Figure S8: Estimated time series of distance from attractor for each pathogen across Canada, Hong Kong, Korea, and the US.** Black lines represent the estimated distance from attractor. Red lines and shaded regions represent the linear regression fits and corresponding 95% confidence intervals. Dashed lines represent the average of the pre-pandemic nearest neighbor distances.






Figure S9: **Predicted timing of when each pathogen will return to their pre-pandemic cycles (zoomed out).** The dashed line in panel B indicates the end of 2024 (current observation time). Error bars represent 95% confidence intervals.



**Figure S10: Estimated time series of distance from attractor for each pathogen across Canada, Hong Kong, Korea, and the US using higher embedding dimensions.** Black lines represent the estimated distance from attractor. Red lines and shaded regions represent the linear regression fits and corresponding 95% confidence intervals. Dashed lines represent the average of the pre-pandemic nearest neighbor distances. A lower threshold is used for determining embedding dimensions with the false nearest neighbors approach, thereby yielding a higher embedding dimension.



**Figure S11: Summary of resilience estimates using higher embedding dimensions.** (A) Estimated pathogen resilience. The gray horizontal line represents the intrinsic resilience of pre-vaccination measles dynamics. (B) Predicted timing of when each pathogen will return to their pre-pandemic cycles. The dashed line in panel B indicates the end of 2024 (current observation time). Error bars represent 95% confidence intervals. A lower threshold is used for determining embedding dimensions with the false nearest neighbors approach, thereby yielding a higher embedding dimension.




Figure S12: **Comparison of resilience estimates.** Each point represents a resilience estimate for a specific pathogen at a specific country. The x values represent the original resilience estimates presented in Figure 4. The y values represent the original resilience estimates presented in Supplementary Figure S11.

## 390 References

- 391 [1] Edward A Bender, Ted J Case, and Michael E Gilpin. Perturbation experiments  
392 in community ecology: theory and practice. *Ecology*, 65(1):1–13, 1984.
- 393 [2] Anthony R Ives and Stephen R Carpenter. Stability and diversity of ecosystems.  
394 *science*, 317(5834):58–62, 2007.
- 395 [3] Marten Scheffer, Jordi Bascompte, William A Brock, Victor Brovkin, Stephen R  
396 Carpenter, Vasilis Dakos, Hermann Held, Egbert H Van Nes, Max Rietkerk,  
397 and George Sugihara. Early-warning signals for critical transitions. *Nature*,  
398 461(7260):53–59, 2009.
- 399 [4] Stuart L Pimm. The structure of food webs. *Theoretical population biology*,  
400 16(2):144–158, 1979.
- 401 [5] Michael G Neubert and Hal Caswell. Alternatives to resilience for measuring  
402 the responses of ecological systems to perturbations. *Ecology*, 78(3):653–665,  
403 1997.
- 404 [6] Lance H Gunderson. Ecological resilience—in theory and application. *Annual  
405 review of ecology and systematics*, 31(1):425–439, 2000.
- 406 [7] Vasilis Dakos and Sonia Kéfi. Ecological resilience: what to measure and how.  
407 *Environmental Research Letters*, 17(4):043003, 2022.
- 408 [8] Jeanne C Chambers, Craig R Allen, and Samuel A Cushman. Operationalizing  
409 ecological resilience concepts for managing species and ecosystems at risk.  
410 *Frontiers in Ecology and Evolution*, 7:241, 2019.
- 411 [9] Rachel E Baker, Sang Woo Park, Wenchang Yang, Gabriel A Vecchi, C Jessica E  
412 Metcalf, and Bryan T Grenfell. The impact of COVID-19 nonpharmaceutical  
413 interventions on the future dynamics of endemic infections. *Proceedings of the  
414 National Academy of Sciences*, 117(48):30547–30553, 2020.
- 415 [10] Gabriela B Gomez, Cedric Mahé, and Sandra S Chaves. Uncertain effects of the  
416 pandemic on respiratory viruses. *Science*, 372(6546):1043–1044, 2021.
- 417 [11] Mihaly Koltai, Fabienne Krauer, David Hodgson, Edwin van Leeuwen, Marina  
418 Treskova-Schwarzbach, Mark Jit, and Stefan Flasche. Determinants of RSV  
419 epidemiology following suppression through pandemic contact restrictions. *Epi-  
420 demics*, 40:100614, 2022.
- 421 [12] Sang Woo Park, Brooklyn Noble, Emily Howerton, Bjarke F Nielsen, Sarah  
422 Lentz, Lilliam Ambroggio, Samuel Dominguez, Kevin Messacar, and Bryan T  
423 Grenfell. Predicting the impact of non-pharmaceutical interventions against  
424 COVID-19 on *Mycoplasma pneumoniae* in the United States. *Epidemics*,  
425 49:100808, 2024.

- 426 [13] Eric J Chow, Timothy M Uyeki, and Helen Y Chu. The effects of the COVID-19  
427 pandemic on community respiratory virus activity. *Nature Reviews Microbiology*, 21(3):195–210, 2023.
- 429 [14] Georgiy V Bobashev, Stephen P Ellner, Douglas W Nychka, and Bryan T  
430 Grenfell. Reconstructing susceptible and recruitment dynamics from measles  
431 epidemic data. *Mathematical Population Studies*, 8(1):1–29, 2000.
- 432 [15] Bärbel F Finkenstädt and Bryan T Grenfell. Time series modelling of childhood  
433 diseases: a dynamical systems approach. *Journal of the Royal Statistical Society  
434 Series C: Applied Statistics*, 49(2):187–205, 2000.
- 435 [16] Floris Takens. Detecting strange attractors in turbulence. In *Dynamical Sys-  
436 tems and Turbulence, Warwick 1980: proceedings of a symposium held at the  
437 University of Warwick 1979/80*, pages 366–381. Springer, 2006.
- 438 [17] Alan Hastings, Karen C Abbott, Kim Cuddington, Tessa Francis, Gabriel Gell-  
439 ner, Ying-Cheng Lai, Andrew Morozov, Sergei Petrovskii, Katherine Scran-  
440 ton, and Mary Lou Zeeman. Transient phenomena in ecology. *Science*,  
441 361(6406):eaat6412, 2018.
- 442 [18] Matthew B Kennel, Reggie Brown, and Henry DI Abarbanel. Determining  
443 embedding dimension for phase-space reconstruction using a geometrical con-  
444 struction. *Physical review A*, 45(6):3403, 1992.
- 445 [19] Eugene Tan, Shannon Algar, Débora Corrêa, Michael Small, Thomas Stemler,  
446 and David Walker. Selecting embedding delays: An overview of embedding  
447 techniques and a new method using persistent homology. *Chaos: An Interdis-  
448 ciplinary Journal of Nonlinear Science*, 33(3), 2023.
- 449 [20] Samit Bhattacharyya, Per H Gesteland, Kent Korgenski, Ottar N Bjørnstad,  
450 and Frederick R Adler. Cross-immunity between strains explains the dynamical  
451 pattern of paramyxoviruses. *Proceedings of the National Academy of Sciences*,  
452 112(43):13396–13400, 2015.
- 453 [21] Bryan T Grenfell, Ottar N Bjørnstad, and Bärbel F Finkenstädt. Dynamics of  
454 measles epidemics: scaling noise, determinism, and predictability with the TSIR  
455 model. *Ecological monographs*, 72(2):185–202, 2002.
- 456 [22] Virginia E Pitzer, Cécile Viboud, Vladimir J Alonso, Tanya Wilcox, C Jessica  
457 Metcalf, Claudia A Steiner, Amber K Haynes, and Bryan T Grenfell. Environ-  
458 mental drivers of the spatiotemporal dynamics of respiratory syncytial virus in  
459 the United States. *PLoS pathogens*, 11(1):e1004591, 2015.
- 460 [23] Katharine R Dean, Fabienne Krauer, Lars Walløe, Ole Christian Lingjærde, Bar-  
461 bara Bramanti, Nils Chr Stenseth, and Boris V Schmid. Human ectoparasites

- 462 and the spread of plague in Europe during the Second Pandemic. *Proceedings*  
463 *of the National Academy of Sciences*, 115(6):1304–1309, 2018.
- 464 [24] Margarita Pons-Salort and Nicholas C Grassly. Serotype-specific immunity  
465 explains the incidence of diseases caused by human enteroviruses. *Science*,  
466 361(6404):800–803, 2018.
- 467 [25] Sarah Cobey and Edward B Baskerville. Limits to causal inference with state-  
468 space reconstruction for infectious disease. *PloS one*, 11(12):e0169050, 2016.