

Susceptible host dynamics explain pathogen resilience to perturbations

Sang Woo Park^{1,2,*} Bjarke Frost Nielsen^{2,3} Emily Howerton^{2,3} Bryan T. Grenfell^{2,3,4,5} Sarah Cobey¹

¹ Department of Ecology and Evolution, University of Chicago, Chicago, IL, USA

² [School of Biological Sciences, Seoul National University, Seoul, Republic of Korea](#)

³ Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, USA

^{3,4} High Meadows Environmental Institute, Princeton University, Princeton, NJ, USA

^{4,5} Princeton School of Public and International Affairs, Princeton, NJ, USA

*Corresponding author: swp2sangwoopark@uchicago.edu snu.ac.kr

Abstract

Interventions to slow the spread of SARS-CoV-2 significantly disrupted the transmission of other pathogens. As interventions lifted, whether and when human pathogens would eventually return to their pre-pandemic dynamics remains to be answered. Here, we present a framework for estimating pathogen resilience based on how fast epidemic patterns return to their pre-pandemic dynamics. By analyzing time series data from Hong Kong, Canada, Korea, and the US, we quantify the resilience of common respiratory pathogens and further predict when each pathogen will eventually return to its pre-pandemic dynamics. Our predictions are able to distinguish which pathogens should have returned already, and deviations from these predictions reveal long-term impacts of pandemic perturbations. We find a faster rate of susceptible replenishment underlies pathogen resilience and sensitivity to both large and small perturbations. Overall, our analysis highlights the persistent nature of common respiratory pathogens compared to vaccine-preventable infections, such as measles.

Significance Statement

COVID-19 interventions slowed the transmission of ~~other respiratory pathogens~~ [many respiratory pathogens in different ways](#), raising questions about ~~the mechanisms driving differences~~ [mechanisms driving the variation](#) in responses to COVID-19 intervention measures. To address this gap, we characterized ~~pathogen resilience~~ [the sensitivity of pathogen transmission](#) to perturbations by quantifying how fast each pathogen returned to its pre-pandemic ~~epidemic cycles~~. ~~We applied the resulting~~

37 ~~framework-to-circulation patterns.~~ We analyzed data from Hong Kong, Canada,
38 Korea, and the US, and showed that common respiratory pathogens are ~~much more~~
39 ~~resilient-far less sensitive to perturbations~~ than measles, a vaccine-preventable in-
40 fection. Finally, we showed that the speed of replenishment of the susceptible
41 population—for example, through waning immunity—largely determines ~~a pathogen's~~
42 ~~resilience to perturbations, including large interventions and small stochastic changes~~
43 ~~in the dynamics~~this sensitivity, suggesting that persistence of common respiratory
44 pathogens is likely driven by rapid susceptible replenishment.

Introduction

Non-pharmaceutical interventions to slow the spread of SARS-CoV-2 disrupted the transmission of other human respiratory pathogens, adding uncertainties to their future epidemic dynamics and their public health burden [1]. As interventions lifted, large ~~heterogeneities~~ differences in outbreak dynamics were observed across different pathogens in different countries, with some pathogens exhibiting earlier and faster resurgences than others [2, 3, 4]. ~~Heterogeneities in the overall reduction in transmission and the timing of re-emergence likely reflect differences~~ These differences likely reflect variation in intervention patterns, pathogen characteristics, immigration/importation from other countries, and pre-pandemic pathogen dynamics [5]. Therefore, comparing the differential impact of the pandemic perturbations across pathogens can provide unique opportunities to learn about underlying pathogen characteristics across different populations, such as their transmissibility or duration of immunity ~~, from heterogeneities in re-emergence patterns~~ [6].




Figure 1: Observed heterogeneity in responses to pandemic perturbations across respiratory pathogens and norovirus in (A) Canada, (B) Hong Kong, (C) Korea, and (D) US. Red points and lines represent data before 2020. Blue points and lines represent data since 2020. Gray lines and shaded regions represent the mean seasonal patterns and corresponding 95% confidence intervals around the mean. Mean seasonal patterns were calculated by aggregating cases before 2020 into 52 weekly bins and taking the average in each week. Cases were scaled to account for changes in testing patterns (Materials and Methods).

Even though more than five years have passed since the emergence of SARS-CoV-2, we still observe persistent changes in pathogen dynamics following the pandemic perturbations. For example, compared to pre-pandemic, seasonal patterns, human metapneumovirus in Korea seems to circulate at lower levels, whereas RSV in Korea seems to exhibit different seasonality (Figure 1). These observations suggest the possibility of a long-term change in pathogen dynamics following the pandemic perturbations, which might be driven by a long-term shift in human behavior or population-level immunity [7, 8]. For example, the emergence of SARS-CoV-2 could have caused a long-term shift in population-level immunity through its interactions with other pathogens [9], especially with seasonal coronaviruses [7, 10, 11]. The possibility of a long-lasting impact of the pandemic perturbations poses an important question for future infectious disease dynamics: can we predict whether and when other pathogens will eventually return to their pre-pandemic dynamics?

So far, most analyses of respiratory pathogens after pandemic perturbations have focused on characterizing the timing of rebound ~~[1, 12, 5].~~ (i.e., when the incidence first returns above historical thresholds) and linking this timing to the relaxation of intervention measures or changes in population susceptibility [1, 12, 5]. While quantifying this timing is particularly useful for public health agencies as it allows them to prepare for future outbreaks, epidemic rebound differs from the recovery of the underlying dynamics—the rebound marks the initial return of pathogen activity without providing any information about the subsequent dynamics. Instead, we seek to characterize how fast (and whether) a pathogen returns to its pre-pandemic dynamics by utilizing all of dynamical information following the perturbation, rather than simply focusing on the initial rebound. These two concepts have a subtle but important difference. For example, it took more than 3 years for human metapneumovirus to rebound in Hong Kong, but the observed epidemic patterns in 2024 appear similar to the pre-pandemic seasonal mean, suggesting a ~~possible potential,~~ rapid return to pre-pandemic dynamics ~~; though confirmation may require multiple seasons within 2 years~~ (Figure 1). Measuring this rate of return is useful because it allows us to quantify the ecological resilience of a host-pathogen system, which can inform responses to future interventions [13, 14, 15, 16].

In this study, we lay out theoretical and statistical approaches to characterizing the resilience of a host-pathogen system based on how fast the system recovers from perturbation. We begin by laying out a few representative scenarios that capture the potential impact of pandemic perturbations on endemic pathogen dynamics and illustrate how resilience can be measured by comparing the pre- and post-pandemic dynamics of susceptible and infected hosts. In practice, information on susceptible hosts is often unavailable, making this theoretical approach infeasible. Instead, we utilize a mathematical technique to reconstruct attractors from the data [17], which allows us to measure the rate at which the host-pathogen system approaches this empirical attractor after a perturbation; we define this rate to be the empirical resilience of the host-pathogen system. We use this method to analyze pathogen surveillance data for respiratory and non-respiratory pathogens from Canada, Hong

Kong, Korea, and the US. Finally, we show that susceptible host dynamics explain variation in pathogen resilience and demonstrate that more resilient pathogens will be less sensitive to perturbations caused by demographic stochasticity, thereby providing a direct link between pathogen resilience and persistence.

Conceptual introduction to pathogen resilience

In the classical ecological literature, the resilience of an ecological system is measured by the rate at which the system ~~returns to~~ is expected to return to its reference state following a perturbation [13, 14, 15, 16]. This rate corresponds to the largest real part of the eigenvalues of the linearized system near equilibrium—here, we refer to this value as the *intrinsic* resilience of the system, ~~which represents the expected rate of return from perturbed states~~. In practice, we rarely know the ~~true model describing~~ underlying data generating process that describes the dynamics of common respiratory pathogens, ~~limiting our ability to~~ which can depend on many hidden variables, such as changes in population-level susceptibility. ~~Instead, we have to rely on simplifying model assumptions, raising questions about whether we can~~ infer the intrinsic resilience of a system ~~without measurements of all variables that describe the observed dynamics~~. ~~Nonetheless~~, we can measure the *empirical* resilience of a host-pathogen system by looking at how fast the system returns to the pre-perturbation endemic dynamics after the perturbation has ended. The COVID-19 pandemic provides a crucial example of a major perturbation, providing unique opportunities to measure the resilience of a host-pathogen system across different countries. As we show later, this measure of resilience depends on the rate of susceptible replenishment, where faster replenishment (e.g., through faster waning immunity) causes a pathogen to be more resilient, and further determines how the system responds to small, stochastic perturbations. Therefore, quantifying pathogen resilience provides key information about their dynamical characteristics.

Resilience of a single-strain system under a short-term perturbation.
As an example, we begin with a simple Susceptible-Infected-Recovered-Susceptible (SIRS) model with seasonally forced transmission and demography (i.e., birth and death) ~~—The SIRS model is the simplest model that allows for the waning of immunity and is commonly used for modeling the dynamics of respiratory pathogens [18]. First, consider [18]. Here,~~ a pandemic perturbation that reduces transmission by 50% for 6 months starting in 2020 ~~—which causes epidemic patterns~~ causes the epidemic to deviate from ~~their~~ its original stable annual cycle ~~for a short period of time and eventually come back and eventually return~~ (Figure 2A). To ~~measure the resilience of this system empirically, we first need to be able to quantify resilience,~~ we measure the distance from ~~its~~ the pre-pandemic ~~attractor, which is defined as a~~ attractor—a set of points in state space or phase plane that the system is pulled towards [19]. ~~There are many ways we can measure the distance from the attractor, but for illustrative purposes, we choose one of the most parsimonious approaches:~~

we look at how the susceptible (S) and infected (I) populations change over time and measure the —by taking Euclidean distance on the SI phase plane, —using relative to the counterfactual unperturbed phase plane as a reference trajectory (Figure 2B; Materials and Methods). In this simple case, This distance decreases exponentially with time on average, as demonstrated by the locally estimated scatterplot smoothing (LOESS) fit indicates that the distance from the attractor decreases exponentially (linearly on a log scale) with time on average (Figure 2C). Furthermore, and the overall rate of return approximates matches the intrinsic resilience of the seasonally unforced system (Figure 2C).

Resilience of a single-strain system under a long-term perturbation. Alternatively, pandemic perturbations can have a lasting impact on the forces driving pathogen dynamics through a long-term reduction in transmission or permanent change in immunity. As an example, we consider a scenario in which a For example, if a 10% reduction in transmission persists even after the major pandemic perturbations are lifted, the system will converge to a new attractor (Figure 2D–F). In such cases, we cannot know whether the pathogen will return to its original cycle or a different eyele, but we may not be able to confirm the convergence to a new attractor in practice until many years have passed, and we cannot a priori measure the distance to the new unknown attractor that the system might eventually approach. Nonetheless, we can still measure the distance from the pre-pandemic attractor and ask how the distance changes over time (Figure 2E). The LOESS fit suggests that the distance from the pre-pandemic attractor will initially decrease, which initially decreases exponentially on average (equivalently, linearly on a log scale) and eventually plateau plateaus (Figure 2F). Here, a permanent 10% reduction in transmission rate slows the system, which causes causing a slower decrease in the distance from the pre-pandemic attractor initially to decrease at a slower rate attractor (Figure 2F) than it would have otherwise (Figure 2C) before plateauing at a fixed distance between the two attractors. This example shows that resilience is not necessarily an intrinsic property of a specific pathogen. Instead, pathogen resilience is a property of a specific attractor that a host-pathogen system approaches, which depends on both pathogen and host characteristics.

Resilience of a single-strain system with long-term transients. Finally, transient phenomena can further complicate the picture (Figure 2G–I). For example, a stage-structured model that accounts for reduction in secondary susceptibility initially exhibits a stable annual cycle, but perturbations from a 1050% reduction in transmission for 6 months cause the epidemic to shift to biennial cycles (Figure 2G). The system a transient biennial cycle before it eventually approaches the original pre-pandemic attractor (Figure 2H), suggesting that this biennial cycle is a transient. The LOESS fit indicates that 2G,H). As indicated by the LOESS fit, the distance from the attractor initially decreases exponentially at a rate that is consistent with the intrinsic resilience of the seasonally unforced stage-structured system, but the approach to the attractor rate of return slows down with the damped oscillations (Figure 2I). This behavior is also referred to as a ghost attractor, which causes long




Figure 2: A simple method to measure pathogen resilience following pandemic perturbations across different scenarios. (A, D, G) Simulated epidemic trajectories across various models. Red and blue solid lines represent epidemic dynamics before and after pandemic perturbations are introduced, respectively. Red dashed lines represent counterfactual epidemic dynamics in the absence of perturbations. Gray regions indicate the duration of perturbations with the dark gray region representing a 50% transmission reduction and the light gray region representing a 10% transmission reduction. (B, E, H) Phase plane representation of the time series in panels A, D, and G alongside the corresponding susceptible host dynamics. Red and blue solid lines represent epidemic trajectories on an SI phase plane before and after perturbations are introduced, respectively. (C, F, I) Changes in distance from the attractor over time on a log scale. Blue lines represent the distance from the attractor. Orange lines represent the locally estimated scatterplot smoothing (LOESS) fits to the logged distance from the attractor. Dotted lines show the intrinsic resilience of the seasonally unforced system.

transient dynamics and slow transitions [20]. Strong seasonal forcing in transmission can also lead to transient phenomena for a simple SIRS model, causing a slow return to pre-perturbation dynamics (Supplementary Figure S1). However, we note that a small amount of importation of infections can cause this transient to disappear and cause the system to become much more resilient, even though it has negligible effect on overall dynamics or their empirical resilience under non-seasonal or moderately seasonal transmission (Supplementary Figure S2); under non-seasonal transmission, importation has limited effect on overall resilience of the system unless the amount of importation is increased to a unrealistically high value (Supplementary Figure S3).

Resilience of a two-strain system. This empirical approach allows us to measure the resilience of a two-strain host-pathogen system as well even when we have incomplete observation of the infection dynamics. Simulations from a simple two-strain competition system illustrate that separate analyses of individual strain dynamics (e.g., RSV subtype A vs B) and a joint analysis of total infections (e.g., total RSV infections) yield identical resilience estimates (Supplementary Figure S2, S4, 5). This is expected because eigenvalues determine the dynamics of the entire system around the equilibrium, meaning that both strains should exhibit identical rates of return following a perturbation. Analogous to a single-strain system, strong seasonal forcing in transmission can cause the two-strain system to slow down through transient phenomena (Supplementary Figure S4S6).

These observations yield three insights. First, we can directly estimate the empirical resilience of a host-pathogen system by measuring the rate at which the system approaches an attractor, ~~provided that we have a way to quantify the distance from the attractor—as we discuss later, the attractor of a system can be reconstructed from data from mathematical theory without making assumptions about the underlying model.~~ The empirical approach to estimating pathogen resilience is particularly convenient because it does not require us to know the true ~~underlying model, underlying data generating process~~; estimating the intrinsic resilience from fitting misspecified models can lead to biased estimates (Supplementary Figure S5S7). Second, resilience estimates allow us to make phenomenological predictions about the dynamics of a host-pathogen system following a perturbation. Assuming that an attractor has not changed and the distance from the attractor will decrease exponentially over time, we can estimate when the system should reach an attractor. Finally, a change in the (exponential) rate of approach can provide information about whether the system has reached an alternative attractor, or a ghost attractor, that is different from the original, pre-pandemic attractor. These alternative attractors may reflect permanent changes in transmission patterns as well as changes in immune landscapes. ~~There will be periods of time when it is difficult to tell whether pathogen dynamics are still diverging from the original attractor due to a long-term perturbation, or have entered the basin of attraction of a new attractor.~~ Now that several years have passed since major interventions have been lifted, many respiratory pathogens may have had sufficient time to begin returning to their post-intervention attractors—empirical comparisons between current cases and pre-pandemic cases are consistent

with this hypothesis (Figure 1). With recent data, we can start to evaluate whether we see early signs of convergence to the former attractor or a new one.

Inferring pathogen resilience from real data

Based on these patterns, we now lay out our approach to estimating pathogen resilience from real data (Figure 3). We first tested this approach against simulations and ~~applied it to real data. Specifically, we~~ analyzed case time series of respiratory pathogens from four countries: Canada, Hong Kong, Korea, and the US.

So far, we have focused on simple examples that assume a constant transmission reduction during the pandemic. However, in practice, the impact of pandemic perturbations on pathogen transmission is likely more complex (Figure 3A), reflecting the introduction and relaxation of various intervention strategies. ~~In some cases, strong perturbations likely caused local fadeouts, requiring immigration/importation from another location for epidemic rebound. Such complexities could lead to longer delays between the introduction of pandemic perturbations and pathogen rebound as well as temporal variation in outbreak sizes (Figure 3B); in this~~ For example, continued transmission reduction from interventions ~~limits~~ can limit the size of the first outbreak ~~in 2021~~ following the rebound, allowing for a larger outbreak in ~~2022~~ when interventions are further relaxed (Figure 3B).

Previously, we relied on the dynamics of susceptible and infected hosts to compute the distance from the attractor (Figure 2), but information on susceptible hosts is rarely available in practice. In addition, uncertainties in case counts due to observation error, strain evolution, and multiannual cycles in the observed epidemic dynamics (e.g., adenovirus circulation patterns in Hong Kong and Korea) add challenges to defining pre-pandemic attractors, ~~which limits our ability to measure the distance from the attractor~~. To address these challenges, we can reconstruct an empirical attractor by utilizing Takens’ theorem [17], which states that an attractor of a nonlinear multidimensional system can be mapped onto a delayed embedding (Materials and Methods). ~~For example, we can use delayed~~ Specifically, by taking the logged values of pre-pandemic cases ~~$C(t)$ (Figure 3C) to reconstruct the attractor~~ $\log(C(t) + 1)$ and creating delayed copies, Takens’ theorem allows us to define the pre-pandemic attractor as a points on an M -dimensional space:

$$\langle \log(C(t) + 1), \log(C(t - \tau) + 1), \dots, \log(C(t - (M - 1)\tau) + 1) \rangle, \quad (1)$$

where the delay τ and embedding dimension M are determined based on autocorrelations and false nearest neighbors, respectively [21, 22]. ~~This allows us to define the pre-pandemic attractor as a points on an M -dimensional space. Intuitively, the time series of case counts is generated from a complex system that depends on many hidden variables, such as changes in susceptibility. Takens’ theorem tells us that, delayed copies of the observed series preserves the dynamics of the underlying multi-dimensional attractor.~~ We can then apply the same delay and embedding dimensions to the entire time series to determine the position in multi-dimensional state

space (Figure 3D) ,~~which allows us to and~~ measure the nearest neighbor distance between the current state of the system and the empirical pre-pandemic attractor (Figure 3E). ~~Specifically, the nearest neighbor distance is calculated by computing the distance between the current position on the M -dimensional space and all points in the empirical attractor set and taking the minimum value. In theory, we can now quantify how fast this distance decreases. In theory, the rate at which the distance from this attractor decreases, which can be estimated~~ by fitting a linear regression on a log scale, ~~where the slope of the linear regression empirically measures pathogen resilience, with a steeper slope corresponding to a higher resilience estimate provides an empirical measure of pathogen resilience~~ (Figure 3E). However, resulting estimates of pathogen resilience can be sensitive to choices about embedding delays and dimensions.~~For example, using, where~~ longer delays and higher dimensions tends to smooth out temporal variations in the distance from the attractor (Supplementary Figure S6S8).




Figure 3: A schematic diagram explaining the inference of pathogen resilience from synthetic data. (A) A realistic example of simulated pandemic perturbations, represented by a relative reduction in transmission. (B) The impact of the synthetic pandemic perturbation on epidemic dynamics simulated using a SIRS model with demographic stochasticity. (C) Generating delayed copies of the logged time series allows us to obtain an embedding. (D) Two dimensional representation of an embedding. (E) Delayed embedding allows us to calculate the nearest neighbor distance from the empirical attractor, which is determined based on the pre-pandemic time series. Blue lines and dashed regions represent the linear regression fit and the associated 95% confidence interval. This distance time series can be used to infer pathogen resilience after choosing an appropriate window for linear regression. For illustration purposes, an arbitrary fitting window was chosen.

~~Complex changes in the distance from the attractor suggest. We note~~ that estimating pathogen resilience from linear regression ~~will be particularly sensitive to our choice of fitting windows for the regression is sensitive to choices about regression fitting windows~~ (Figure 3E). Therefore, ~~before we tried estimating resilience from real data, we explored we developed~~ an automated window selection criteria for lin-

ear regression and tested it against randomized, stochastic simulations across a range of realistic pandemic perturbation shapes. ~~In doing so, we~~ We also explored optimal choices for embedding dimensions and ~~evaluated our choices of~~ fitting window parameters ~~and embedding dimensions~~ by quantifying correlation coefficients between the estimated resilience and the intrinsic resilience of a seasonally unforced system (Materials and Methods). ~~Overall, we found large variation in estimation performances with~~ Estimation performances varied widely (correlation coefficients ranging from 0.21 to 0.61) ~~(Supplementary Figure S7). In almost all cases, the automated window selection approach~~ (Supplementary Figure S7), but the automated approach consistently outperformed a naive approach, which ~~performs regression using~~ relies on all data from ~~the the timing of~~ peak distance to ~~current time the present~~ (Supplementary Figure S7S9).

~~Based on~~ We then applied the best-performing window selection criteria and embedding dimension ~~, we applied this approach~~ to pathogen surveillance data presented in Figure 1 (Materials and Methods). For each time series, we applied Takens' theorem independently to reconstruct the empirical attractor and obtained the corresponding time series of distances from attractors (Supplementary Figure ~~S8~~ S10). Then, we ~~used the automated window selection criteria to fit~~ fitted a linear regression and estimated the empirical resilience for each pathogen in each country (Supplementary Figure ~~S8~~ S10). ~~The~~ window selection criteria gave poor regression windows in three cases (norovirus in Hong Kong and Korea and rhinovirus/enterovirus in the US), leading to unrealistically low resilience estimates, and so we used ad-hoc regression windows instead (Supplementary Figure ~~S9~~ S11; Materials and Methods).

For all pathogens we considered, resilience estimates fall between 0.4/year and 1.8/year (Figure 4A). We estimated the mean resilience of common respiratory pathogens to be 0.99/year (95% CI: 0.81/year–1.18/year). For reference, this is ≈ 7.5 times higher than the intrinsic resilience of pre-vaccination measles in England and Wales (≈ 0.13 /year). Finally, resilience estimates for norovirus, a gastrointestinal pathogen, were comparable to those of common respiratory pathogens: 1.44/year (95% CI: 1.01/year–1.87/year) for Hong Kong and 1.07/year (95% CI: 0.86/year–1.29/year) for Korea. Based on a simple ANOVA test, we did not find significant differences in resilience estimates across countries ($p = 0.25$) or pathogens ($p = 0.67$).

Using resilience estimates, we predicted when each pathogen would hypothetically return to their pre-pandemic dynamics, assuming no long-term change in the attractor. Specifically, we extended our linear regression fits to distance-from-attractor time series and ask when the predicted regression line will cross a threshold value; since we relied on nearest neighbor distances, pre-pandemic distances are always greater than zero (Figure 3E), meaning that we can use the mean of pre-pandemic distances as our threshold.

We predicted that a return to pre-pandemic cycles has occurred or would be imminent for most pathogens (Figure 4B). In particular, we predicted that 12 out of 23 pathogen-country pairs should have already returned before the end of 2024. For almost all pathogens that were predicted to have returned already, the observed epidemic dynamics showed clear convergence towards their pre-pandemic seasonal




Figure 4: **Summary of resilience estimates and predictions for return time.** (A) Estimated pathogen resilience. The gray horizontal line represents the intrinsic resilience of pre-vaccination measles dynamics. (B) Predicted timing of when each pathogen will return to their pre-pandemic cycles. The dashed line in panel B indicates the end of 2024. Error bars represent 95% confidence intervals. (C) Observed dynamics for pathogen that are predicted to have returned before the end of 2024. Red points and lines represent data before 2020. Gray lines and shaded regions represent the mean seasonal patterns and corresponding 95% confidence intervals around the mean, previously shown in Figure 1. Orange vertical lines represent the predicted timing of return.

329 averages, confirming our predictions (Figure 4C). However, there were a few ex-
 330 ceptions, including norovirus in Hong Kong and rhinovirus/enterovirus in the US,
 331 where the observed epidemic dynamics in 2024 exhibit clear deviation from their pre-
 332 pandemic seasonal averages (Figure 4C; Figure S9S11). These observations suggest a
 333 possibility that some common respiratory pathogens may have converged to different
 334 attractors or are still exhibiting non-equilibrium dynamics. In contrast, pathogens

that were predicted to have not returned yet also showed clear differences from their pre-pandemic seasonal averages; as many of these pathogens are predicted to return in 2025–2026, we may be able to test these predictions in near future (Supplementary Figure S10S12). Our reconstructions of distance time series and estimates of pathogen resilience and expected return time were generally robust to choices of embedding dimensions (Supplementary Figure S11–12S13–14).

~~To evaluate~~ We also evaluated the ability of our approach to make out-of-sample predictions ~~we applied our framework to the same data set by~~ using observations up to the 26th week of 2023 (Supplementary Figure S13–14). ~~Limiting the data~~ S15–16. This led to greater uncertainty in resilience estimates and corresponding predictions of the expected return time (Supplementary Figure S14AS16A,B). The expected and observed return times exhibit weak positive correlation ($\rho = 0.36$; 95% CI: -0.34–0.81), and 7 out of 10 confidence intervals contain the true value (Supplementary Figure S14CS16C). We found moderate correlations ($\rho = 0.51$; 95% CI: 0.01–0.80) in predictions for the expected return time between partial regression fits, which rely on observations up to the 26th week of 2023, and full regression fits, which rely on the entire data set (Supplementary Figure S14DS16D); this implies consistency in our predictions. However, resilience estimates from partial regression fits and full regression fits were weakly correlated ($\rho = 0.30$; 95% CI: -0.23–0.70), demonstrating challenges in estimating pathogen resilience from limited data (Supplementary Figure S14ES16E).

Susceptible host dynamics explain variation in pathogen resilience

So far, we have focused on quantifying pathogen resilience from the observed patterns of pathogen re-emergence following pandemic perturbations. But what factors determine the resilience of a host-pathogen system? ~~To address this question, we used the SIRS model to explore how changes in~~ Here, we illustrate how susceptible host dynamics affect pathogen resilience ~~. To do so, we varied by varying the basic reproduction number $\mathcal{R}_0$, which represents the average number of secondary infections caused by a newly infected individual in a fully susceptible population, and the duration of immunity and computed intrinsic resilience for each parameter of the SIRS model.~~

~~We found that an~~ An increase in $\mathcal{R}_0$ and a decrease in the duration of immunity ~~correspond to an increase in pathogen resilience leads to a higher resilience value~~ (Figure 5A). ~~Similarly, an increase in $\mathcal{R}_0$ and a decrease in the duration of immunity corresponds to a~~ and a faster per-capita rate of susceptible replenishment, ~~which is defined as the ratio between absolute total~~ rate at which new susceptibles enter the population and the equilibrium number of ~~susceptible individuals in the population~~ susceptibles, S^* (Figure 5B). ~~We note that~~ Specifically, a higher $\mathcal{R}_0$ drives a faster per-capita susceptible replenishment rate by decreasing ~~the susceptible~~




Figure 5: Linking pathogen resilience to epidemiological parameters and susceptible host dynamics. (A) Intrinsic resilience as a function of the basic reproduction number $\mathcal{R}_0$ and the duration of immunity. (B) Per-capita susceptible replenishment rate as a function of the basic reproduction number $\mathcal{R}_0$ and the duration of immunity. The standard SIRS model without seasonal forcing is used to compute intrinsic resilience and the per-capita susceptible replenishment rate. Lines correspond to a set of parameters that are consistent with mean resilience estimates for each pathogen, averaged across different countries.

375 ~~proportion at equilibrium, S^* .~~ For a simple SIR model that assumes a life-long
 376 immunity, ~~we can show analytically that~~ pathogen resilience is proportional to the
 377 per-capita rate of susceptible replenishment (Materials and Methods). Overall, these
 378 observations suggest that a faster per-capita susceptible replenishment rate causes
 379 the system to be more resilient.

380 By taking the average values of empirical resilience for each pathogen, we were
 381 able to map each pathogen onto a set of parameters of the SIRS model that are
 382 consistent with corresponding resilience estimates (Figure 5A). Across all pathogens
 383 we considered, we estimated that the average duration of immunity is likely to be
 384 short (< 4 years) across a plausible range of $\mathcal{R}_0$ (< 6). ~~We were also able to obtain~~
 385 ~~a plausible range of susceptible replenishment rates for each pathogen (Figure 5B);~~
 386 but in contrast, there was a large uncertainty in the estimates for susceptible re-
 387 plenishment rates due to a lack of one-to-one correspondence between susceptible
 388 replenishment rates and pathogen resilience (Figure 5B).

Pathogen resilience determines sensitivity to stochastic perturbations

Even in the absence of major pandemic perturbations, host-pathogen systems are expected to experience continued perturbations of varying degrees from changes in epidemiological conditions, such as human behavior, climate, and viral evolution. These perturbations can also arise from demographic stochasticity, which is inherent to any ecological system. Here, ~~we used~~ a seasonally unforced SIRS model with constant birth and death rates ~~to explore how~~ illustrates that resilience of a host-pathogen system determines the sensitivity to perturbations caused by demographic stochasticity ~~(Materials and Methods).~~

Linking resilience of a host-pathogen system to its sensitivity to stochastic perturbations. ~~(A–C) Epidemic trajectories of a stochastic SIRS model across three different resilience values: low, intermediate, and high. The relative prevalence was calculated by dividing infection prevalence by its mean value. (D) Intrinsic resilience of a system as a function of the basic reproduction number $\mathcal{R}_0$ and the duration of immunity. (E) Epidemic amplitude as a function of the basic reproduction number $\mathcal{R}_0$ and the duration of immunity. The epidemic amplitude corresponds to $(\max I - \min I)/(2\bar{I})$, where $\bar{I}$ represents the mean prevalence. Labels A–C in panels D and E correspond to scenarios shown in panels A–C. (F) The relationship between pathogen resilience and epidemic amplitude.~~

~~We found that resilience of a host-pathogen system determines the amount of, where a less resilient system experiences a greater deviation from the deterministic trajectory caused by demographic stochasticity, with less resilient systems experiencing larger deviations (Figure 6). Notably, less~~ Materials and Methods). Less resilient systems also exhibited slower epidemic cycles (Figure 6A–C). The periodicity of this epidemic cycle matched those predicted by the intrinsic periodicity of the system (Supplementary Figure S15) ~~where the intrinsic resilience of the system~~ S17), which is inversely proportional to ~~its intrinsic periodicity~~ the intrinsic resilience (Supplementary Figure S16). ~~However, we~~ S18). We note that the interplay between seasonal transmission and intrinsic periodicity can also lead to complex cycles, as illustrated by a recent analysis of *Mycoplasma pneumoniae* dynamics [23].

We also note that the intrinsic resilience is not the sole determinant for how sensitive the system is to stochastic perturbations. For example, the population size and average duration of infection also affect the amount of deviation from the deterministic trajectory caused by demographic stochasticity, even though these quantities have little to no impact on the intrinsic resilience (Supplementary Figure S17). These conclusions were robust for the seasonally forced SIRS model, where the amount of deviation from the deterministic trajectory depends on the corresponding resilience of the seasonally unforced system (Supplementary Figure ~~S18~~ S19).




Figure 6: Linking resilience of a host-pathogen system to its sensitivity to stochastic perturbations. (A–C) Epidemic trajectories of a stochastic SIRS model across three different resilience values: low, intermediate, and high. The relative prevalence was calculated by dividing infection prevalence by its mean value. (D) Intrinsic resilience of a system as a function of the basic reproduction number R_0 and the duration of immunity. (E) Epidemic amplitude as a function of the basic reproduction number R_0 and the duration of immunity. The epidemic amplitude corresponds to $(\max I - \min I)/(2\bar{I})$, where $\bar{I}$ represents the mean prevalence. Labels A–C in panels D and E correspond to scenarios shown in panels A–C. (F) The relationship between pathogen resilience and epidemic amplitude.

Discussion

COVID-19 pandemic interventions caused major disruptions to circulation patterns of both respiratory and non-respiratory pathogens, adding challenges to predicting their future dynamics [1, 2, 3, 4]. However, these perturbations offer large-scale natural experiments for understanding how different pathogens respond to perturbations. In this study, we showed that pathogen re-emergence patterns following pandemic perturbations can be characterized through the lens of ecological resilience and presented a new method for estimating pathogen resilience from time series data. We showed that variation in pathogen resilience can be explained by the differences in susceptible host dynamics, where faster replenishment of the susceptible pool corresponds to a more resilient host-pathogen system. Finally, we showed that pathogen resilience also determines the sensitivity to stochastic perturbations.

We analyzed case time series of common respiratory infections and norovirus infections from Canada, Hong Kong, Korea, and the US to estimate their resilience.

Overall, we estimated the resilience of these pathogens to range from 0.4/year to 1.8/year, which is 3–14 times more resilient than prevaccination measles. Consistent with other epidemiological evidence [24, 25, 26, 27], these resilience estimates indicate that common respiratory pathogens and norovirus likely exhibit faster susceptible replenishment and are therefore more persistent, indicating potential challenges in controlling these pathogens.

Based on our resilience estimates, we made phenomenological predictions about when each pathogen will return to their endemic cycles. Our main regression analysis could accurately distinguish which pathogens should have already returned before the end of 2024. However, there were two main exceptions (i.e., norovirus in Hong Kong and rhinovirus/enterovirus in the US), ~~suggesting that these pathogens may be converging to new endemic cycles or experiencing long-term transient behavior~~ hinting at long-lasting impact of pandemic perturbations. These changes may reflect changes in surveillance or actual shift in the dynamics, ~~caused by permanent changes in behavior or population-level immunity~~. While it may seem unlikely that ~~permanent changes in behavior~~ behavioral changes would only affect a few pathogens and not others, we cannot rule out this possibility given differences in the observed mean age of infections and therefore the differences in age groups that primarily drive transmission [28, 29]. Differences in the mode of transmission between respiratory vs gastrointestinal pathogens may also contribute to the differences in responses to pandemic perturbations.

For almost half of the pathogens we considered, we predicted that their return to original epidemic patterns is imminent. However, our evaluation of out-of-sample predictions suggested major challenges in predicting the exact timing of pathogen return and estimating pathogen resilience from limited data. Therefore, our estimates of prediction timing should be taken as a ballpark estimate. We will need a few more years of data to test whether these pathogens will eventually return to their original dynamics or eventually converge to a different attractor. We also cannot rule out the possibility that some pathogens may exhibit long-term transient behaviors following pandemic perturbations. Overall, these observations echo earlier studies that highlighted the long-lasting impact of pandemic perturbations [8, 30, 31, 4, 23].

We showed that susceptible host dynamics shape pathogen resilience, where faster replenishment of the susceptible population causes the pathogen to be more resilient. For simplicity, we focused on waning immunity and birth as the main drivers of the susceptible host dynamics but other mechanisms can also contribute to the replenishment of the susceptible population. In particular, pathogen evolution, especially the emergence of antigenically novel strains, can cause effective waning of immunity in the population; therefore, we hypothesize that the rate of antigenic evolution is likely a key feature of pathogen resilience. Future studies should explore the relationship between the rate of evolution and resilience for antigenically evolving pathogens. This result also highlights the importance of detailed measurements of changes in the susceptible population through immune assays for understanding pathogen dynamics [32].

~~Quantifying pathogen resilience~~ There are many other factors that may affect the resilience of a host-pathogen system beyond susceptible replenishment rate. For example, we showed that importation can also increase the resilience of the system (Supplementary Figures S2, 3). Interestingly, this effect is minimal without seasonal forcing but substantially amplified under strong seasonal forcing, suggesting that analyzing the eigenvalues of seasonally unforced system may be insufficient to capture key aspects of the system’s resilience. Future studies should consider both analytical and numerical analyses of more detailed, mechanistic models to understand how other factors—such as the generation-interval distribution [33], immune boosting [34], strain competition [35], and post-infection mortality [36]—may alter the resilience of a host-pathogen system. Quantifying relative changes in pathogen resilience with respect to underlying parameters of a model—analogue to sensitivity and elasticity analyses in ecology [37, 38]—may allow one to infer relative importance of different factors that affect pathogen resilience.

Quantifying pathogen resilience also offers novel approaches to validating population-level epidemiological models. So far, most model validation in infectious disease ecology is based on the ability of a model to reproduce the observed epidemic dynamics and to predict future dynamics [39, 40, 26, 41, 42]. However, many models can perform similarly under these criteria. For example, two major RSV models have been proposed to explain biennial epidemic patterns: (1) a stage- and age-structured model that allows disease severity to vary with number of past infections and age of infection [26] and (2) a pathogen-interaction model that accounts for cross immunity between RSV and human metapneumovirus [40]. Since both models can accurately reproduce the observed epidemic patterns, standard criteria for model validation do not allow us to distinguish between these two models from population-level data alone. Instead, it would be possible to measure the empirical resilience of each model by simulating various perturbations and comparing the simulations to estimates of empirical resilience from data, using pandemic perturbations as a reference.

There are several limitations to our work. First, we did not extensively explore other approaches to reconstructing the attractor. Recent studies showed that more sophisticated approaches, such as using non-uniform embedding, can provide more robust reconstruction for noisy data [22]. In the context of causal inference, choices about embedding can have major impact on the resulting inference [43]. Our resilience estimates are likely overly confident given a lack of uncertainties in attractor reconstruction as well as the simplicity of our statistical framework. Nonetheless, as illustrated in our sensitivity analyses, inferences about pathogen resilience in our SIRS model appear to be robust to decisions about embedding lags and dimensions—this is because tracking the rate at which the system approaches the attractor is likely a much simpler problem than making inferences about causal directionality. Short pre-pandemic time series also limit our ability to accurately reconstruct the attractor and contribute to the crudeness of our resilience estimates; although this is less likely a problem for respiratory pathogens that are strongly annual, our attractor reconstruction may be inaccurate for those exhibiting multi-annual cycles, such as

adenovirus in Hong Kong and Korea. Our framework also does not allow us to distinguish whether a system has settled to a new attractor or is experiencing long-term transient behavior.

Uncertainties in pathogen dynamics due to changes in testing patterns further contribute to the crudeness of our resilience estimates. We tried to account for changes in testing patterns by either scaling the case counts (for Canada, Hong Kong, and Korea) or deriving incidence proxy (for the US). We were unable to derive incidence proxy for Canada and Hong Kong due to instability of testing positivity, including long-term reduction in test positivity associated with an increase in testing volume. Consistency in resilience estimates across different countries provide some robustness to our conclusions, our exact estimates are likely sensitive to changes in testing. We also cannot rule out the possibility that potential long-term impacts of pandemic perturbations may simply reflect changes in testing, rather than actual dynamical changes. Future studies should investigate improved methods for accounting for changes in testing patterns to obtain a more accurate perspective on the differences between current and past outbreak patterns of endemic pathogens.

While attractor reconstruction allows us to make model-free inferences of pathogen resilience, it does not allow us to tease apart how different mechanisms contribute to the resilience a host-pathogen system. Using the simple SIRS model, we illustrated that susceptible host dynamics are key determinants of pathogen resilience, but we also found that there isn't a one-to-one correspondence between per capita replenishment rate of the susceptible population and pathogen resilience estimates. Future studies should explore using mechanistic models to explain heterogeneity in resilience estimates across different pathogens.

Finally, our simulation-based analyses primarily focused on single-strain systems, but real-world pathogens can interact with other pathogens, which can result in complex dynamics [44, 45]. To address this limitation, we considered a simple model of two competing strains (via cross immunity) and showed that the resilience of a coupled system can be measured by studying the dynamics of either strain. However, this conclusion likely depends on the strength and mechanism of strain interactions. For example, ecological interference between two unrelated pathogens [44] will likely generate weaker coupling than cross-immunity between related pathogens; in the former case, we do not necessarily expect two unrelated pathogens to have same resilience despite their ecological interference. Some pathogen strains can also exhibit positive interactions where infection by one strain can lead to an increased transmission of another competing strain. For example, previous studies showed that an increased dengue transmission through antibody-dependent enhancement can permit coexistence and persistence of competing strains [46]; since pathogen transmissibility is a major determinant of pathogen resilience, we tentatively hypothesize that positive interactions such as antibody-dependent enhancement may increase the resilience of a system. Future studies should explore how different mechanisms of pathogen interactions contribute to the resilience of each competing pathogen as well as the entire system. Despite these limitations, our study illustrates that quantifying pathogen

resilience can provide novel insights into pathogen dynamics. Furthermore, our qualitative prediction that common respiratory pathogens are more resilient than pre-vaccination measles is also likely to be robust, given how rapidly many respiratory pathogens returned to their original cycles following pandemic perturbations.

Predicting the impact of anthropogenic changes on infectious disease dynamics is a fundamental aim of infectious disease research in a rapidly changing world. Our study illustrates that how a host-pathogen system responds to both small and large perturbations is largely predictable through the lens of ecological resilience. In particular, quantifying the resilience of a host-pathogen system offers a unique insight into questions about endemic pathogens' responses to pandemic perturbations, including whether some pathogens will exhibit long-lasting impact from the pandemic perturbation or not. More broadly, a detailed understanding of the determinants of pathogen resilience can provide deeper understanding of pathogen persistence.

Materials and Methods

Data

We gathered time series on respiratory infections from Canada, Hong Kong, Korea, and United States (US). As a reference, we also included ~~time series data on~~ norovirus infections when available. In contrast to respiratory pathogens, we hypothesized gastrointestinal viruses, such as norovirus, to be differently affected by pandemic perturbations.

Weekly time series of respiratory infection cases in Canada came from a publicly available website by the Respiratory Virus Detection Surveillance System, which collects data from select laboratories across Canada [47]. Weekly time series of respiratory infection cases in Hong Kong came from a publicly available website by the Centre for Health Protection, Department of Health [48, 49]. Weekly time series of acute respiratory infection cases in Korea came from a publicly available website by the Korea Disease Control and Prevention Agency [50]. Finally, weekly time series of respiratory infection cases in the US were obtained from the National Respiratory and Enteric Virus Surveillance System (NREVSS). Readers can request the data from NREVSS at nrevss@cdc.gov. Time series on number of tests were also available in Canada, Hong Kong, and the US, but not in Korea.

Data processing

For all time series, we rounded every year to 52 weeks by taking the average number of cases and tests between the 52nd and 53rd week. We also rescaled all time series to account for changes in testing patterns, which were then used for the actual analysis.

For Canada, an increase in testing was observed from 2013 to 2024 (Supplementary Figure ~~S17~~S21); this increase is associated with a decrease in test positivity during the same period (Supplementary Figure S22). To account for this increase,

we calculated a 2 year moving average for the number of tests for each pathogen, which we used as a proxy for testing effort. Then, we divided the smoothed testing patterns by the smoothed value at the final week such that the testing effort has a maximum of 1. We then divided weekly cases by the testing effort to obtain a scaled case time series. A similar approach was used earlier for an analysis of RSV time series in the US to account for changes in testing patterns [26].

For Hong Kong, we applied the same scaling procedure to the time series as we did for Canada. In this case, we only adjusted for testing efforts up to the end of 2019 because there was a major reduction in testing for common respiratory pathogens between 2020 and 2023 (Supplementary Figure S18-S23). [Analogous to patterns in Canada, an increase in testing volume is associated with a decrease in test positivity \(Supplementary Figure S24\).](#)

For Korea, while we did not have information on testing, the reported number of respiratory infections consistently increased from 2013 to the end of 2019, which we interpreted as changes in testing patterns (Supplementary Figure S19-S25). Since we did not have testing numbers, we used the weekly sum of all acute respiratory viral infection cases as a proxy for testing, which were further smoothed with moving average and scaled to have a maximum of 1. For Korea, we also only adjusted for testing efforts up to the end of 2019.

In the US, there has been a large increase in testing for some respiratory pathogens, especially RSV, which could not be corrected by simple scaling (Supplementary Figure S20-S26). Instead, we derived an incidence proxy by multiplying the test positivity with influenza-like illness positivity, which was taken from <https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>. This method of estimating an incidence proxy has been recently applied in the analysis of seasonal coronaviruses [7] and *Mycoplasma pneumoniae* infections [4]. Detailed assumptions and justifications are provided in [51].

Data summary

To make qualitative comparisons between pre- and post-perturbation dynamics of respiratory pathogen circulation patterns, we calculated the mean seasonal patterns using time series of either rescaled cases or incidence proxy estimates before 2020. We did so by taking the mean value in each week across all years before 2020. Confidence intervals around the means were calculated using a simple t test.

Estimating pathogen resilience

In order to measure pathogen resilience from surveillance data, we first reconstructed the empirical pre-pandemic attractor of the system using Takens' embedding theorem [17]. Specifically, for a given pathogen, we took the pre-pandemic (before 2020) case time series $C(t)$ and reconstructed the attractor using delayed embedding with a

uniform delay of τ and dimension M :

$$X_{\tau,m}(t) = \langle \log(C(t) + 1), \log(C(t - \tau) + 1), \dots, \log(C(t - (M - 1)\tau) + 1) \rangle. \quad (2)$$

Here, the delay τ was determined by calculating the autocorrelation of the logged pre-pandemic time series and asking when the autocorrelation crosses 0 for the first time [22]; a typical delay for an annual outbreak is around 13 weeks.

Then, for a given delay τ , we determined the embedding dimension M using the false nearest neighbors approach [21, 22]. To do so, we started with an embedding dimension e and constructed a set of points $A_{\tau,e} = \{X_{\tau,e}(t) | t < 2020\}$. Then, for each point $X_{\tau,e}(t)$, we determined the nearest neighbor from the set $A_{\tau,e}$, which we denote $X_{\tau,e}(t_{nn})$ for $t \neq t_{nn}$. Then, if the distance between these two points in the $e+1$ dimension, $D_{\tau,e+1}(t) = \|X_{\tau,e+1}(t_{nn}) - X_{\tau,e+1}(t)\|_2$, is larger than their distance in the e dimension, $D_{\tau,e}(t) = \|X_{\tau,e}(t_{nn}) - X_{\tau,e}(t)\|_2$, these two points are deemed to be false nearest neighbors; specifically, we used a threshold R for the ratio between two distances $D_{\tau,e+1}(t)/D_{\tau,e}(t)$ to determine false nearest neighbors. The first embedding dimension e that does not have any false nearest neighbors corresponds to the embedding dimension M for a given pathogen-country pair. For the main analysis, we used $R = 10$, which was chosen from a sensitivity analysis against simulated data (Supplementary Text). Once we determined the embedding lag τ and dimension M , we apply the embedding to the entire time series and calculate the nearest neighbor distance against the attractor $A_{\tau,M}$ to obtain a time series of distance from the attractor $D_{\tau,M}(t)$.

From a time series of distances from the attractor, we estimated pathogen resilience by fitting a linear regression to an appropriate window. To automatically select fitting windows, we began by smoothing the distance time series using locally estimated scatterplot smoothing (LOESS) to obtain $\hat{D}_{\tau,M}(t)$, where the smoothing is performed on a log scale and exponentiated afterwards. This smoothing allowed us to find appropriate threshold values for selecting fitting windows that are insensitive to errors in our estimates of distance from the attractor. Then, we determined threshold values (T_{start} and T_{end}) for the smoothed distances and choose the fitting window based on when $\hat{D}_{\tau,M}(t)$ crosses these threshold values for the first time. These thresholds were determined by first calculating the maximum distance,

$$\max \hat{D} = \max \hat{D}_{\tau,M}(t), \quad (3)$$

and the mean pre-pandemic distance,

$$\hat{D}_{\text{mean}} = \frac{1}{N_{t < 2020}} \sum_{t < 2020} \hat{D}_{\tau,M}(t), \quad (4)$$

as a reference, and then dividing their ratios into K equal bins,

$$T_{\text{start}} = \hat{D}_{\text{mean}} \times \left(\frac{\max \hat{D}}{\hat{D}_{\text{mean}}} \right)^{(K-a)/K}, \quad (5)$$

$$T_{\text{end}} = \hat{D}_{\text{mean}} \times \left(\frac{\max \hat{D}}{\hat{D}_{\text{mean}}} \right)^{a/K}, \quad (6)$$

where a represents the truncation threshold. This allows us to discard the initial period during which the distance increases (from the introduction of intervention measures) and the final period during which the distance plateaus (as the system reaches an attractor). The fitting window is determined based on when the smoothed distance $\hat{D}_{\tau,M}(t)$ crosses these threshold values for the first time; then, we fit a linear regression to logged (unsmoothed) distances $\log D_{\tau,M}(t)$ using that window. Alongside the threshold R for the false nearest neighbors approach, we tested optimal choices for K and a values using simulations (Supplementary Text). We used $K = 19$ and $a = 2$ throughout the paper based on the simulation results. We excluded data sets from the regression analysis if the logged average of the estimated distance from the attractor during the last year is at least 2-fold greater than the logged average of the pre-pandemic distance from the attractor.

To evaluate the ability to predict the precise return time for out-of-sample observations, we applied the same framework using data up to the 26th week of 2023. In doing so, we compared the predicted return time with the observed return time, which was determined based on when the smoothed empirical distance from the attractor became close enough to pre-pandemic levels. As a pathogen returns to its pre-pandemic cycle, the the smoothed empirical distance from the attractor is expected to converge to the pre-pandemic levels $\hat{D}_{\text{mean}}$, but the empirical distance may not become smaller than $\hat{D}_{\text{mean}}$. Therefore, we used $1.2\hat{D}_{\text{mean}}$ as our threshold to obtain an approximate return time. We note that this approach necessarily underestimates the observed return time and therefore should be interpreted as a conservative estimate. For additional validation, we compared these out-of-sample predictions with return time predictions based on full time series.

Mathematical modeling

Throughout the paper, we use a series of mathematical models to illustrate the concept of pathogen resilience and to understand the determinants of pathogen resilience. In general, the intrinsic resilience of a given system is given by the largest real part of the eigenvalues of the linearized system at endemic equilibrium. Here, we focus on the SIRS model with demography (birth and death) and present the details of other models in Supplementary Text. The SIRS (Susceptible-Infected-Recovered-Susceptible) model is the simplest model that allows for waning of immunity, where recovered (immune) individuals are assumed to become fully susceptible after an average of $1/\delta$ time period. The dynamics of the SIRS model is described by the following set of differential equations:

$$\frac{dS}{dt} = \mu - \beta(t)SI - \mu S + \delta R \quad (7)$$

$$\frac{dI}{dt} = \beta(t)SI - (\gamma + \mu)I \quad (8)$$

$$\frac{dR}{dt} = \gamma I - (\delta + \mu)R \quad (9)$$

where μ represents the birth and death rates, $\beta(t)$ represents the time-varying transmission rate, and γ represents the recovery rate. The basic reproduction number $\mathcal{R}_0(t) = \beta(t)/(\gamma + \mu)$ is defined as the average number of secondary infections that a single infected individual would cause in a fully susceptible population at time t and measures the intrinsic transmissibility of a pathogen.

When we introduced the idea of pathogen resilience (Figure 2), we imposed sinusoidal changes to the transmission rate to account for seasonal transmission,

$$\beta(t) = b_1(1 + \theta \cos(2\pi(t - \phi)))\alpha(t), \quad (10)$$

where b_1 represents the baseline transmission rate, θ represents the seasonal amplitude, and ϕ represents the seasonal offset term. Here, we also introduced an extra multiplicative term $\alpha(t)$ to account for the impact of pandemic perturbations, where $\alpha(t) < 1$ indicates transmission reduction. Figure 2A and 2B were generated assuming $b_1 = 3 \times (365/7 + 1/50)/\text{years}$, $\theta = 0.2$, $\phi = 0$, $\mu = 1/50/\text{years}$, $\gamma = 365/7/\text{years}$, and $\delta = 1/2/\text{years}$. Specifically, $b_1 = 3 \times (365/7 + 1/50)/\text{years}$ implies $\mathcal{R}_0 = 3$, where $(365/7 + 1/50)/\text{years}$ represent the rate of recovery. In Figure 2A, we imposed a 50% transmission reduction for 6 months from 2020:

$$\alpha(t) = \begin{cases} 0.5 & 2020 \leq t < 2020.5 \\ 1 & \text{otherwise} \end{cases} \quad (11)$$

In Figure 2B, we imposed a 50% transmission reduction for 6 months from 2020 and a permanent 10% reduction onward:

$$\alpha(t) = \begin{cases} 1 & t < 2020 \\ 0.5 & 2020 \leq t < 2020.5 \\ 0.9 & 2020.5 \leq t \end{cases} \quad (12)$$

In both scenarios, we simulated the SIRS model from the same initial conditions ($S(0) = 1/\mathcal{R}_0$, $I(0) = 1 \times 10^{-6}$, and $R(0) = 1 - S(0) - I(0)$) from 1900 until 2030. When we include importation in Supplementary Figures S2 and S3, we model the force of infection as $\beta(t)(I + \omega)$ instead and vary ω between 10^{-7} and 10^{-4} per day. Throughout the paper, all deterministic models were solved using the `lsoda` solver from the `deSolve` package [52] in R [53].

To measure the empirical resilience of the SIRS model (Figure 2C and 2F), we computed the normalized distance between post-intervention susceptible and logged infected proportions and their corresponding unperturbed values at the same time:

$$\sqrt{\left(\frac{S(t) - S_{\text{unperturbed}}(t)}{\sigma_S}\right)^2 + \left(\frac{\log I(t) - \log I_{\text{unperturbed}}(t)}{\sigma_{\log I}}\right)^2}, \quad (13)$$

740 where σ_S and $\sigma_{\log I}$ represent the standard deviation in the unperturbed susceptible
 741 and logged infected proportions. The unperturbed values were obtained by simulat-
 742 ing the same SIRS model without pandemic perturbations ($\alpha = 1$). We normalized
 743 the differences in susceptible and logged infected proportions to allow both quantities
 744 to equally contribute to the changes in distance from the attractor. We used logged
 745 prevalence, instead of absolute prevalence, in order to capture epidemic dynamics
 746 in deep troughs during the intervention period. In Supplementary Materials, we
 747 also compared how the degree of seasonal transmission affects empirical resilience
 748 by varying θ from 0 to 0.4; when we assumed no seasonality ($\theta = 0$), we did not
 749 normalize the distance because the standard deviation of pre-intervention dynamics
 750 are zero.

751 We used the SIRS model to understand how underlying epidemiological param-
 752 eters affect pathogen resilience and determine the relationship to underlying sus-
 753 ceptible host dynamics. For the simple SIRS model without seasonal transmission
 754 ($\theta = 0$), the intrinsic resilience equals

$$-\frac{\text{Re}(\lambda)}{2}\text{Re}(\lambda) = \frac{\delta + \beta I^* + \mu}{2}. \quad (14)$$

755 Here, I^* represents the prevalence at endemic equilibrium:

$$I^* = \frac{(\delta + \mu)(\beta - (\gamma + \mu))}{\beta(\delta + \gamma + \mu)}. \quad (15)$$

756 The susceptible replenishment rate is given by

$$S_{\text{replenish}} = \frac{\mu + \delta R}{S^*}, \quad (16)$$

757 where $S^* = 1/\mathcal{R}_0$ represents the equilibrium proportion of susceptible individuals.
 758 We varied the basic reproduction number $\mathcal{R}_0$ between 1.1 to 6 and the average
 759 duration of immunity $1/\delta$ between 2 to 4 years, and computed these two quantities.
 760 In doing so, we fixed all other parameters: $\mu = 1/80/\text{years}$ and $\gamma = 365/7/\text{years}$.
 761 When infection provides a life-long immunity ($\delta = 0$), the model collapses to the SIR
 762 model and the intrinsic resilience is ~~inversely~~ exactly proportional to the susceptible
 763 replenishment rate:

$$-\frac{\text{Re}(\lambda)}{2}\text{Re}(\lambda) = \frac{\mu}{2S^*} = \frac{S_{\text{replenish}}}{2}. \quad (17)$$

764 Finally, we used a seasonally unforced stochastic SIRS model without demog-
 765 raphy to understand how pathogen resilience affects sensitivity of the system to
 766 demographic stochasticity (see Supplementary Text for the details of the stochas-
 767 tic SIRS model). By varying the basic reproduction number $\mathcal{R}_0$ between 2 to 20
 768 and the average duration of immunity $1/\delta$ between 1 to 40 years, we ran the SIRS
 769 model for 100 years and computed the epidemic amplitude, which we defined as

770 $(\max I - \min I)/(2\bar{I})$. Each simulation began from the equilibrium, and we trun-
771 cated the initial 25 years before computing the epidemic amplitude. In doing so,
772 we assumed $\gamma = 365/7/\text{years}$ and fixed the population size to 1 billion to prevent
773 any fadeouts. We also considered a seasonally forced stochastic SIRS model without
774 demography, assuming an amplitude of seasonal forcing of 0.04; in this case, we com-
775 puted the relative epidemic amplitude by comparing the deterministic and stochastic
776 trajectories (Supplementary Materials).

777 Acknowledgement

778 We thank Anthony R. Ives for helpful discussion. We thank Centers for Disease
779 Control and Prevention, National Respiratory and Enteric Virus Surveillance System
780 (NREVSS) for providing time series data for respiratory infection cases in the US.

781 Data availability

782 All data and code are stored in a publicly available GitHub repository (<https://github.com/cobeylab/return-time>).
783

784 Funding

785 S.W.P. is a Peter and Carmen Lucia Buck Foundation Awardee of the Life Sciences
786 Research Foundation. B.F.N. acknowledges financial support from the Carlsberg
787 Foundation (grant no. CF23-0173). B.T.G is supported by Princeton Catalysis Ini-
788 tiative and Princeton Precision Health. S.C. is supported by Federal funds from the
789 National Institute of Allergy and Infectious Diseases, National Institutes of Health,
790 Department of Health and Human Services under CEIRR contract 75N93021C00015
791 [Subcontract 77789](#). The content is solely the responsibility of the authors and does
792 not necessarily represent the official views of the NIAID or the National Institutes
793 of Health. This project has been funded in whole or in part with Federal funds from
794 the National Cancer Institute, National Institutes of Health, under Prime Contract
795 No. 75N91019D00024, Task Order No. 75N91023F00016 (B.F.N., E.H., and B.T.G).
796 The content of this publication does not necessarily reflect the views or policies of
797 the Department of Health and Human Services, nor does mention of trade names,
798 commercial products or organizations imply endorsement by the U.S. Government.

References

- [1] Rachel E Baker, Sang Woo Park, Wenchang Yang, Gabriel A Vecchi, C Jessica E Metcalf, and Bryan T Grenfell. The impact of COVID-19 nonpharmaceutical interventions on the future dynamics of endemic infections. *Proceedings of the National Academy of Sciences*, 117(48):30547–30553, 2020.
- [2] Gabriela B Gomez, Cedric Mahé, and Sandra S Chaves. Uncertain effects of the pandemic on respiratory viruses. *Science*, 372(6546):1043–1044, 2021.
- [3] Mihaly Koltai, Fabienne Krauer, David Hodgson, Edwin van Leeuwen, Marina Treskova-Schwarzbach, Mark Jit, and Stefan Flasche. Determinants of RSV epidemiology following suppression through pandemic contact restrictions. *Epidemics*, 40:100614, 2022.
- [4] Sang Woo Park, Brooklyn Noble, Emily Howerton, Bjarke F Nielsen, Sarah Lentz, Lilliam Ambroggio, Samuel Dominguez, Kevin Messacar, and Bryan T Grenfell. Predicting the impact of non-pharmaceutical interventions against COVID-19 on *Mycoplasma pneumoniae* in the United States. *Epidemics*, 49:100808, 2024.
- [5] Amanda C Perofsky, Chelsea L Hansen, Roy Burstein, Shanda Boyle, Robin Prentice, Cooper Marshall, David Reinhart, Ben Capodanno, Melissa Truong, Kristen Schwabe-Fry, et al. Impacts of human mobility on the citywide transmission dynamics of 18 respiratory viruses in pre-and post-COVID-19 pandemic years. *Nature communications*, 15(1):4164, 2024.
- [6] Eric J Chow, Timothy M Uyeki, and Helen Y Chu. The effects of the COVID-19 pandemic on community respiratory virus activity. *Nature Reviews Microbiology*, 21(3):195–210, 2023.
- [7] Stephen M Kissler, Christine Tedijanto, Edward Goldstein, Yonatan H Grad, and Marc Lipsitch. Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period. *Science*, 368(6493):860–868, 2020.
- [8] Rachel E Baker, Chadi M Saad-Roy, Sang Woo Park, Jeremy Farrar, C Jessica E Metcalf, and Bryan T Grenfell. Long-term benefits of nonpharmaceutical interventions for endemic infections are shaped by respiratory pathogen dynamics. *Proceedings of the National Academy of Sciences*, 119(49):e2208895119, 2022.
- [9] Maaïke C Swets, Clark D Russell, Ewen M Harrison, Annemarie B Docherty, Nazir Lone, Michelle Girvan, Hayley E Hardwick, Leonardus G Visser, Peter JM Openshaw, Geert H Groeneveld, et al. SARS-CoV-2 co-infection with influenza viruses, respiratory syncytial virus, or adenoviruses. *The Lancet*, 399(10334):1463–1464, 2022.

- [10] Chun-Yang Lin, Joshua Wolf, David C Brice, Yilun Sun, Macauley Locke, Sean Cherry, Ashley H Castellaw, Marie Wehenkel, Jeremy Chase Crawford, Veronika I Zarnitsyna, et al. Pre-existing humoral immunity to human common cold coronaviruses negatively impacts the protective SARS-CoV-2 antibody response. *Cell host & microbe*, 30(1):83–96, 2022.
- [11] Sam M Murray, Azim M Ansari, John Frater, Paul Klenerman, Susanna Dunachie, Eleanor Barnes, and Ane Ogbe. The impact of pre-existing cross-reactive immunity on SARS-CoV-2 infection and vaccine responses. *Nature Reviews Immunology*, 23(5):304–316, 2023.
- [12] John-Sebastian Eden, Chisha Sikazwe, Ruopeng Xie, Yi-Mo Deng, Sheena G Sullivan, Alice Michie, Avram Levy, Elena Cutmore, Christopher C Blyth, Philip N Britton, et al. Off-season RSV epidemics in Australia after easing of COVID-19 restrictions. *Nature communications*, 13(1):2884, 2022.
- [13] Stuart L Pimm. The structure of food webs. *Theoretical population biology*, 16(2):144–158, 1979.
- [14] Michael G Neubert and Hal Caswell. Alternatives to resilience for measuring the responses of ecological systems to perturbations. *Ecology*, 78(3):653–665, 1997.
- [15] Lance H Gunderson. Ecological resilience—in theory and application. *Annual review of ecology and systematics*, 31(1):425–439, 2000.
- [16] Vasilis Dakos and Sonia Kéfi. Ecological resilience: what to measure and how. *Environmental Research Letters*, 17(4):043003, 2022.
- [17] Floris Takens. Detecting strange attractors in turbulence. In *Dynamical Systems and Turbulence, Warwick 1980: proceedings of a symposium held at the University of Warwick 1979/80*, pages 366–381. Springer, 2006.
- [18] Jonathan Dushoff, Joshua B Plotkin, Simon A Levin, and David JD Earn. Dynamical resonance can account for seasonality of influenza epidemics. *Proceedings of the National Academy of Sciences*, 101(48):16915–16916, 2004.
- [19] Alan Hastings, Carole L Hom, Stephen Ellner, Peter Turchin, and H Charles J Godfray. Chaos in ecology: is mother nature a strange attractor? *Annual review of ecology and systematics*, pages 1–33, 1993.
- [20] Alan Hastings, Karen C Abbott, Kim Cuddington, Tessa Francis, Gabriel Gellner, Ying-Cheng Lai, Andrew Morozov, Sergei Petrovskii, Katherine Scranton, and Mary Lou Zeeman. Transient phenomena in ecology. *Science*, 361(6406):eaat6412, 2018.

- [21] Matthew B Kennel, Reggie Brown, and Henry DI Abarbanel. Determining embedding dimension for phase-space reconstruction using a geometrical construction. *Physical review A*, 45(6):3403, 1992.
- [22] Eugene Tan, Shannon Algar, Débora Corrêa, Michael Small, Thomas Stemler, and David Walker. Selecting embedding delays: An overview of embedding techniques and a new method using persistent homology. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 33(3), 2023.
- [23] Bjarke Frost Nielsen, Sang Woo Park, Emily Howerton, Olivia Frost Lorentzen, Mogens H Jensen, and Bryan T Grenfell. Complex multiannual cycles of *Mycoplasma pneumoniae*: persistence and the role of stochasticity. *arXiv*, 2025.
- [24] Samantha Bosis, Susanna Esposito, HGM Niesters, GV Zuccotti, G Marseglia, Marcello Lanari, Giovanna Zuin, C Pelucchi, ADME Osterhaus, and Nicola Principi. Role of respiratory pathogens in infants hospitalized for a first episode of wheezing and their impact on recurrences. *Clinical microbiology and infection*, 14(7):677–684, 2008.
- [25] Miguel L O’Ryan, Yalda Lucero, Valeria Prado, María Elena Santolaya, Marcela Rabello, Yanahara Solis, Daniela Berrios, Miguel A O’Ryan-Soriano, Hector Cortés, and Nora Mamani. Symptomatic and asymptomatic rotavirus and norovirus infections during infancy in a Chilean birth cohort. *The Pediatric infectious disease journal*, 28(10):879–884, 2009.
- [26] Virginia E Pitzer, Cécile Viboud, Wladimir J Alonso, Tanya Wilcox, C Jessica Metcalf, Claudia A Steiner, Amber K Haynes, and Bryan T Grenfell. Environmental drivers of the spatiotemporal dynamics of respiratory syncytial virus in the United States. *PLoS pathogens*, 11(1):e1004591, 2015.
- [27] Arthur WD Edridge, Joanna Kaczorowska, Alexis CR Hoste, Margreet Bakker, Michelle Klein, Katherine Loens, Maarten F Jebbink, Amy Matser, Cormac M Kinsella, Paloma Rueda, et al. Seasonal coronavirus protective immunity is short-lasting. *Nature medicine*, 26(11):1691–1693, 2020.
- [28] Jennifer M Radin, Anthony W Hawksworth, Peter E Kammerer, Melinda Balansay, Rema Raman, Suzanne P Lindsay, and Gary T Brice. Epidemiology of pathogen-specific respiratory infections among three US populations. *PLoS One*, 9(12):e114871, 2014.
- [29] Guojian Lv, Limei Shi, Yi Liu, Xuecheng Sun, and Kai Mu. Epidemiological characteristics of common respiratory pathogens in children. *Scientific Reports*, 14(1):16299, 2024.
- [30] Saverio Caini, Adam Meijer, Marta C Nunes, Laetitia Henaff, Malaika Zounon, Bronke Boudewijns, Marco Del Riccio, and John Paget. Probable extinction of

- 907 influenza b/yamagata and its public health implications: a systematic literature
908 review and assessment of global surveillance databases. *The Lancet Microbe*,
909 2024.
- 910 [31] Zhiyuan Chen, Joseph L-H Tsui, Bernardo Gutierrez, Simon Busch Moreno,
911 Louis du Plessis, Xiaowei Deng, Jun Cai, Sumali Bajaj, Marc A Suchard,
912 Oliver G Pybus, et al. COVID-19 pandemic interventions reshaped the global
913 dispersal of seasonal influenza viruses. *Science*, 386(6722):eadq3003, 2024.
- 914 [32] Hai Nguyen-Tran, Sang Woo Park, Kevin Messacar, Samuel R Dominguez,
915 Matthew R Vogt, Sallie Permar, Perdita Permaul, Michelle Hernandez, Daniel C
916 Douek, Adrian B McDermott, et al. Enterovirus D68: a test case for the use of
917 immunological surveillance to develop tools to mitigate the pandemic potential
918 of emerging pathogens. *The Lancet Microbe*, 3(2):e83–e85, 2022.
- 919 [33] Alun L Lloyd. Realistic distributions of infectious periods in epidemic models:
920 changing patterns of persistence and dynamics. *Theoretical population biology*,
921 60(1):59–71, 2001.
- 922 [34] Jennie S Lavine, Aaron A King, and Ottar N Bjørnstad. Natural immune
923 boosting in pertussis dynamics and the potential for long-term vaccine failure.
924 *Proceedings of the National Academy of Sciences*, 108(17):7259–7264, 2011.
- 925 [35] Nicholas G Reich, Sourya Shrestha, Aaron A King, Pejman Rohani, Justin
926 Lessler, Siripen Kalayanarooj, In-Kyu Yoon, Robert V Gibbons, Donald S
927 Burke, and Derek AT Cummings. Interactions between serotypes of dengue
928 highlight epidemiological impact of cross-immunity. *Journal of The Royal Soci-*
929 *ety Interface*, 10(86):20130414, 2013.
- 930 [36] Chadi M Saad-Roy, Simon A Levin, Bryan T Grenfell, and Mike Boots. Epidemi-
931 ological impacts of post-infection mortality. *Proceedings of the Royal Society B*,
932 290(2002):20230343, 2023.
- 933 [37] Tim G Benton and Alastair Grant. Elasticity analysis as an important tool in
934 evolutionary and population ecology. *Trends in ecology & evolution*, 14(12):467–
935 471, 1999.
- 936 [38] Amy Matser, Nienke Hartemink, Hans Heesterbeek, Alison Galvani, and
937 Stephen Davis. Elasticity analysis in epidemiology: an application to tick-borne
938 infections. *Ecology Letters*, 12(12):1298–1305, 2009.
- 939 [39] Bryan T Grenfell, Ottar N Bjørnstad, and Bärbel F Finkenstädt. Dynamics of
940 measles epidemics: scaling noise, determinism, and predictability with the TSIR
941 model. *Ecological monographs*, 72(2):185–202, 2002.

- [40] Samit Bhattacharyya, Per H Gesteland, Kent Korgenski, Ottar N Bjørnstad, and Frederick R Adler. Cross-immunity between strains explains the dynamical pattern of paramyxoviruses. *Proceedings of the National Academy of Sciences*, 112(43):13396–13400, 2015.
- [41] Katharine R Dean, Fabienne Krauer, Lars Walløe, Ole Christian Lingjærde, Barbara Bramanti, Nils Chr Stenseth, and Boris V Schmid. Human ectoparasites and the spread of plague in Europe during the Second Pandemic. *Proceedings of the National Academy of Sciences*, 115(6):1304–1309, 2018.
- [42] Margarita Pons-Salort and Nicholas C Grassly. Serotype-specific immunity explains the incidence of diseases caused by human enteroviruses. *Science*, 361(6404):800–803, 2018.
- [43] Sarah Cobey and Edward B Baskerville. Limits to causal inference with state-space reconstruction for infectious disease. *PloS one*, 11(12):e0169050, 2016.
- [44] P Rohani, CJ Green, NB Mantilla-Beniers, and Bryan T Grenfell. Ecological interference between fatal diseases. *Nature*, 422(6934):885–888, 2003.
- [45] Sema Nickbakhsh, Colette Mair, Louise Matthews, Richard Reeve, Paul CD Johnson, Fiona Thorburn, Beatrix Von Wissmann, Arlene Reynolds, James McMenamin, Rory N Gunson, et al. Virus–virus interactions impact the population dynamics of influenza and the common cold. *Proceedings of the National Academy of Sciences*, 116(52):27142–27150, 2019.
- [46] Neil Ferguson, Roy Anderson, and Sunetra Gupta. The effect of antibody-dependent enhancement on the transmission dynamics and persistence of multiple-strain pathogens. *Proceedings of the National Academy of Sciences*, 96(2):790–794, 1999.
- [47] Public Health Agency of Canada. Respiratory virus detections in Canada. 2024. <https://www.canada.ca/en/public-health/services/surveillance/respiratory-virus-detections-canada.html>. Accessed Sep 4, 2024.
- [48] Centre for Health Protection. Detection of pathogens from respiratory specimens. 2024. <https://www.chp.gov.hk/en/statistics/data/10/641/642/2274.html>. Accessed Nov 20, 2024.
- [49] Centre for Health Protection. Detection of gastroenteritis viruses from faecal specimens. 2024. <https://www.chp.gov.hk/en/statistics/data/10/641/717/3957.html>. Accessed Nov 20, 2024.
- [50] Korea Disease Control and Prevention Agency. Acute respiratory infection, Sentinel surveillance infectious diseases. 2024. <https://dportal.kdca.go.kr/pot/is/st/ari.do>. Accessed Nov 20, 2024.

- 979 [51] Edward Goldstein, Sarah Cobey, Saki Takahashi, Joel C Miller, and Marc Lip-
980 sitch. Predicting the epidemic sizes of influenza A/H1N1, A/H3N2, and B: a
981 statistical method. *PLoS medicine*, 8(7):e1001051, 2011.
- 982 [52] Karline Soetaert, Thomas Petzoldt, and R Woodrow Setzer. Solving differential
983 equations in R: package deSolve. *Journal of statistical software*, 33:1–25, 2010.
- 984 [53] R Core Team. *R: A Language and Environment for Statistical Computing*. R
985 Foundation for Statistical Computing, Vienna, Austria, 2023.