

Data-driven design of targeted gene panels for estimating immunotherapy biomarkers

Jacob R. Bradley, Timothy I. Cannings

January 2021



THE UNIVERSITY
of EDINBURGH



Abstract

- ❶ Exome-wide biomarkers such as tumour mutation burden (TMB) are useful predictors of response to immunotherapy.
- ❷ While whole-exome sequencing directly measures TMB, its cost prevents it from being standard-of-care.
- ❸ We develop a data-driven framework both for selecting targeted gene panels and for using them to intelligently estimate immunotherapy biomarkers.
- ❹ To do this, we utilise an exome-wide generative model of mutation, whose structure can be chosen to reflect biological assumptions.

Outline

1 Biological/Clinical Background

- Cancer and immunotherapy
- Exome-wide biomarkers
- Targeted gene panels


2 Generative/predictive modelling framework

- Generative models: what and why?
- From generative models to predictive models


3 Application to non-small cell lung cancer

Cancer is a disease of the genome

Cancer arises when DNA in cells changes (mutates) [1].




(a) Non-cancer cell: Normal DNA.




(b) Cancer cell: Mutated DNA.

Immunotherapy enables the immune system

Immunotherapy 'releases the brakes' on the immune system in order for it to attack tumours [2].



(a) Low-damage cell: **unrecognisable** to immune system.



(b) High-damage cell: **recognisable** to the immune system.

However, the immune system can only attack tumours that it recognises!

Tumour mutation burden stratifies patients for immunotherapy response

As a simple proxy for likelihood of immune response, we can use **tumour mutation burden**: the total count of non-synonymous mutations throughout the tumour exome [3, 4].

Advantages:


- ❶ Calculated from DNA sequencing only (compatible with liquid biopsy [5]).
- ❷ Relevant across cancer types.

Disadvantages:


- ❶ Requires Whole-Exome Sequencing (WES) to measure directly.
- ❷ Fails to incorporate other features of the immune environment.

Targeted panels make genomic biomarkers viable

Rather than the entire exome/genome, targeted gene panels sequence a **subset** of genes.



(a) Whole-exome sequencing: **all genes** sequenced.



(b) Targeted panel sequencing: **subset** of genes sequenced.

Problem Statement

Aims:

- ① Select a discrete set of genes to form a panel (of pre-specified cost).
- ② Estimate TMB from panel sequencing data.

Mutations and biomarkers: some notation

We'll use i , g , and s to refer to:

- 1 A sample i (ranging from $i = 1$ to $i = N$),
- 2 A gene g (belonging to some index set G),
- 3 A variant type s (for example synonymous/non-synonymous).

Finally, we write:

- 4 M_{igs} , to refer to a mutation count of a particular sample, gene, and mutation type.

We then can define the TMB of the i th sample via:

$$T_i = \sum_{\substack{g \in G \\ s: \text{non-synonymous}}} M_{igs}.$$

What is a generative model of mutation?

A generative model of mutation attempts to capture the **underlying distribution** of variants across the entire exome/genome.

A simple example (Poisson):

$$M_{igs} \sim \text{Poisson}(\lambda_{gs}) \quad (1)$$

Why use generative models of mutation?

- ❶ Utilise known biology to inform the structure of the model.
- ❷ (Sometimes) make interpretable inferences.
- ❸ Generate trust that we've thought about it!

Why use generative models of mutation?

- ❶ Utilise known biology to inform the structure of the model.

Choice of distribution, exome organisation, parameter space.

- ❷ (Sometimes) make interpretable inferences.
- ❸ Generate trust that we've thought about it!

Why use generative models of mutation?

- ❶ Utilise known biology to inform the structure of the model.
- ❷ (Sometimes) make interpretable inferences.

Parameter estimates, post-model inference.

- ❸ Generate trust that we've thought about it!

Generative models to predictive models

Aims:

- 1 Select a discrete set of genes to form a panel (of pre-specified cost).
- 2 A mechanism for estimating TMB from panel sequencing data.

Proposed mechanism: Linear estimator.

$$\hat{T} = \sum_{g,s} w_{gs} M_{igs} \quad (2)$$

Optimisation procedure:

$$\hat{w} := \arg \min_w \{ \mathbb{E}[(\hat{T}_i - T_i)^2] + \lambda |w| \}, \quad (3)$$

where λ is a penalty specifying panel cost, and the expectation \mathbb{E} is over the distribution learned by our generative model.

NSCLC: Generative Model Structure




Figure: Generative model overview, as applied to NSCLC dataset [6].

NSCLC: Poisson-based generative model fit




Figure: Generative model parameters. [7]

NSCLC: Poisson-based generative model fit




Figure: Generative model parameters (indel-specific) [7].

NSCLC: Estimation performance




Figure: Predictive model performance: comparison with commercial panels and other predictive methods [7, 8].

References I

- [1] Douglas Hanahan and Robert A. Weinberg. Hallmarks of Cancer: The Next Generation. *Cell*, 144(5):646–674, March 2011.
- [2] Drew M. Pardoll. The blockade of immune checkpoints in cancer immunotherapy. *Nature reviews. Cancer*, 12(4):252–264, March 2012.
- [3] Jiaxin Zhu, Tiantian Zhang, Jiahao Li, Junming Lin, Wenhua Liang, Wenjie Huang, Ning Wan, and Jie Jiang. Association Between Tumor Mutation Burden (TMB) and Outcomes of Cancer Patients Treated With PD-1/PD-L1 Inhibitions: A Meta-Analysis. *Frontiers in Pharmacology*, 10, June 2019.
- [4] Dedong Cao, Huilin Xu, Ximing Xu, Tao Guo, and Wei Ge. High tumor mutation burden predicts better efficacy of immunotherapy: a pooled analysis of 103078 cancer patients. *Oncoimmunology*, 8(9):e1629258, 2019.

References II

- [5] David R. Gandara, Sarah M. Paul, Marcin Kowanetz, Erica Schleifman, Wei Zou, Yan Li, Achim Rittmeyer, Louis Fehrenbacher, Geoff Otto, Christine Malboeuf, Daniel S. Lieber, Doron Lipson, Jacob Silterra, Lukas Amler, Todd Riehl, Craig A. Cummings, Priti S. Hegde, Alan Sandler, Marcus Ballinger, David Fabrizio, Tony Mok, and David S. Shames. Blood-based tumor mutational burden as a predictor of clinical benefit in non-small-cell lung cancer patients treated with atezolizumab. *Nature Medicine*, 24(9):1441–1448, 2018.
- [6] Jacob R. Bradley. Immune Checkpoint BioMarkers (R Package), January 2021.
- [7] Jacob R. Bradley and Timothy I. Cannings. Data-driven design of targeted gene panels for estimating immunotherapy biomarkers, January 2021.
https://github.com/cobrbra/TargetedPanelEstimation_Paper.

References III

- [8] Lijing Yao, Yao Fu, Marghoob Mohiyuddin, and Hugo Y. K. Lam. ecTMB: a robust method to estimate and classify tumor mutational burden. *Scientific Reports*, 10(1):1–10, March 2020. Number: 1 Publisher: Nature Publishing Group.