

ForageModv02 sensitivity analysis

For the sensitivity analysis, we examined how forager burning probabilities and forager movement was affected by those parameters not studied in the paper. Specifically, we looked at varying the available energy in the environment (*veg-type-modifier*) and the thresholds/costs of non-disturbance forager behaviors. To do so, we allowed for essentially unlimited burning behaviors by holding *burn-cost*, *burnt-neighbor-limit*, and *burn-veg-type-threshold* constant. We also tested with one type of environmental scenario - starting in a productive environment and cycling the environment every 100 ticks - and one type of movement model - random walking. We tested the effects of the parameters listed below at a low, medium, and high value.

```
## $natural_ignition
## [1] 0 0.5
## Levels: 0 0.5
##
## $forager_moves
## [1] 5 10 15
## Levels: 5 10 15
##
## $veg_type_modifier
## [1] 200 100 300
## Levels: 100 200 300
##
## $forager_energy_requirement
## [1] 2000 1000 1500
## Levels: 1000 1500 2000
##
## $movement_cost
## [1] 150 50 100
## Levels: 50 100 150
##
## $reproduction_threshold
## [1] 5000 4000 6000
## Levels: 4000 5000 6000
##
## $reproduction_cost
## [1] 1000 500 1500
## Levels: 500 1000 1500
```

What parameters result in population dying off?

```
all.exp = unique(data$exp)

end.tick = unique(data$`tick-limit`)
fd = data %>%
  filter(ticks == end.tick)
finished.exp = unique(fd$exp)

finished.data = data %>%
```

```

  filter(exp %in% finished.exp)
unfinished.data = data %>%
  filter(!(exp %in% finished.exp))
length(unique(unfinished.data$exp))/length(unique(data$exp))

```

```
## [1] 0.5071331
```

Over half of the model runs performed in the sensitivity analysis do not sustain a population to the end of model run (ticks = 500).

```


max.ticks = unfinished.data %>%
  group_by_at(c("exp", parameters)) %>%
  summarize(max.tick = max(ticks))

# lapply(max.ticks[,c(reg.parameters)], summary)

rt.labs = c("repro threshold: 4000", "repro threshold: 5000",
  "repro threshold: 6000")
names(rt.labs) = c(4000, 5000, 6000)
rc.labs = c("repro cost: 500", "repro cost: 1000", "repro cost: 1500")
names(rc.labs) = c(500, 1000, 1500)
fm.labs = c("moves: 5", "moves: 10", "moves: 15")
names(fm.labs) = c(5, 10, 15)
mc.labs = c("move cost: 50", "move cost: 100", "move cost: 150")
names(mc.labs) = c(50, 100, 150)

ggplot(max.ticks) + geom_count(aes(x = veg_type_modifier, y = forager_energy_requirement,
  color = veg_type_modifier)) + facet_grid(forager_moves ~ reproduction_threshold + reproduction_cost,
  labeller = labeller(forager_moves = fm.labs, movement_cost = mc.labs,
    reproduction_threshold = rt.labs, reproduction_cost = rc.labs)) +
  theme(strip.text = element_text(size = 5), legend.position = "bottom")

```



n ● 5 ● 10 ● 15 ● 20 veg_type_modifier ● 100 ● 200 ● 300

The above count graph shows those model runs that do not sustain a population until the end of model run. We can clearly see that limited forager mobility (*forager-moves* = 5) and high movement cost (*movement-cost* = 150), low resource availability (*veg-type-modifier* = 100), and high cost of foraging (*forager-energy-requirement* = 2000) result in population extinction. High reproduction costs and thresholds also can lead to populations dying off before the end of model run. For the final experiments, we chose parameters that allow for populations to frequently be sustained until the end of model run.


When population is maintained, how do the parameters affect this?

```
finished.data$pop.dens = finished.data$pop.count/(51 * 51)

er.labs = c("energy requirement: 1000", "energy requirement: 1500",
"energy requirement: 2000")
names(er.labs) = c(1000, 1500, 2000)


ggplot(finished.data) + geom_smooth(aes(x = ticks, y = pop.count,
color = veg_type_modifier)) + facet_grid(forager_energy_requirement ~
reproduction_threshold + reproduction_cost, labeller = labeller(forager_energy_requirement = er.labs,
reproduction_threshold = rt.labs, reproduction_cost = rc.labs)) +
```

```
theme(strip.text = element_text(size = 5), legend.text = element_text(size = 5),
      legend.position = "bottom", axis.text.x = element_text(size = 5))
```



Access to energy is a critical determinant of population survival and population size. High amounts of energy in the environment (*veg_type_modifier* = 300) and low forager/reproduction costs (top left graphs) result in very large populations, which in turn have high computing requirements for the model. For this reason, we intentionally chose some values for the final experiments that keep the population smaller.

How do the parameters affect burning probabilities?



Across all parameters, burning probability increases. We look at the final average burning probability to compare parameter effects. Specifically, we test for significant differences in resulting burning probabilities using Mann-Whitney (two-sample Wilcoxon) tests. For all tests, we look at the differences separated out by foraging costs (*forager-energy-requirement*) and energy required for reproduction (*reproduction-threshold*).


```
end.data = data %>%
  filter(ticks == 500)

ggboxplot(end.data, x = "veg_type_modifier", y = "mean.burn.prob",
  color = "veg_type_modifier", facet.by = c("reproduction_threshold",
    "forager_energy_requirement"), short.panel.lab = FALSE) +
  stat_compare_means(comparisons = list(c("100", "200"), c("200",
```

```

"300"), c("100", "300")), label = "p.signif", vjust = 0.6,
hide.ns = T, method = "wilcox.test") + theme(strip.text = element_text(size = 6),
legend.title = element_text(size = 6), legend.position = "none")

```




First, what is obvious is that with high enough forager energy requirements, there need to be enough resources available (*veg-type-modifier*) in the environment for populations to survive. With low forager energy requirements, the differences introduced by changes in *veg-type-modifier* are larger. When forager energy requirements are higher, these differences, while still significant, are not as large, particularly at lower reproduction thresholds. For this reason, we preferred middle values of *forager-energy-requirement*, *reproduction-threshold*, and *veg-type-modifier* for the final experiments.

```


ggboxplot(end.data, x = "forager_moves", y = "mean.burn.prob",
color = "forager_moves", facet.by = c("reproduction_threshold",
"forager_energy_requirement"), short.panel.lab = FALSE) +
stat_compare_means(comparisons = list(c("5", "10"), c("10",
"15"), c("5", "15")), label = "p.signif", vjust = 0.6,
hide.ns = T, method = "wilcox.test") + theme(strip.text = element_text(size = 6),
legend.title = element_text(size = 6), legend.position = "none")

```




An alternative population maintenance strategy to enough energy availability in the environment is that agents must be able to move more to access more resources (*forager-moves*). Again, here we see smaller significant differences in final burning probability between the middle and high value of *forager-moves* across all energy requirements/thresholds.

```
ggboxplot(end.data, x = "movement_cost", y = "mean.burn.prob",
  color = "movement_cost", facet.by = c("reproduction_threshold",
    "forager_energy_requirement"), short.panel.lab = FALSE) +
  stat_compare_means(comparisons = list(c("50", "100"), c("100",
    "150"), c("50", "150")), label = "p.signif", vjust = 0.6,
    hide.ns = T, method = "wilcox.test") + theme(strip.text = element_text(size = 6),
    legend.title = element_text(size = 6), legend.position = "none")
```




The cost of movement (*movement-cost*) will play a role in how many patches a forager can access and/or burn during a foraging bout. We can see that the middle and high values for *movement-cost* allow for similar evolution of burning behaviors by the end of model run. It is only when foraging energy requirements are low, allowing for larger populations, that we see significant differences produced by different movement costs.

```
ggboxplot(end.data, x = "reproduction_cost", y = "mean.burn.prob",
  color = "reproduction_cost", facet.by = c("reproduction_threshold",
    "forager_energy_requirement"), short.panel.lab = FALSE) +
  stat_compare_means(comparisons = list(c("500", "1000"), c("1000",
    "1500"), c("500", "1500")), label = "p.signif", vjust = 0.6,
    hide.ns = T, method = "wilcox.test") + theme(strip.text = element_text(size = 6),
  legend.title = element_text(size = 6), legend.position = "none")
```



Changes in reproduction costs produce some significant differences, but these are not consistent across energy requirements/thresholds.


```
ggarrange(ggboxplot(end.data, x = "forager_energy_requirement",
  y = "mean.burn.prob", color = "forager_energy_requirement",
  facet.by = c("reproduction_threshold"), short.panel.lab = FALSE) +
  stat_compare_means(comparisons = list(c("1000", "1500"),
    c("1500", "2000"), c("1000", "2000")), label = "p.signif",
    vjust = 0.6, hide.ns = T, method = "wilcox.test") + theme(strip.text = element_text(size = 5),
  legend.title = element_text(size = 6), legend.position = "none",
  axis.text.x = element_text(size = 6)), ggboxplot(end.data,
  x = "reproduction_threshold", y = "mean.burn.prob", color = "reproduction_threshold",
  facet.by = c("forager_energy_requirement"), short.panel.lab = FALSE) +
  stat_compare_means(comparisons = list(c("4000", "5000"),
    c("5000", "6000"), c("4000", "6000")), label = "p.signif",
    vjust = 0.6, hide.ns = T, method = "wilcox.test") + theme(strip.text = element_text(size = 5),
  legend.title = element_text(size = 6), legend.position = "none",
  axis.text.x = element_text(size = 6)), ncol = 2, nrow = 1)
```



Finally, when comparing across different values for forager energy requirements or reproduction threshold, we see significant differences. Final burning probability increases with higher energy requirements and decreases with higher reproduction thresholds.


While altering these parameters do change the rate of increase of burning probabilities in the model, all parameter settings result in increasing burning probabilities, meaning that final result of the model will always be the same: 100% burning probability. This means that choosing the parameters for the final experiments could take into consideration other factors beside their effect on burning probability.

How do the parameters affect forager movement?



It is clear that the reproduction parameters do not have a large effect on forager movement compared to the parameters that control resource availability (*veg-type-modifier*) and foraging costs (*forager-energy-requirement*). Aside from showing what we already know (that foragers must be able to perform a high enough number of steps per foraging bouts when foraging costs are higher and resource availability is limited), the above results demonstrate that differences caused by the other parameters are minimized with lower forager movement. However, since forager movement needs to be high enough to maintain the population, we chose the middle value for the final experiments.

How do the parameters affect proportion of vegetation types across the landscape?



TODO

Conclusion

For all of the reasons outlined in the above sensitivity analysis, we use the middle values of each of the parameters tested here for the final experiments. These are:

- *forager-moves* = 10
- *movement-cost* = 100
- *veg-type-modifier* = 200
- *forager-energy-requirement* = 1500
- *reproduction-threshold* = 5000
- *reproduction-cost* = 1000

These values allow the model to maintain a reasonable population that does not significantly hinder model performance (i.e., does not drastically increase time complexity or memory requirements). Additionally, these parameters do meaningfully affect our variables of interest - burning probability and forager movement. The effects of the other model parameters are discussed in the manuscript and the supplementary material.

References

Package	Version	Citation
base	4.4.1	(1)
betareg	3.2.1	(2); (3); (4)
fitdistrplus	1.2.1	(5)
formatR	1.14	(6)
ggpubr	0.6.0	(7)
ggsignif	0.6.4	(8)
gtsummary	2.0.2	(9)
here	1.0.1	(10)
jtools	2.3.0	(11)
knitr	1.48	(12); (13); (14)
marginaleffects	0.23.0	(15)
mgcv	1.9.1	(16); (17); (18); (19); (20)
rcompanion	2.4.36	(21)
rmarkdown	2.28	(22); (23); (24)
tidyverse	2.0.0	(25)

1. R Core Team, *R: A language and environment for statistical computing* (R Foundation for Statistical Computing, 2024).
2. F. Cribari-Neto, A. Zeileis, Beta regression in R. *Journal of Statistical Software* **34**, 1–24 (2010).
3. B. Grün, I. Kosmidis, A. Zeileis, Extended beta regression in R: Shaken, stirred, mixed, and partitioned. *Journal of Statistical Software* **48**, 1–25 (2012).
4. I. Kosmidis, A. Zeileis, “Extended-support beta regression for $[0, 1]$ responses” (arXiv.org E-Print Archive, 2024).
5. M. L. Delignette-Muller, C. Dutang, fitdistrplus: An R package for fitting distributions. *Journal of Statistical Software* **64**, 1–34 (2015).
6. Y. Xie, *formatR: Format r code automatically* (2023).
7. A. Kassambara, *ggpubr: “ggplot2” based publication ready plots* (2023).

8. A.-E. Constantin, I. Patil, ggsignif: R package for displaying significance brackets for “ggplot2”. *PsyArxiv* (2021). <https://doi.org/10.31234/osf.io/7awm6>.
9. D. D. Sjoberg, K. Whiting, M. Curry, J. A. Lavery, J. Larmarange, Reproducible summary tables with the gtsummary package. *The R Journal* **13**, 570–580 (2021).
10. K. Müller, *here: A simpler way to find your files* (2020).
11. J. A. Long, *jtools: Analysis and presentation of social scientific data* (2022).
12. Y. Xie, “knitr: A comprehensive tool for reproducible research in R” in *Implementing Reproducible Computational Research*, V. Stodden, F. Leisch, R. D. Peng, Eds. (Chapman; Hall/CRC, 2014).
13. Y. Xie, *Dynamic documents with R and knitr*, 2nd Ed. (Chapman; Hall/CRC, 2015).
14. Y. Xie, *knitr: A general-purpose package for dynamic report generation in r* (2024).
15. V. Arel-Bundock, N. Greifer, A. Heiss, How to interpret statistical models using `marginalEffects` in R and Python. *Journal of Statistical Software* (Forthcoming).
16. S. N. Wood, Thin-plate regression splines. *Journal of the Royal Statistical Society (B)* **65**, 95–114 (2003).
17. S. N. Wood, Stable and efficient multiple smoothing parameter estimation for generalized additive models. *Journal of the American Statistical Association* **99**, 673–686 (2004).
18. S. N. Wood, Fast stable restricted maximum likelihood and marginal likelihood estimation of semi-parametric generalized linear models. *Journal of the Royal Statistical Society (B)* **73**, 3–36 (2011).
19. S. N. Wood, N., Pya, B. S"afken, Smoothing parameter and model selection for general smooth models (with discussion). *Journal of the American Statistical Association* **111**, 1548–1575 (2016).
20. S. N. Wood, *Generalized additive models: An introduction with r*, 2nd Ed. (Chapman; Hall/CRC, 2017).
21. S. S. Mangiafico, *rcompanion: Functions to support extension education program evaluation* (Rutgers Cooperative Extension, 2024).
22. Y. Xie, J. J. Allaire, G. Golemund, *R markdown: The definitive guide* (Chapman; Hall/CRC, 2018).
23. Y. Xie, C. Dervieux, E. Riederer, *R markdown cookbook* (Chapman; Hall/CRC, 2020).
24. J. Allaire, *et al.*, *rmarkdown: Dynamic documents for r* (2024).
25. H. Wickham, *et al.*, Welcome to the tidyverse. *Journal of Open Source Software* **4**, 1686 (2019).