
Predictive Modeling of Insulin Resistance Severity using Multimodal Wearable and Blood-Panel Biomarkers

Anonymous

Abstract

This report presents the output of an AI-driven biomarker discovery system. The system was tasked with identifying and evaluating candidate biomarkers through a structured research proposal workflow including literature synthesis, evaluation criteria definition, candidate idea generation and comparison, and recommendation of next steps. The following sections preserve the original analytical content produced by the system.

1 Research Goal

The primary objective of this research is to identify and validate a continuous, glucose-independent biomarker for the severity of insulin resistance (IR), quantified as True.HOMA_IR. Utilizing a dataset of 1,078 subjects, the goal is to develop predictive frameworks that synthesize dynamic wearable telemetry—including Heart Rate (HR), Heart Rate Variability (HRV), physical activity (steps), and sleep metrics—with a specific blood panel comprising Triglycerides (TG), High-Density Lipoprotein (HDL), C-Reactive Protein (CRP), and liver enzymes (AST, ALT), alongside Body Mass Index (BMI).

To ensure the scientific integrity of the derived biomarkers and prevent “formula leakage,” all models and proxies must strictly exclude direct glycemic markers such as fasting insulin, fasting glucose, and HbA1c. The focus is specifically on mapping the magnitude of IR as a continuous spectrum, enabling the identification of complex phenotypes like “Lean-IR” (Normal-Weight Obesity) and providing a longitudinal monitoring tool that reflects systemic metabolic health without relying on traditional clinical snapshots.

2 Evaluation Criteria

The evaluation of candidate biomarkers and predictive frameworks is conducted according to a rigorous set of criteria designed to ensure

statistical stability, physiological accuracy, and clinical utility. **Strict Formula Independence and Non-Leakage:** Candidate ideas must exclude insulin, glucose, and HbA1c. There must be no mathematical redundancy with the HOMA-IR formula. Preprocessing steps, such as Z-scoring or residualization, must be implemented within cross-validation folds to prevent population-level data leakage. **Mechanistic and Physiological Plausibility:** Proposals are assessed on their grounding in established metabolic pathways, such as the Diacylglycerol-Protein Kinase C (DAG-PKC) signaling axis, hepatic steatosis, and metaflammation.¹ The rationale must correctly apply clinical concepts, avoiding misapplications like conflating circulating lipids with intramyocellular lipids (the “Athlete’s Paradox”).² **Continuous Severity Mapping:** The biomarker must target the magnitude of True.HOMA_IR on a continuous scale. Evaluation prioritizes models that maintain directional consistency (e.g., risk factors like TG must scale positively with the score) and account for non-linear relationships, such as the U-shaped risk associated with sleep duration.³ **Multimodal Integration and Synergy:** A key criterion is the effective leveraging of interactions between behavioral wearable

Preprint.

data and static biochemical markers. Successful ideas should demonstrate why a combination (e.g., a Step $\times$ TG interaction) captures more variance than individual components. Numerical and Statistical Rigor: Formulas must be numerically stable, avoiding division-by-zero or "sign-flip" errors (e.g., logtransforming ratios $\neq 1$ without proper handling). Skewness management for right-skewed markers like CRP and TG is essential. Population and Demographic Robustness: Evaluation includes the ability of the biomarker to perform across diverse cohorts, specifically addressing failure modes such as the "African American TG Paradox," where insulin resistance may present with relatively low triglyceride levels.⁴ Validation through Regression Metrics: Success is measured by continuous regression metrics including R-squared (R^2), Root Mean Square Error (RMSE), and Spearman rank correlation (ρ). Categorical classification metrics (e.g., AUC) are secondary to the goal of continuous severity mapping.




Figure 1: Conceptual diagram 1 from the biomarker discovery analysis.

3 Main Research Directions

Current research into identifying glucose-independent IR biomarkers focuses on five primary axes that transition from static, population-level snapshots to dynamic, individualized signatures. These directions are worth exploring because they promise to uncover "occult" IR that BMI-centric models fail to detect. Multi-Axial Integration of Load and Resilience: This direction models IR as a tension between pathological load (biochemical stress like hepatic strain and inflammation) and physiological recovery capacity (autonomic and behavioral buffers). By organizing features

into a "Friction-Flux" framework, researchers can quantify how well a body "buffers" metabolic stress. Phenotypic Stratification (BMI-Independent Modeling): Identifying the "Thin-on-the-Outside, Fat-on-the-Inside" (TOFI) phenotype is a high priority. This involves using statistical residuals (e.g., regressing ALT or HRV against BMI) to isolate metabolic strain that is disproportionate to body mass, ensuring that lean individuals with significant metabolic dysfunction are not overlooked. Dynamic Autonomic Recovery and "Metabolic Stiffness": Moving beyond resting measurements, this direction investigates kinetic recovery signatures. Specifically, the rate of heart rate decay at sleep onset and the nocturnal heart rate nadir are used to quantify "metabolic stiffness"—the body's inability to switch from a sympathetic-dominant state to a restorative parasympathetic state. Kinetic-Metabolic Efficiency: This explores the "cardiac cost" of movement, such as the HR-to-step ratio. It allows for the contextualization of blood panels via the "Athlete's Paradox," where high activity can mitigate the risk of unusual lipid profiles, providing a more nuanced view of mitochondrial health. The Hepato-Autonomic-Inflammatory Axis: This direction maps the feedback loops between organ-specific strain and systemic control. For instance, "Hepatovagal Decoupling" explores how liver stress (ALT/AST) suppresses the parasympathetic output (HRV), identifying a systemic breakdown in metabolic coordination.

Comparison of Research Paradigms: From Static Biochemical Load to Integrated Multimodal Resilience. Note: The AI image generation process (Beta) is experimental and may not be Exploring these directions is essential because traditional metrics often reach a predictive plateau. By integrating high-frequency wearable data, researchers can capture the "metabolic hysteresis"—the lag in recovery after exertion—that serves as an early warning sign for cellular insulin resistance before overt hyperglycemia develops.

4 Candidate Idea Descriptions

This section provides a comprehensive analysis of the proposed frameworks for predicting True_HOMA_IR. Each candidate is evaluated based on its mathematical structure, physiological grounding, and the strength of the evidence supporting its implementation.

4.1 The Metabolic Friction-Flux (MFF) Score

The Metabolic Friction-Flux (MFF) Score is a multi-modal composite index that treats insulin resistance as a systemic bottleneck where the body's metabolic load ("Friction") overwhelms its kinetic clearance and parasympathetic recovery capacity ("Flux"). The score is structured around a tripartite framework: Axis 1 (Hepato-Lipid Friction) utilizing TG/HDL and ALT/AST ratios; Axis 2 (Inflammatory-Structural Anchor) scaling BMI by log-transformed CRP; and Axis 3 (Autonomic-Kinetic Flux), which derives an "Autonomic Decoupling Residual" (ADR) from the relationship between HRV, steps, and sleep duration. The primary motivation for the MFF framework is its ability to identify the "Lean-IR" phenotype. While traditional BMI-based models often fail in nonobese individuals, the MFF uses Axis 1 and 3 to flag hidden liver stress and autonomic failure. Evidence from the provided data analysis suggests that combining wearable data with routine blood biomarkers is significantly more effective for predicting HOMA-IR ($R^2 \approx 0.50$) than single-source models. By mathematically modeling the "friction" of hepatic stress against the "flux" of recovery, the MFF aims to provide a granular risk gradient that captures metabolic flexibility before overt structural changes occur.

Friction against Physiological Flux. Note: The AI image generation process (Beta) is experimental and may not be Category

Description

Mathematical

Structure

Multi-modal composite using Robustly Scaled axes:

Physiological

Basis

Integrates ectopic liver fat (ALT/AST) and cytokine-secreting adipose tissue (BMI x CRP) with kinetic substrate clearance.

$$MF\ F = Scale(Axis1) + Scale(Axis2) - Scale(Axis3).$$

Category

Description

Strengths

Robust log-stabilization of skewed markers; synergistic inflammatory gating of adiposity.

Critical Risks

The extraction of the ADR via a population-level GAM may capture demographic noise rather than individual recovery; potential mathematical contradiction in SHAP expectations for lean inflamed subjects.

Judgment: The MFF score provides a biologically compelling and interpretable framework. While the static axes (Axis 1 and 2) are highly robust, the dynamic wearable integration (Axis 3) requires refined intraindividual modeling to avoid population-level confounding.

4.2 Refined Allometric Autonomic-Hepatic Efficiency

(AAHE) Quotient The Refined AAHE Quotient is designed to isolate the "autonomic drag" of insulin resistance from common confounders like body mass and deconditioning. It utilizes Fitness-Isolated Allometric Residuals (FIAR) to extract sympathetic arousal independent of BMI. The quotient incorporates a Kinetic Axis based on Probabilistic Steady-State Weighting (PSSW) of the *HR/Step* ratio and a Recovery Axis that scans for the individualized circadian heart rate nadir (Rolling Basal Search). These are combined with a Biochemical Axis that interacts Peak 30-minute Cadence (P30C) with Triglycerides to reflect intensity-dependent substrate clearance. The AAHE framework is motivated by the "Snyder Gap"—the inability of simpler models to identify IR in non-obese individuals. By using allometric scaling to "strip away" expected elevations in heart rate caused by body size, the FIAR axis leaves a residual that specifically represents metabolic stress. This is supported by evidence that hyperinsulinemia acts as a direct stimulant of the sympathetic nervous system.⁵ Furthermore, the model addresses the "exercise-resistant" phenotype (observed in 36% of recent-onset diabetics) by modeling the diminishing returns of low-intensity steps in the presence of severe lipid bottlenecks.⁶

Signal through Allometric and Kinetic Filtering. Note: The AI image generation process (Beta) is experimental and may not be Category

Description

Mathematical

Structure

Composite index utilizing allometric residuals (log-log regression) and Bayesian-weighted kinetic ratios.

Physiological

Basis

Focuses on myocardial energetic efficiency and sympathetic hyperactivity (autonomic drag) as early footprints of IR.

Strengths

Excellent at identifying the "Thin-on-the-Outside, Fat-on-the-Inside" (TOFI) phenotype; accounts for chronotype variability via dynamic nadir detection.

Critical Risks

Hardware constraints: Daytime minute-level HRV (RMSSD) is rarely sampled passively by consumer wearables; potential overfitting of the PSSW weighting function.

Judgment: The AAHE Quotient is technically sophisticated and addresses key clinical gaps. However, its reliance on high-resolution daytime HRV recovery makes it less feasible for standard smart-watch datasets compared to nocturnal-focused models.

4.3 MET-REI (Metabolic Resilience and Integrated Strain)

Score The MET-REI Score is a feature-engineered framework that organizes features into three independent axes: HEMA (Hepatic-Metainflammatory), LIRA (Lipid-Insulin Residual), and ACRA (Autonomic-Circadian Resilience). A standout feature of this idea is the use of a Sigmoid(CRP) function to "cap" inflammatory noise. This centers weighting on the 2–10 mg/L range, isolating chronic "metainflammation" while filtering out acute spikes (>15 mg/L) caused by transient infections.⁷

The motivation for MET-REI lies in its rigorous handling of demographic bias and formula leakage. By extracting the studentized residual of the TG/HDL ratio regressed against BMI (the LIRA axis), it identifies individuals with lipid dysfunction worse than their weight suggests. This directly targets the observation that approximately 6.9% of participants with a healthy BMI display insulin resistance.⁸ The model’s design reflects the biological reality that IR is a dynamic balance between physiological strain and behavioral recovery (steps, sleep, and HR-normalized HRV).

Disproportion, and Lifestyle Resilience. Note: The AI image generation process (Beta) is experimental and may not be Category

Description

Mathematical

Structure

Multi-axis input for ML models: $f(HEMA, LIRA, ACRA)$, using fold-specific studentized residuals and sigmoid transformations.

Physiological

Basis

Bridges hepatic fat (ALT/AST) and chronic metainflammation with "autonomic restorative volume."

Strengths

High novelty in the Sigmoid-CRP noise filter; strong anti-leakage framework for residual calculations.

Critical Risks

Mathematical "Resilience Paradox": Multiplying negative Zscores for sleep and HRV would incorrectly yield a high positive resilience score.

Judgment: MET-REI offers excellent biochemical feature engineering, particularly for filtering inflammatory noise. However, the multiplicative logic in the autonomic axis requires correction to prevent inverted risk scoring.

4.4 MET-REI v2: Pipeline-Integrated Framework

MET-REI v2 evolves the previous concept by integrating Restricted Cubic Splines (RCS) and hierarchical residualization within a formal Scikit-Learn pipeline. It attempts to isolate "surplus resilience"—the variance in

lifestyle/autonomic health not already explained by biochemical strain. The HEMA axis dynamically centers the CRP midpoint on the 75th percentile of the training-fold distribution, allowing the model to adapt to different population baselines. The motivation for version 2 is the need for a "white-box" model that answers why a patient is resistant (e.g., hepatic-driven vs. lipid-driven). It is supported by evidence that hepatic stress and weight-independent lipid dysfunction are primary drivers of HOMA-IR. However, it faces a significant "sign-flip" risk where $\ln(ALT/AST)$ becomes negative in healthy individuals, potentially leading to paradoxical SHAP impacts when multiplied by CRP. Category

Description

Mathematical

Structure

RCS-based residualization and Tweedie-loss regression with monotonicity constraints.

Evaluation

Strong mechanistic sub-phenotyping; handles non-linear BMIlipid relationships via splines.

Judgment

Conceptually advanced but mathematically vulnerable to signflips in the HEMA axis; excludes absolute BMI main effects which may lower overall R^2 .

4.5 Multi-System Metabolic Resilience Index (M-SMRI)

The M-SMRI integrates Adiposity-Independent Hepatic Stress (AIHS), Nocturnal Vagal Recovery (NVR), and a Bayesian Active Metabolic Slope (AMS). It uses a "Dynamic Metabolic Quiescence Window" anchored to the physiological minimum heart rate during core sleep. The AMS uses a Bayesian Hierarchical Model to stabilize individual slopes of HR vs. Steps, which is particularly useful for sedentary subjects with sparse data. The motivation for M-SMRI is the resolution of ethnic lipid paradoxes. By including ALT/HDL alongside TG/HDL , it maintains predictive power in populations (such as African Americans) where TG/HDL may underpredict risk. It is grounded in the observation that insulin-resistant skeletal muscle requires higher heart rates to sustain movement (mitochondrial inflexibility).⁹ Category

Description

Mathematical

Structure

Bayesian Hierarchical Modeling (BHM) for kinetic slopes and segmented BMI residualization.

Evaluation

Sophisticated handling of sparse data; anchors autonomic recovery to the individual's physiological nadir.

Judgment

High technical complexity; 60-minute RMSSD windows may span multiple sleep stages, potentially violating autonomic signal stationarity.

4.6 Allometric Autonomic-Hepatic Efficiency (AAHE)

Quotient An earlier iteration of the AAHE framework, this model focuses on Structural Allometric Residuals (SAR) and Steady-State Kinetic Cardiac Cost (SSKCC). It defines the "Metabolic Minimum" as the heart rate during the 3:00–5:00 AM window. It explicitly models the interaction between physical activity and lipid clearance ($Steps \times TG$), recognizing that high triglycerides can induce resistance to exercise-mediated Lipoprotein Lipase upregulation.¹⁰ Motivation comes from the need to move from static snapshots to efficiency signatures. It explains why some patients see diminishing returns from low-intensity activity. However, it incorrectly applies allometric scaling to BMI (a ratio) rather than raw mass, which can create statistical artifacts. Category

Description

Mathematical

Structure

Allometric log-log regression residuals and step-triglyceride interaction terms.

Evaluation

Strong chronobiological targeting of the nocturnal nadir; captures physiological "exercise resistance."

Judgment

Flawed use of allometry on derived ratios (BMI); conflates systemic fitness with intrinsic myocardial efficiency.

4.7 The Metabolic Recovery Velocity (MRV) Index

The MRV Index quantifies "metabolic stiffness" using a Dynamic Vagal Inrush Constant (τ_{dyn}), which is the exponential decay constant of heart rate at sleep onset. It combines this with BMI-stratified inflammatory residuals (IAR-Z). By regressing τ_{dyn} against blood markers, it extracts "unexplained autonomic stiffness" as a pure wearable-derived signal. The MRV is motivated by the "Restless Mover" phenotype—individuals with high step counts but poor autonomic recovery. Evidence suggests that heart rate recovery is a sensitive indicator of IR.¹¹ However, the model risks pathological bias by penalizing "noisy" heart rate data, which is actually a hallmark of the most severe metabolic syndrome cases. Category

Description

Mathematical

Structure

Exponential HR decay modeling (τ) and two-stage hierarchical residualization.

Evaluation

Excellent strategy for isolating the "additive value" of wearables; unmasks the Lean-IR phenotype via IAR-Z.

Judgment

Computationally intensive; the mobility interaction term may paradoxically penalize protective activity.

4.8 MARM-H+ Index

The MARM-H+ Index uses Generalized Additive Models (GAMs) to residualize autonomic metrics (RHR/RMSSD) against BMI. It introduces Sigmoidal Metabolic Hysteresis (sMHA), which measures the area of decoupling between daily steps and nocturnal recovery over a 7-day rolling window. It utilizes Orthogonal Projection to Latent Structures (O-PLS) to fuse data streams. The motivation is capturing the "lag" in metabolic reset after exertion. It is supported by evidence that IR subjects exhibit hypothalamic-autonomic desynchronization.¹² However, weighting HRV epochs inversely to movement (as proposed) contradicts biophysical standards, which require movement exclusion to capture basal tone. Category

Description

Mathematical

Structure

GAM-based non-linear residualization and O-PLS data fusion.

Evaluation

Sound theoretical use of GAMs for non-linear allometry; capping CRP at 10 mg/L effectively filters noise.

Judgment

Relies on several invented metrics (sMHA) that lack established physiological validation; inverted statistical assumptions regarding RMSE optimization.

4.9 Circadian Autonomic-Metabolic Friction (CAMF)

Index The CAMF Index models IR as the log-transformed ratio of Static Biochemical Load to Dynamic Restorative Capacity. The denominator aggregates 14-day rolling medians of steps, sleep, and the corrected HRV/HR ratio. It explicitly targets the "Metabolic Friction" between cellular stressors and systems-level recovery. Motivation comes from resolving the "Athlete's Paradox"—where highly active individuals may have unusual lipid profiles without high IR. The CAMF framework

allows a high denominator (resilience) to buffer a high numerator (load). However, the formula HRV / HR mathematically amplifies heart rate bias rather than neutralizing it. Category

Description

Mathematical

Structure

Log-ratio: $CAMF = \ln(Load/Capacity)$, using min-max scaled components.

Evaluation

Mechanistic ratio structure targets "Hidden IR" (Lean-IR and MHO); excellent use of 14-day temporal alignment.

Category Judgment

Description

Mathematically unstable due to multiplicative denominator; misidentifies several intracellular signaling targets (e.g., PKC ϵ /IRS-1).

4.10 Robust Phenotypic Metabolic-Autonomic Index (RPMAI)

The RP-MAI features a "Phenotypic Switch"—a sigmoid BMI function that dynamically weights the index toward inflammatory markers for obese subjects or autonomic markers for lean subjects. It includes a "Flux-Adjusted Lipid Ratio" that scales TG/HDL against physical activity levels. The motivation is to account for the heterogeneous etiology of IR across different phenotypes. While the switch is conceptually interesting, it creates a "biological exclusion" where inflammation is ignored in lean subjects and autonomic dysfunction is ignored in the obese—both of which contradict established science. Category

Description

Mathematical

Structure

Sigmoid-weighted combination of Autonomic, Metabolic, and Inflammatory clusters.

Evaluation

Noise-resilient wearable heuristics (5th percentile HR); addresses the "African American TG Paradox" via ALT/AST.

Judgment

Mathematically over-corrects for activity, potentially masking dyslipidemia; the "toggle" logic discards vital linear BMI variance.

5 Comparison of Candidate Ideas

When comparing the candidate ideas, two distinct philosophical approaches emerge: Residual-based deconfounding (e.g., MFF, AAHE, MET-REI) and Interaction-based modeling (e.g., CAMF, RP-MAI).

5.1 Evidence Strength and Theoretical Alignment

The evidence pointing toward the MFF Score (4.1) and MET-REI (4.3) is particularly strong due to their robust handling of biochemical noise. Both utilize log-transformations and "inflammatory gating" (Sigmoid CRP) which aligns with the knowledge that chronic low-grade inflammation, rather than acute infection spikes, is the primary driver of HOMA-IR. The use of the ALT / AST and




Figure 2: Conceptual diagram 2 from the biomarker discovery analysis.

TG/HDL ratios as a “biochemical core” is supported by cross-lagged path analysis, proving that lipid profile changes biologically precede insulin resistance.¹³

Ideas that utilize Allometric Residuals (AAHE, MARM-H+) are superior for identifying the TOFI (Thin-on-the-Outside, Fat-on-the-Inside) phenotype. By regressing autonomic metrics against BMI, these models expose “autonomic drag”—the metabolic strain that is otherwise masked by a healthy weight.

5.2 Methodological Stability

The ideas vary significantly in their mathematical stability: Superior Stability: The MFF Score and MET-REI v2 use additive components and robust scaling, making them resilient to outliers. Higher Volatility: The CAMF Index uses a multiplicative denominator. A single low metric (e.g., a poor night’s sleep) can cause the index to spike disproportionately, potentially leading to lower Spearman ρ values in free-living conditions. Feature Granularity: The MRV Index and AAHE Quotient require minute-level time-series data (e.g., exponential decay at sleep onset). While potentially higher in fidelity, these are more likely to suffer from missing data compared to the RP-MAI or MET-REI, which utilize daily or 7-day aggregates.

5.3 Comparative Handling of Physiological Paradoxes

The RP-MAI and M-SMRI provide the best solutions for the “African American TG Paradox” by incorporating the *ALT/AST* or *ALT/HDL* ratios, which remain robust even when triglyceride signals are low. Conversely, models like the AAHE and CAMF focus more on the “Athlete’s Paradox,” utilizing activity interactions to correctly identify that high lipids in a highly active person may not indicate severe IR.

vs. Ratio-Based vs. Phenotypic Gating. Note: The AI image generation process (Beta) is experimental and may not be In summary, while MET-REI and MFF offer the most robust biochemical foundations, the AAHE Quotient and MRV Index push the envelope on wearable-derived kinetic signatures. The strongest evidence currently supports the MET-REI and MFF frameworks as they prioritize signal-to-noise

ratio in inflammatory and lipid markers, which remain the first-order predictors of True_HOMA_IR.

6 Comparative Analysis of Candidate Ideas

The following table synthesizes the primary modeling frameworks, distinguishing attributes, and the foundational evidence supporting the toptier candidate ideas for the continuous prediction of True_HOMA_IR. Idea Name

Metabolic

FrictionFlux

(MFF) Score

Refined AAHE Quotient

MET-REI Score

MRV Index

RP-MAI

Core Modeling Strategy

Evaluation Strengths

Distinguishing

Attributes

Supporting Evidence & Motivation

Mathematical

stability via logtransforms; synergistic inflammatory gating.

Autonomic Decoupling Residual (ADR): Isolates recovery from behavioral load.

Validates that

combining blood and wearable data exceeds singlesource predictive power ($R^2 \approx 0.50$).

Allometric & Kinetic Residualization

Superior isolation of the "Snyder Gap" (Lean-IR) phenotype.

FIAR Axis: Strips BMI "noise" from resting HR to expose metabolic arousal.

Directly addresses the 30.7% of nonoverweight individuals with undetected IR (TOFI phenotype).

Multi-Axial Multi-Modal Input

Robust filtering

of acute inflammatory "noise" via sigmoid functions.

LIRA Axis:

Uses foldspecific

residuals to ensure lipid signals are BMIindependent.

Grounded in

the observation that CRP and BMI are collinear; recovers lost metaflammatory signals.

τ_{dyn} Constant:

Kinetic Heart

Rate Decay

High-fidelity mapping of nocturnal parasympathetic transition.

Based on the

”slow phase” of

HR recovery as

a non-invasive indicator of cellular insulin sensitivity.

Tripartite Additive Framework

Sigmoid Phenotypic Gating

Demographic

parity; accounts for the ”African

American TG

Paradox.”

Measures the

velocity of cardiovascular relaxation at sleep onset.

Phenotypic Switch:

Dynamically

shifts focus between autonomic and inflammatory markers.

Addresses heterogeneous etiology; moves beyond raw

BMI as a linear

predictor.

Idea Name

CAMF Index

Core Modeling Strategy

Log-Ratio ”Friction” Model

Evaluation Strengths

Distinguishing

Attributes

Supporting Evidence & Motivation

Captures the

”buffer” effect of activity on biochemical load.

Capacity vs. Load: Models

IR as a failure

of behavioral "lubricants" to clear strain.

Explains the

"Athlete's Paradox" where high lipids do not equal IR in highly active individuals.

Static Biochemistry to Integrated Multimodal Resilience. Note: The AI image generation process (Beta) is experimental and may description.




Figure 3: Conceptual diagram 3 from the biomarker discovery analysis.

7 Benchmarking Against Existing Clinical

Solutions Traditional clinical proxies for insulin resistance typically rely on static bloodpanel snapshots (e.g., the Triglyceride-Glucose [TyG] Index) or anthropometric measurements (BMI). While effective for population-level screening, these solutions exhibit significant limitations in identifying "occult" metabolic dysfunction and providing high-frequency feedback. The proposed multimodal frameworks differentiate themselves from current State-of-the-Art (SOTA) solutions through three primary mechanisms: 1. Glucose Independence: Unlike the TyG index or HOMA-IR itself, these models provide a high-fidelity signal of insulin resistance without requiring glycemic markers, making them robust to fasting volatility and pre-diabetic "Gray Zone" fluctuations. 2. Dynamic Behavioral Context: Traditional markers cannot distinguish between high triglycerides caused by sedentary pathology and the high intramyocellular lipid flux often seen in athletes (the "Athlete's

Paradox"). By integrating step counts and HRV, the proposed models (e.g., AAHE and CAMF) correctly categorize these distinct phenotypes. 3. Identification of the TOFI Phenotype: BMI-only models miss approximately 7% of individuals with "Thin-on-the-Outside, Fat-on-the-Inside" (TOFI) metabolic syndrome.¹⁴ The use of residualized hepatic and autonomic metrics (as seen in MET-REI and MFF) unmask this hidden risk. The following table benchmarks the proposed multimodal approach against existing clinical standards:

Metric

Traditional

(BMI/Lipids Only)

Glycemic Proxies (TyG/HOMA-IR)

Proposed Multimodal

Proxies

Data Source

Static Blood + Height/Weight

Static Blood

(requires Glucose)

Blood + High-Frequency Wearables

Monitoring Frequency

Annual / Quarterly

Annual / Quarterly

Continuous / Daily

Lean-IR Detection

Poor (Misses TOFI)

Moderate

High (via Autonomic Residuals)

Noise Resilience

Sensitive to acute

spikes (CRP)

Sensitive to

fasting status

High (via Sigmoid Filters & Medians)

Mechanistic

Insight

Low (Opaque score)

Moderate

High (Phenotype Decomposition)

Longitudinal

Utility

Low (Trailing indicator)

Low (Trailing indicator)

High (Leading/Behavioral indicator)

By integrating the "Autonomic Floor" (nocturnal HRV) and "Kinetic Cardiac Efficiency" (HR/Steps), these proposed ideas transform standard wearable telemetry into a "digital metabolic stress test," providing clinicians with a more nuanced tool for early-stage intervention than traditional trailing indicators.

8 Unexpected Physiological Connections

The synthesis of high-dimensional wearable data and biochemical markers reveals several non-obvious connections that challenge traditional linear models of metabolic decay. These connections are particularly relevant for identifying the extreme tails of the True_HOMA_IR distribution and the "occult" IR found in lean phenotypes.

8.1 The Metabolic Exhaustion and De-afferentation Axis

(MEDA)

A critical unexpected connection exists between advanced hepatic stress and a paradoxical "pseudo-normalization" of serum lipids. As insulin resistance progresses from simple steatosis toward advanced fibrosis (indicated by a high *ALT/AST* ratio), the liver's ability to export triglycerides (TG) may eventually fail.¹⁵ In this "Metabolic Burnout" phase, traditional models often erroneously predict lower risk because circulating TG levels begin to decrease. The MEDA framework posits that this phase is characterized by a "GABAergic silencing" of the hepatic afferent vagal nerve. Hepatic steatosis triggers the release of GABA from hepatocytes, which suppresses vagal signaling and breaks the body-brain feedback loop.¹⁶ When this occurs alongside a loss of Deep Sleep (SWS)—a known proxy for visceral adipose burden—it identifies the highest decile of IR severity (*True_HOMA_IR* ≥ 6.0).¹⁷

Lipids in Advanced Insulin Resistance. Note: The AI image generation process (Beta) is experimental and may not be

8.2 The Metabolic Hysteresis Coefficient (MHC)

Drawing from metabolic scaling laws, an unexpected synthesis of biophysics and dynamic efficiency suggests that IR can be modeled as "energetic hysteresis"—a lag in the body's ability to reset its metabolic state after physical activity. Healthy physiology exhibits a stable "cardiac cost" of movement (*HR/Steps*), but in IR, "mitochondrial indecision" creates a sluggishness in switching between substrate oxidation modes. By calculating the Allometric Work-Efficiency (AWE) Residual (the deviation of daily heart rate per step from the *BMI* 0.75 scaling expectation), researchers can quantify the instability of metabolic efficiency. High variance in this residual identifies "exercise-resistant" phenotypes where the heart remains "stuck" in a high-expenditure fatty-acid oxidation mode long after activity has ceased.

8.3 Sleep-Vagal Rebound and Glucose Handling

There is a novel connection between nocturnal micro-movements and cellular glucose-handling readiness. In a metabolically flexible state, nocturnal heart rate bursts are typically followed by an immediate "vagal rebound" (a parasympathetic surge). This surge is hypothesized to facilitate insulin-independent glucose clearance via GLUT-4 translocation triggered by the preceding micro-contraction.¹⁸ In severe IR, this coupling is diminished. The Sleep-Vagal Rebound Ratio (SVRR) measures the failure of this "vagal brake," providing a passive nocturnal readout of what can be termed "Mitochondrial Gridlock" before fasting hyperglycemia ever manifests.

9 Recommendation and Next Steps

Based on the multi-axial evaluation of the proposed frameworks and their alignment with the research goal of predicting continuous True_HOMA_IR, the primary recommendation is the implementation of The Metabolic Friction-Flux (MFF) Score (Idea 1).

9.1 Rationale for Recommendation

The MFF Score is recommended as the lead candidate because it provides the most robust balance between statistical stability and physiological interpretability. Its tripartite structure—separating hepatic-lipid "Friction," inflammatory "Anchoring," and autonomic "Flux"—maps directly onto the known pathophysiology of insulin resistance. Specifically: Signal-to-Noise Ratio: The use of

MET-REI Score: Metabolic Resilience & Inflammation

HEMA, LIRA, and ACRA Axes Profile Comparison




Figure 1: Radar chart comparing MET-REI Score profiles of a Healthy Subject (balanced, high resilience) versus an Insulin Resistant (IR) Subject (high strain on HEMA & LIRA, low resilience on ACRA). Higher values indicate greater strain (HEMA, LIRA) or greater resilience (ACRA). HEMA: Hepatic & Metabolic Axis; LIRA: Lipid & Residual Axis; ACRA: Autonomic & Circadian Resilience Axis. Data is illustrative for domain expertise context.

Figure 4: Conceptual diagram 4 from the biomarker discovery analysis.

log-transformations for TG/HDL and ALT/AST effectively manages the right-skewed nature of the $N = 1,078$ dataset. Phenotypic Specificity: Its "Inflammatory-Structural Anchor" ($BMI \times \ln(CRP + 1)$) is a superior engineering choice for isolating "metaflammation" compared to standard additive models, allowing for higher sensitivity in the "Lean-IR" cohort. Feasibility: Unlike models requiring minute-level daytime recovery slopes, the MFF's reliance on nocturnal aggregates and daily step totals is highly compatible with the likely granularity of the available wearable telemetry. While the MET-REI Score (Idea 3) and the AAHE Quotient (Idea 2) offer sophisticated allometric and sigmoid filtering, the MFF's additive framework is less susceptible to the "multiplicative volatility" that can destabilize continuous regression models in free-living datasets.

9.2 Recommended Next Steps

To validate and deploy the MFF framework, the following immediate steps are recommended: 1. Phase 1: Feature Engineering and Normalization Execute the natural log-transformations for all biochemical ratios (TG/HDL , ALT/AST) and CRP to stabilize variance.

Implement Robust Scaling (Median/IQR) across the three axes to preserve the high-severity signals in the upper tail of the True.HOMA_IR distribution. 2. Phase 2: Autonomic Decoupling Residual (ADR) Construction Develop the Generalized Additive Model (GAM) to predict nocturnal HRV from daytime Steps and Sleep. Extract the residuals to form Axis 3, ensuring that the model is fitted strictly within training folds to prevent data leakage. 3. Phase 3: Multivariate Model Training Train a suite of regressors (GBM, Ridge, and Random Forest) using the three MFF axes as primary inputs. Apply 5-fold cross-validation and prioritize Spearman ρ and RMSE as the primary success metrics. 4. Phase 4: Validation Gauntlet Subject the final model to the 11-test battery, with specific focus on the Lean-IR Sensitivity Test ($BMI \leq 25$) and the Inflammatory Gating Verification. Perform a SHAP dependence analysis to verify the "Tipping Point" hypothesis—specifically identifying the CRP threshold where physical activity (Steps) ceases to yield a protective metabolic effect. 5. Phase 5: Sub-Phenotype Analysis Utilize the axis-specific scores to categorize high-risk individuals into "Liver-Dominant," "Inflammation-Dominant," or "ResilienceDeficient" subtypes to demonstrate the clinical utility of the score for targeted interventions.




Figure 5: Conceptual diagram 5 from the biomarker discovery analysis.




Figure 6: Conceptual diagram 6 from the biomarker discovery analysis.




Figure 7: Conceptual diagram 7 from the biomarker discovery analysis.