P^{4161} S $^{2/2014}$ Acas in Software Verification

Exam

This take-l Monday 10 Nov 2014, 08:00 am and is due on Tuesday II Nov 2014, 07:59 am. We will accept plain text files, PDF files, and Isabelle theory files (.thy); submission instructions are posted on the website.

WeChat: cstutorcs http://www.cse.unsw.edu.au/~cs4161/exam.html

Many of the questions require you to write Isabelle proofs. For these questions you should be included the life of the proofs of

As usual, you may also use any lemmas in theories imported by AutoCorres: the find_theorems command helps find such lemmas. Finally, all work must be your own the exam policy is more restrictive than for assignments:

You must not discuss the exam with anyone except the lecturers of this course before the exam is due. Do not give or receive assistance.

You are allowed to use all lecture material, sindes, and assignment solutions from the web. You are also allowed to use other passive internet resources such as Google, the Isabelle tutorial or Isabelle documentation. You are not allowed to ask for assistance on mailing lists, forums, or anywhere else. You are allowed to clarify questions with the lecturers.

Hints: Each question is divided into a number of sub-questions. Many of these ask you to prove results that can, and should, be used to help prove later lemmas in that question. If you get stuck trying to prove a result, use the **sorry** command and move on to the next part of the question. This way, you can still use an earlier, unproved result to solve a later goal if needed. If you correctly prove a later sub-question using a sorried lemma from a previous question, you will still earn full marks for that later question.

1 Lam (15 marks)

Consider the consideration and the

- (b) Wha
- (c) Provide a step-by-step derivation of its type, i.e. give a pen-and-paper proof of your answer to part (b) (10 marks)

WeChat: cstutorcs

- 2 Induction (45 marks)
 - (a) A differential Exam Help

In this question we will be deriving step-by-step a different induction principle for lists that says a property holds for all lists if it holds for:

(1) the empty list (2) all lists with a single element, and (3) all lists formed by adding new elements to the front and back of an existing list for which the property holds.

To guide your proof, the development is split into the following subparts $\frac{1}{1}$

- (i) Using the **inductive_set** command, define the set pal_lists that contains: the empty list, all lists with a single element, and all lists formed by adding new elements to the front and back of an existing list in pal_lists. (3 marks)
- (ii) Prove that if a list xs is in pal_lists, then x # xs ∈ pal_lists,
 i.e. prove xs ∈ pal_lists ⇒ x # xs ∈ pal_lists. (10 marks)
- (iii) Prove that pal_lists contains all lists, i.e. prove xs ∈ pal_lists.(4 marks)
- (iv) Using this result, or otherwise, prove the induction principle above. (4 marks)
- (b) The function palindrome mechanically decides if a list is a palindrome, making use of the in-built function rev that reverses a list.

palindrome $xs \equiv rev xs = xs$

Implement palindrome using the **definition** command. (1 mark)

(c) Prove [] is a palindrome, then a = b, i.e. palin (4 m: $\frac{1}{2}$) \Rightarrow a = b

- (d) Inductive palindromes that contains all palindromes and in the inductive set command. (5 marks)
- (e) Prove that palindrome xs if and only if $xs \in palindromes$, i.e.

palind pacific pacific

(14 marks)

Assignment Project Exam Help

3 C Verification (40 marks)

We will no Email: intutores @ 163.com

- (a) After being processed by AutoCorres with the unsigned_word_abs enabled for both functions, what are the names of the functions produced by AutoCorres that represent the semantics of the divmod() and even() functions, and what are their types? Which monad has AutoCorres used to represent both of them? (5 marks)
- (b) Is divmod() guaranteed to always terminate? Explain and justify your answer. (5 marks)

(c) We c specification for the behaviour of divmod() via t] n.

if $domod \neq 0$ then n mod m else n div m

Prove that

ovalid (λ_{-} . True) (divmod' n m domod) (λ r s. r = divmod_spec n m domod)

(8 malWeChat: cstutorcs

- (d) Write and prove correct a weakest precondition rule for divmod', i.e. replace the term ?PRE in the following statement with the weakest precondition and the galantee that strongisting that prove the statement elpower correct: ovalid ?PRE (divmod' n m domod) Q (12 marks)
- (e) Prove that even (returns correct answers i.e. prove that even (returns correct answers i.e. prove ovalid (λ_{-} . True) (even n) (λ_{-} s. r = (if even n then 1 else 0))

(5 ma/Q)O: 749389476

end

(f) Is even() guaranteed to always terminate? Explain and justify your answer. (5 marks)

https://tutorcs.com

4