

程序代写代做 CS编程辅导



COM

Foundations of Computer Science

WeChat: cstutorcs

Lecture 14: Combinatorics

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



UNSW
SYDNEY

Topic 4: Probability

程序代写代做 CS编程辅导



Probability

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com [LLM] [RW] [Rosen]

Week 9	Combinatorics	Ch. 14	Ch. 5	Ch. 6, 8
Week 10	Probability	Ch 16, 17	Ch. 9	Ch. 7
Week 10	Statistics	Ch. 18	Ch. 9	Ch. 7

QQ: 749389476

<https://tutorcs.com>

Combinatorics in Computer Science

程序代写代做 CS编程辅导



Combinatorics:

- Computing cost functions in algorithmic analysis
- Identifying (in-)efficiencies in data management
- Developing effective techniques for enumerating objects
- Probability calculations

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Probability in Computer Science

Probability:

程序代写代做 CS编程辅导

- Artificial Intelligence
 - Machine Learning
 - Decision theory
 - Image processing
 - Speech recognition



WeChat: cstutorcs

- Algorithms
 - Algorithm analysis
 - Big Data sampling and analysis
- Security
 - Cryptography
 - Quantum computing

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

- Networks
 - Network traffic modelling
 - Reliability modelling

<https://tutorcs.com>

Statistics in Computer Science

程序代写代做 CS编程辅导



Statistics:

- Sampling from large data sets

- Identifying anomalies

- Making predictions

WeChat: tutormcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Outline

程序代写代做 CS编程辅导

Counting Principles

Basic Counting Rules

Basic Counting Rules: Product

WeChat: cstutorcs

Combinations and Permutations

Assignment Project Exam Help

Alternative Techniques

Email: tutorcs@163.com

Difficult Counting Problems

QQ: 749389476

<https://tutorcs.com>



Outline

程序代写代做 CS编程辅导

Counting Principles



Basic Counting Rules

Basic Counting Rules: Product

WeChat: cstutorcs

Combinations and Permutations

Assignment Project Exam Help

Alternative Techniques

Email: tutorcs@163.com

Difficult Counting Problems

QQ: 749389476

<https://tutorcs.com>

Counting Techniques

General idea: find methods, algorithms or precise formulae to count the number of elements in various sets or collections derived, in a structured way, from basic sets.



Examples

Single base set $S = \{s_1, \dots, s_n\}$, $|S| = n$; find the number of

- all subsets of S
- ordered selections of r different elements of S
- unordered selections of r different elements of S
- selections of r elements from S such that ...
- functions $S \rightarrow S$ (onto, 1-1)
- partitions of S into k equivalence classes
- graphs/trees with elements of S as labelled vertices/leaves

Example

程序代写代做 CS编程辅导

Example

A restaurant has the menu:



Starters	Courses	Desserts
Soup	Fish	Ice-cream
Bread	Beef	Fruit
	Pork	Cheese
	Chicken	

WeChat: cstutorcs

Assignment Project Exam Help

How many:

Email: tutorcs@163.com

- 3 course meals (Starter-Main-Dessert) are possible?
- 3 course meals (Any item for each course) are possible?
- 3 course meals (Any item, no duplicates) are possible?
- Meals consisting of 3 items (order is unimportant)?

QQ: 749389476

<https://tutorcs.com>

Example

程序代写代做 CS编程辅导

Example

A restaurant has the menu:



Starters	Course	Dessert
Soup	Fish	Ice-cream
Bread	Beef	Fruit
	Pork	Cheese
	Chicken	

WeChat: cstutorcs

Assignment Project Exam Help

How many:

Email: tutorcs@163.com

- Starter-Main-Dessert?
- Any item for 3 courses?
- Any item, no duplicates, for 3 courses?
- Meals of 3 items?

QQ: 749389476

<https://tutorcs.com>

Basic Counting Rules: Principles

程序代写代做 CS编程辅导

Two simple rules:



- **Union rule** (“or”): If S and T are disjoint $|S \cup T| = |S| + |T|$
- **Product rule** (“and”): $|S \times T| = |S| \cdot |T|$

These cover many examples, though the rule application is not always obvious.

WeChat: [cstutorcs](#)

Common strategies: [Assignment Project Exam Help](#)

- Direct application of the rule
- Relate unknown quantities to known quantities (e.g. $|S| + |T| = |S \cup T| + |S \cap T|$)
- Find a bijection to a set that can be counted

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Outline

程序代写代做 CS编程辅导

Counting Principles

Basic Counting Rules

Basic Counting Rules: Product

Combinations and Permutations

Alternative Techniques

Difficult Counting Problems



WeChat: cstutorcs

Assignment Project Exam Help


Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

The Union Rule

Union rule — S and T disjoint 程序代写代做 CS编程辅导


$$|S \cup T| = |S| + |T|$$

S_1, S_2, \dots, S_n pairwise disjoint ($S_i \cap S_j = \emptyset$ for $i \neq j$)

$$|S_1 \cup \dots \cup S_n| = \sum |S_i|$$

WeChat: cstutorcs

Assignment Project Exam Help

Example

How many numbers in $A = [1, 2, \dots, 999]$ are divisible by 31 or 41? Email: tutorcs@163.com

$\lfloor 999/31 \rfloor = 32$ divisible by 31 QQ: 749389476

$\lfloor 999/41 \rfloor = 24$ divisible by 41 https://tutorcs.com

No number in A divisible by both

Hence, $32 + 24 = 56$ divisible by 31 or 41

Consequences of the Union Rule

程序代写代做 CS编程辅导



Fact

For any sets X, Y, Z

$$|Y \setminus X| = |Y| - |X \cap Y|$$

WeChat: cstutorcs

$$|X \cup Y| = |X| + |Y| - |X \cap Y|$$

Assignment Project Exam Help

$$|X \cup Y \cup Z| = |X| + |Y| + |Z| - |X \cap Y| - |Y \cap Z| - |Z \cap X| + |X \cap Y \cap Z|$$

Email: tutorcs@163.com

QQ: 749389476

<https://tutores.com>

Fact

程序代写代做 CS编程辅导

- If $|S \cup T| = |S| + |T|$ then S and T are disjoint
- If $|\bigcup_{i=1}^n S_i| = \sum |S_i|$ then S_i are pairwise disjoint
- If $|T \setminus S| = |T|$ then $S \subseteq T$



These properties can serve to identify cases when sets are disjoint (resp. one is contained in the other).

WeChat: cstutorcs

Assignment Project Exam Help

Proof.

Email: tutorcs@163.com

$|S| + |T| = |S \cup T|$ means $|S \cap T| = 0$

QQ: 749389476

$|T \setminus S| = |T|$ means $|S \cap T| = |S|$ means $S \subseteq T$

http://tutorcs.com



Exercises

程序代写代做 CS编程辅导

Exercises

RW: 5.3.1 200 people swim or jog, 85 swim and 60 do both.
How many jog?



WeChat: cstutorcs

Assignment Project Exam Help

RW: 5.6.38 (Supp) There are 100 problems, 75 of which are 'easy' and 40 'important'.
What's the smallest number of easy *and* important problems?

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Outline

程序代写代做 CS编程辅导

Counting Principles

Basic Counting Rules



Basic Counting Rules: Product

WeChat: cstutorcs

Combinations and Permutations

Assignment Project Exam Help

Alternative Techniques

Email: tutorcs@163.com

Difficult Counting Problems

QQ: 749389476

<https://tutorcs.com>

The Product Rule

Product rule

程序代写代做 CS编程辅导

$$|S_1 \times \dots \times S_k| = |S_1| \cdot |S_2| \cdots |S_k| = \prod_{i=1}^k |S_i|$$

If all $S_i = S$ (the same set) and $|S| = m$ then $|S^k| = m^k$



NB

This counts the number of sequences where the first item is from S_1 , the second is from S_2 , and so on.

WeChat: cs_tutorcs

Assignment Project Exam Help

Example

Email: tutorcs@163.com

Let $\Sigma = \{a, b, c, d, e, f, g\}$

How many 5-letter words?

QQ: 749389476

<https://tutorcs.com>

$$|\Sigma^5| = |\Sigma|^5 = 7^5 = 16,807$$

How many with no letter repeated?


Product rule: Sequences of selections

Question

程序代写代做 CS编程辅导

How can we count sequences when the underlying set changes?

To count sequences  placement:

- Define an order  underlying set
- Select from $[1, n]$, where n is the size of the “remaining” set, and a selection of i represents choosing the i -th element in that set

Assignment Project Exam Help

Example

Email: tutorcs@163.com

Let $\Sigma = \{a, b, c, d, e, f, g\}$.

How many 5-letter words with no letter repeated?

QQ: 749389476

<https://tutorcs.com>

$$\prod_{i=0}^4 (|\Sigma| - i) = 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 = 2,520$$

Exercises

Exercises

程序代写代做 CS编程辅导

S, T finite. How many functions $S \rightarrow T$ are there?



RW: 5.1.19

Consider a complete graph on n vertices.

WeChat: tutorcs

(a) No. of paths of length 3

Assignment Project Exam Help

Email: tutorcs@163.com

(b) paths of length 3 with all vertices distinct

QQ: 749389476

<https://tutorcs.com>

(c) paths of length 3 with all edges distinct

Exercise

Exercise

程序代写代做 CS编程辅导

RW: 5.3.2 $S = [100 \dots 999]$, thus $|S| = 900$.

(a) How many numbers have at least one digit that is a 3 or 7?



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

(b) How many numbers have a 3 *and* a 7?

Combinatorial Symmetry

A **symmetry** of a mathematical object is a bijective mapping from the object to itself which preserves "structure".

A (combinatorial) symmetry defines an equivalence relation where the equivalence classes all have the same size.



We are often interested in counting a set "up to symmetry". That is, counting the number of equivalence classes.

This can also be stated as a constraint that identifies a specific item in each equivalence class (**symmetric constraint**).

Definition

A *k-to-1 function* is a function that maps exactly k inputs to an output.

NB

A *k-to-1 function* defines the equivalence relation of a combinatorial symmetry and vice-versa.

程序代写代做CS编程辅导

WeChat: cstutores

Assignment Project Exam Help

Email: tutores@163.com

QQ: 749389476

<https://tutores.com>

Product rule: Symmetries and duplications

程序代写代做 CS编程辅导

Question

- How can we count the number of outcomes when we have symmetric constraints?
- How can we count the number of outcomes when we have duplicates?



Example

WeChat: cstutorcs

Let $\Sigma = \{a, b, c, d, e\}$

Assignment Project Exam Help

- How many 5-letter words with no letter repeated and a before b before c ?
- How many 5-letter words can be made from a, a, a, d, e ?

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

NB

The answer will be the same.

Product rule: Symmetries and duplications

程序代写代做 CS编程辅导

- S_1 = sequences
- S_2 = symmetric
- S = sequences without symmetry



WeChat: [tutorcs](#)

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

Alternatively, $\frac{1}{|S_2|}$ of the $|S|$ sequences meet the symmetric constraint.

<https://tutorcs.com>

Product rule: Symmetries and duplications

程序代写代做 CS编程辅导

Example

Let $\Sigma = \{a, b, c, d, e\}$

How many 5-letter words with no letter repeated and a before b before c ?



Let $\Sigma' = \{a, b, c\}$.

WeChat: cstutorcs

Assignment Project Exam Help

$$S = \prod_{i=0}^4 (|\Sigma| - i) = 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 120$$

Email: tutorcs@163.com

$$S_2 = \prod_{i=0}^2 (|\Sigma'| - i) = 3 \cdot 2 \cdot 1 = 6$$

QQ: 749389476

$$\text{So } S_1 = 120/6 = 20$$

<https://tutorcs.com>

Outline

程序代写代做 CS编程辅导

Counting Principles

Basic Counting Rules

Basic Counting Rules: Product

Combinations and Permutations

Alternative Techniques

Difficult Counting Problems



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Combinatorial Objects: How Many?

permutations

程序代写代做 CS编程辅导

Ordering of all objects from a set S ; equivalently: Selecting all objects while *recognising* the order of selection.

The number of permutations of n elements is



$$n! = n \cdot (n-1) \cdots 1, \quad 0! = 1! = 1$$

WeChat: cstutorcs

r -permutations (sequences without repetition)

Assignment Project Exam Help

Selecting any r objects from a set S of size n without repetition while *recognising* the order of selection.

Email: tutorcs@163.com

Their number is

QQ: 749389476

$$(n)_r = {}^n P_r = n \cdot (n-1) \cdots (n-r+1) = \frac{n!}{(n-r)!}$$

<https://tutorcs.com>

Permutations with duplicates

程序代写代做 CS编程辅导



Example

How many anagrams of $SSSS$?

Label S's: $AS_1S_2ES_3S_4$: $6!$

In each anagram we can label the S's in $4!$ ways.

Suppose there are m anagrams. So $m \cdot 4! = 6!$, i.e. $m = \frac{6!}{4!}$

WeChat: tutormcs
Assignment Project Exam Help

Example

Email: tutormcs@163.com

Number of anagrams of MISSISSIPPI? $\frac{11!}{3!4!2!}$

QQ: 749389476

<https://tutormcs.com>

***r*-selections** (or: ***r*-combinations**)

Collecting any r distinct objects without repetition;
equivalently: selecting r objects from a set S of size n and *not* recognising the order of selection.

Their number is

$$\binom{n}{r} = \frac{(n)_r}{r!} = \frac{n \cdot (n-1) \cdots (n-r+1)}{1 \cdot 2 \cdots r}$$

WeChat: cstutorcs

NB

Assignment Project Exam Help

These numbers are usually called binomial coefficients due to
Email: tutorcs@163.com

$$(a+b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + b^n = \sum_{i=0}^n \binom{n}{i} a^{n-i} b^i$$

<https://tutorcs.com>

Also defined for any $\alpha \in \mathbb{R}$ as $\binom{\alpha}{r} = \frac{\alpha(\alpha-1) \cdots (\alpha-r+1)}{r!}$

Simple Counting Problems

Example

程序代写代做 CS编程辅导

RW: 5.1.2 Give an example of a counting problem whose answer is

(a) $(26)_{10}$

(b) $\binom{26}{10}$



Draw 10 cards from a half-deck (just black cards only)

(a) the cards are recorded in the order of appearance

(b) only the complete draw is recorded

WeChat: tutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

Examples

QQ: 749389476

- Number of edges in a complete graph K_n
- Number of diagonals in a convex polygon
- Number of poker hands
- Decisions in games, lotteries etc.

<https://tutorcs.com>

Exercises

Exercises

程序代写代做 CS编程辅导

RW: 5.1.6 From a group of 12 men and 16 women, how many committees can be formed consisting of



(a) 7 members?

(b) 3 men and 4 women?

WeChat: cstutorcs

(c) 7 women or 7 men?

Assignment Project Exam Help

RW: 5.1.7 As above, but any 4 people (male or female) out of 9 and two, Alice and Bob, unwilling to serve on the same committee.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Counting Poker Hands

程序代写代做 CS编程辅导

Exercises

RW: 5.1.15 A poker hand consists of 5 cards drawn without replacement from a standard deck of 52 cards



$\{A, 2-10, J, Q, K\} \times \{\text{club, spade, heart, diamond}\}$

WeChat: cstutorcs

(a) Number of “4 of a kind” hands (e.g. 4 Jacks)

Assignment Project Exam Help

(b) Number of non-straight flushes, i.e. all cards of same suit but *not* consecutive (e.g. 8,9,10,J,K)

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Selecting items summary

程序代写代做 CS编程辅导

Selecting k items from n items:



With replacement	Order matters	Examples	Formula
Yes	Yes	Words of length k (sequences of length k)	n^k
No	Yes	k -permutations	$(n)_k$
No	No	Subsets of size k	$\binom{n}{k}$
Yes	No	Multisets of size k	$\binom{n+k-1}{k}$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutors@163.com

QQ: 749389476

<https://tutorcs.com>

“Balls in boxes”

Have n “distinguishable” boxes. 程序代写代做 CS编程辅导

Have k balls which are

- ① Indistinguishable
- ② Distinguishable



How many ways to place balls in boxes with

- A At most one WeChat: cstutorcs
- B Any number of Assignment Project Exam Help

balls per box?

Email: tutorcs@163.com

NB

QQ: 749389476
Suppose K is a set with $|K| = k$ and N is a set with $|N| = n$:

- $2A$ counts the number of injective functions from K to N
- $2B$ counts the number of functions from K to N

https://tutorcs.com

“Balls in boxes”

程序代写代做 CS编程辅导



Case	Balls per box		Number
1A	Indist.	At most 1	$\binom{n}{k}$
1B	Indist.	Any number	$\binom{n+k-1}{k}$
2A	Dist.	At most 1	$(n)_k$
2B	Dist.	Any number	n^k

QQ: 749389476

<https://tutorcs.com>

Outline

程序代写代做 CS编程辅导

Counting Principles

Basic Counting Rules

Basic Counting Rules: Product

Combinations and Permutations

Alternative Techniques

Difficult Counting Problems



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Alternative techniques

程序代写代做 CS编程辅导



What if the current techniques are unwieldy?

Other techniques for obtaining an exact count:

- Find a different approach for counting
- Make use of symmetries
- Make use of recursion
- Write a program (running time?)

WeChat: tutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example

程序代写代做 CS编程辅导



Example

How many sequences of 15 coin flips have an even number of heads?

WeChat: cstutorcs

- Using “balls in boxes”: $\binom{15}{0} + \binom{15}{2} + \dots + \binom{15}{14}$
- Use symmetry: $\frac{1}{2} \times 2^{15}$

Assignment Project Exam Help

- Use recursion: $\text{Even}(n) = \text{Odd}(n-1) + \text{Even}(n-1);$
 $\text{Odd}(n) = \text{Even}(n-1) + \text{Odd}(n-1)$

Email: tutors@163.com

QQ: 749389476

<https://tutorcs.com>

Example

Example

程序代写代做 CS编程辅导

How many sequences of n coin flips do not contain HH ?



$$C(0)$$

$$C(1)$$

$$C(n) = C(n-1) + C(n-2) + 2^{n-2}$$

$$N(0) = 1$$

$$N(1) = 2$$

$$N(2) = 3$$

$$N(n) = N(n-1) + N(n-2)$$

WeChat: cstutorcs


Assignment Project Exam Help

Email: tutorcs1@163.com

QQ: 749389476

<https://tutorcs.com>

We can summarise all possible outcomes in a **recursive tree**



Outline

程序代写代做 CS编程辅导

Counting Principles

Basic Counting Rules

Basic Counting Rules: Product

Combinations and Permutations

Alternative Techniques

Difficult Counting Problems



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Difficult Counting Problems

程序代写代做 CS编程辅导

Example (Ramsay)



An example of a Ramsey number is $R(3, 3) = 6$, meaning that

" K_6 is the smallest complete graph such that if all edges are painted using two colours, then there must be at least one monochromatic triangle"

Assignment Project Exam Help

This serves as the basis of a game called S-I-M (invented by Simmons), where two adversaries connect six dots, respectively using blue and red lines. The objective is to *avoid* closing a triangle of one's own colour. The second player has a winning strategy, but the full analysis requires a computer program.

Email: adversaries@163.com

QQ: 749389476

<https://tutores.com>

Using Programs to Count

程序代写代做 CS编程辅导

Two dice, a red die and a black die, are rolled.

(Note: one *die*, two *dice*)



Write a program to list all pairs $\{(R, B) : R > B\}$

Similarly, for three dice, list all triples $R > B > G$

WeChat: cstutorcs

Generally, for n dice, all of which are m -sided ($n < m$), list all decreasing n -tuples

Assignment Project Exam Help

Email: tutorcs@163.com

NB

QQ: 749389476

In order to just find the number of such n -tuples, it is not necessary to list them all. One can write a recurrence relation for these numbers and compute (or try to solve) it.

<https://tutorcs.com>

Approximate Counting

程序代写代做 CS编程辅导

NB

A Count may be a precise one or an **estimate**.

The latter should be *asymptotically correct* or at least give a good *asymptotic bound*, whether upper or lower. If S is the base set, $|S| = n$ its size, and we denote by $c(S)$ some collection of objects from S we are interested in, then we seek constants a, b such that

Email: tutors@163.com

$$a \leq \lim_{n \rightarrow \infty} \frac{\text{est}(|c(S)|)}{|c(S)|} \leq b$$

QQ: 749389476

In other words $\text{est}(|c(S)|) \in \Theta(|c(S)|)$

<https://tutors.com>

