

程序代写代做 CS编程辅导



COM

Foundations of Computer Science

WeChat: cstutorcs

Lecture 15: Probability

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Outline

程序代写代做 CS编程辅导

Elementary Discrete



Independence

Infinite Sample Spaces

WeChat: cstutorcs

Recursive Probability Computations

Assignment Project Exam Help

Conditional Probability

Email: tutorcs@163.com

Indpendence, revisited

QQ: 749389476

<https://tutorcs.com>

# Outline

程序代写代做 CS编程辅导

Elementary Discrete



Independence

Infinite Sample Spaces

WeChat: cstutorcs

Recursive Probability Computations

Assignment Project Exam Help

Conditional Probability

Email: tutorcs@163.com

Indpendence, revisited

QQ: 749389476

<https://tutorcs.com>

# Elementary Probability

## Definition

程序代写代做 CS编程辅导

Sample space:



$\omega_1, \dots, \omega_n\}$

Each point represents an outcome.

Event: a collection of outcomes = subset of  $\Omega$

WeChat: cstutorcs

Probability distribution: A function  $P: \text{Pow}(\Omega) \rightarrow \mathbb{R}$  such that:

- $P(\Omega) = 1$  Email: tutorcs@163.com
- $E$  and  $F$  disjoint events then  $P(E \cup F) = P(E) + P(F)$ .  
QQ: 749389476

## Fact

<https://tutorcs.com>

$$P(\emptyset) = 0, \quad P(E^c) = 1 - P(E)$$

# Examples

程序代写代做 CS编程辅导

## Examples

Tossing a coin:  $\Omega = \{H, T\}$



$$P(H) = P(T) = 0.5$$

Rolling a die:  $\Omega = \{1, 2, 3, 4, 5, 6\}$

$$P(1) = P(2) = P(3) = P(4) = P(5) = P(6) = \frac{1}{6}$$

QQ: 749389476

<https://tutorcs.com>

# Uniform distribution

Each outcome  $\omega_i$  equally likely.

$$P(\omega_1) = \dots = P(\omega_n) = \frac{1}{n}$$

This is called a uniform probability distribution over  $\Omega$ .

## Examples

WeChat: cstutorcs

Tossing a coin:  $\Omega = \{H, T\}$

Email: tutorcs@163.com

Rolling a die:  $\Omega = \{1, 2, 3, 4, 5, 6\}$

$$P(1) = P(2) = P(3) = P(4) = P(5) = P(6) = \frac{1}{6}$$

# Computing Probabilities by Counting

程序代写代做 CS编程辅导

Computing probabilities with respect to a *uniform* distribution comes down to counting the size of the event.

If  $E = \{e_1, \dots, e_k\}$  then



$$P(E) = \sum_{i=1}^k P(e_i) = \sum_{i=1}^k \frac{1}{|\Omega|} = \frac{|E|}{|\Omega|}$$

Assignment Project Exam Help

Most of the counting rules carry over to probabilities wrt. a uniform distribution. Email: [tutorcs@163.com](mailto:tutorcs@163.com)

**Important!**

QQ: 749389476

The expression “selected at random”, when not further qualified, means:

<https://tutorcs.com>

“subject to / according to / ... a *uniform* distribution.”

# Combining events

程序代写代做 CS编程辅导

We can create complex events by combining simpler ones.  
Common constructions:



- $A$  and  $B$ :  $A \cap B$
- $A$  or  $B$ :  $A \cup B$
- Not  $A$ :  $\Omega \setminus A$
- $A$  followed by  $B$

The first three involve events from the same set of outcomes. The last may involve events from different sets of outcomes (e.g. roll die and flip coin).

<https://tutorcs.com>

# Inclusion-exclusion rule

程序代写代做 CS编程辅导

Fact



$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

WeChat: cstutorcs

$$P(A \cup B \cup C) = P(A) + P(B) + P(C)$$

Assignment Project Exam Help  
- $P(A \cap B)$  - $P(B \cap C)$  - $P(C \cap A)$

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

# Exercises

## Exercises

程序代写代做 CS编程辅导

RW: 5.2.7 Suppose an experiment leads to events  $A, B$  with probabilities  $P(A) =$   = 0.8,  $P(A \cap B) = 0.4$ .

Find

- $P(B^c)$
- $P(A \cup B)$
- $P(A^c \cup B^c)$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

RW: 5.2.8 Given  $P(A) = 0.6$ ,  $P(B) = 0.7$ , show  $P(A \cap B) \geq 0.3$

<https://tutorcs.com>

# Examples

程序代写代做 CS编程辅导

## Example

A four-digit number  is selected at random (i.e. randomly from  $[1000 \dots 9999]$ ). Find the probability  $p$  that  $n$  has each of 0, 1, 2 among its digits.

Let  $q = 1 - p$  be the complementary probability and define

**WeChat: cstutorcs**  
**Assignment Project Exam Help**  
 $A_i = \{n : \text{no digit } i\}, A_{ij} = \{n : \text{no digits } i, j\}, A_{ijk} = \{n : \text{no } i, j, k\}$

**Email: tutorcs@163.com**

Then define

**QQ: 749389476**

$T = A_0 \cup A_1 \cup A_2 = \{n : \text{missing at least one of 0, 1, 2}\}$

$S = (A_0 \cup A_1 \cup A_2)^c = \{n : \text{containing each of 0, 1, 2}\}$

**https://tutorcs.com**

# Examples

程序代写代做 CS编程辅导

## Example (cont'd)

Once we find the car



$T$ , the solution is

$$q = \frac{|T|}{9000}, \quad p = 1 - q$$

Assignment Project Exam Help

To find  $|A_i|$ ,  $|A_{ij}|$ ,  $|A_{ijk}|$  we reflect on how many choices are available for the first digit, for the second etc. A special case is the leading digit, which must be  $1, \dots, 9$

<https://tutorcs.com>

# Examples

## Example (cont'd) 程序代写代做 CS编程辅导

$$|A_0| = 9^4, \quad |A_1| = 8^4 = 8 \cdot 9^3$$

$$|A_{01}| = |A_{02}| = 8^3 = 7 \cdot 8^3$$

$$|A_{012}| = 7^4$$



WeChat: cstutorcs

$$|T| = |A_0 \cup A_1 \cup A_2|$$

$$= |A_0| + |A_1| + |A_2| - |A_0 \cap A_1| - |A_0 \cap A_2| - |A_1 \cap A_2|$$

$$+ |A_0 \cap A_1 \cap A_2|$$

$$= 9^4 + 2 \cdot 8 \cdot 9^3 - 2 \cdot 8^4 - 7 \cdot 8^3 + 7^4$$

QQ: 749389476

$$= 25 \cdot 9^3 - 23 \cdot 8^3 + 7^4 = 8850$$

<https://tutorcs.com>

$$q = \frac{8850}{9000}, \quad p = 1 - q \approx 0.01667$$

# Examples

程序代写代做 CS编程辅导

## Example

Previous example get  
having all of 0,1,2,3



Probability of an  $r$ -digit number  
having all of 0,1,2,3 digits.

We use the previous notation:  $A_i$  — set of numbers  $n$  missing digit  $i$ , and similarly for all  $A_j$ .

WeChat: cstuorcs

We aim to find the size of  $T = A_0 \cup A_1 \cup A_2 \cup A_3$ , and then to compute  $|S| = 9 \cdot 10^{r-1}$ .

Email: tutorcs@163.com

QQ: 749389476

+ sum of  $|A_i \cap A_j \cap A_k|$

- sum of  $|A_i \cap A_j \cap A_k \cap A_l|$

# Exercises

程序代写代做 CS编程辅导

## Exercises

RW: 5.6.38 (Supp) Consider  $n$  problems, 75 are 'easy' and 40 'important'.

(b)  $n$  problems chosen randomly. What is the probability that all  $n$  are important?

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



WeChat: **tutorcs**

What's app: **tutorcs**

# Exercises

## Exercises

程序代写代做 CS编程辅导

- RW: 5.2.3 A 4-letter word is selected at random from  $\Sigma^4$ , where  $\Sigma = \{a, b, c, d, e\}$ . What is the probability that
- (a) the letters in the word are distinct?



WeChat: cstutorcs

- (b) there are no vowels (“a”, “e”) in the word?

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

- (c) the word begins with a vowel?

<https://tutorcs.com>

# Outline

程序代写代做 CS编程辅导

Elementary Discrete



Independence

Infinite Sample Spaces

WeChat: cstutorcs

Recursive Probability Computations

Assignment Project Exam Help

Conditional Probability

Email: tutorcs@163.com

Indpendence, revisited

QQ: 749389476

<https://tutorcs.com>

# Unifying sets of outcomes

To combine events from different sets of outcomes we unify the sample space using the ~~product space~~:  $\Omega_1 \times \Omega_2 \times \dots \times \Omega_n$ .

## Example

Flipping a coin and rolling a die:



$$\Omega_1 = \{\text{heads, tails}\} \quad \Omega_2 = \{1, 2, 3, 4, 5, 6\}$$

Assignment Project Exam Help

$$\Omega = \Omega_1 \times \Omega_2 = \{(\text{heads}, 1), (\text{heads}, 2), \dots\}$$

Email: [tutorcs@163.com](mailto:tutorcs@163.com)  
QQ: 749389476

NB

<https://tutorcs.com>

*This approach can also be used to model sequences of outcomes.*

# Events in the product space

Events are lifted into the product space by restricting the appropriate co-ordinate. E.g.  $A \subseteq \Omega_1$  translates to  $A' = A \times \Omega_2 \times \dots \times \Omega_n$

## Example



Coin shows heads and die shows an even number:

WeChat: cstutorcs  
 $\Omega_1 = \{\text{heads, tails}\}$        $A = \{\text{heads}\}$   
 $\Omega_2 = \{1, 2, 3, 4, 5, 6\}$        $B = \{2, 4, 6\}$

Email: tutorcs@163.com

$$\Omega = \Omega_1 \times \Omega_2 = \{(\text{heads}, 1), (\text{heads}, 2), \dots\}$$

QQ: 749389476

$$A' = A \times \Omega_2 \quad B' = \Omega_1 \times B$$

“ $A$  and  $B$ ” or “ $A$  followed by  $B$ ” corresponds to:

$$A' \cap B' = (A \times \Omega_2) \cap (\Omega_1 \times B) = A \times B$$

# Probability in the product space

NB

程序代写代做 CS编程辅导

Cannot assume that  $P(A \times B) = P(A)P(B)$



## Example

Toss two coins.

- A: First coin shows heads
- B: Both coins show tails

$$\Omega_1 = \{H, T\}$$

WeChat: cstutorcs  
Assignment Project Exam Help

$$A = \{H\}$$

$$\Omega_2 = \{HH, HT, TH, TT\}$$

$$B = \{TT\}$$

$$A' = \{(H, HH), (H, HT), (H, TH), (H, TT)\}$$

$$B' = \{(H, TT), (T, TT)\}$$

$$A' \cap B' = A \times B = \{(H, TT)\}$$

Email: tutorcs@163.com  
QQ: 749389476

<https://tutorcs.com>

$$P(A) = \frac{1}{2} \quad P(B) = \frac{1}{4} \quad P(A' \cap B') = 0$$

# Product distribution

程序代写代做 CS编程辅导

Given probability distributions on the component spaces, there is a natural probability distribution on the product space:

$$P(E_1 \times E_2 \times \dots) = P_1(E_1) \cdot P_2(E_2) \cdots P_n(E_n)$$

WeChat: cstutorcs

Intuitively, the probability of an event in one dimension is not affected by the outcomes in the other dimensions.

Email: tutorcs@163.com

Fact

QQ: 749389476

If the  $P_i$  are uniform distributions then so is the product distribution.

<https://tutorcs.com>

# Independence

Informally, events are *independent* if the outcomes in one do not affect the outcomes in the other.

More generally, we discuss dependence on events of the **same** sample space.



## Definition

$A$  and  $B$  are (*stochastically*) **independent** (notation:  $A \perp B$ ) if  
 $P(A \cap B) = P(A) \cdot P(B)$

WeChat: cstutorcs  
**Assignment Project Exam Help**

## NB

*Informal notion of independence corresponds to the stochastic independence of the "lifted" events  $A'$  and  $B'$*

## Important!

<https://tutorcs.com>

Unless specified otherwise, we assume independence when unifying events (where appropriate).

# Independence of multiple events

Independence of  $A_1, \dots, A_n$  (程序代写代做 CS 编程辅导)  $(A_1 \perp\!\!\! \perp A_2 \perp\!\!\! \perp \dots \perp\!\!\! \perp A_n)$

$$P(A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_k}) = P(A_{i_1}) \cdot P(A_{i_2}) \cdots P(A_{i_k})$$



This is often called (for emphasis) a *full* independence

WeChat: cstutorcs

Pairwise independence is a weaker concept.

Assignment Project Exam Help

## Example

Toss of two coins

$A = \langle$ first coin H $\rangle$

$B = \langle$ second coin H $\rangle$

$C = \langle$ exactly one H $\rangle$

Email: tutorcs@163.com

$$\left. \begin{array}{l} P(A) = P(B) = P(C) = \frac{1}{2} \\ P(A \cap B) = P(A \cap C) = P(B \cap C) = \frac{1}{4} \end{array} \right\}$$

QQ: 749389476

$$\text{However: } P(A \cap B \cap C) = 0$$

<https://tutorcs.com>

One can similarly construct a set of  $n$  events where any  $k$  of them are independent, while any  $k + 1$  are dependent (for  $k < n$ ).

# Example: Dependent events

程序代写代做 CS编程辅导

## Example

Basic non-independent events (assuming non-trivial probabilities)



- $A \subseteq B$

WeChat: cstutorcs

- $A \cap B = \emptyset$

Assignment Project Exam Help

- Any pair of one-point events  $A = \{x\}$ ,  $B = \{y\}$ :  
either  $x = y$  and  $A \subseteq B$

Email: tutorcs@163.com

or  $x \neq y$  and  $A \cap B = \emptyset$

QQ: 749389476

<https://tutorcs.com>

# Exercise

## Exercise

程序代写代做 CS编程辅导

RW: 9.1.25

Does  $A \perp B \perp C$  imply  $(A \cap B) \perp (A \cap C)$  ?



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

## Example: Sequences of independent events

### Example

程序代写代做 CS编程辅导

Team A has probability  $p = 0.5$  of winning a game against B.

What is the probability of A winning a best-of-seven match if

- a A already won three games?
- b A already won two games?
- c A already won two out of the first three games?

(a) Sample space  $S$  — 6-sequences, formed from wins (W) and losses (L)

Email:[tutorcs@163.com](mailto:tutorcs@163.com)

Favourable sequences — those with three to six W

$$|F| = \binom{6}{3} + \binom{6}{4} + \binom{6}{5} + \binom{6}{6} = 20 + 15 + 6 + 1 = 42$$

Therefore  $P_{0.5} = \frac{42}{64} \approx 66\%$

(b) Sample space  $S$  — 5-sequences of W and L

# Binomial distribution

程序代写代做 CS编程辅导

A useful corollary:

## Fact

*In a sequence of  $n$  independent trials, each with a probability of  $p$  of success:*



WeChat: cstutorcs

$$P(\text{exactly } k \text{ successes}) = \binom{n}{k} p^k q^{n-k}$$

Assignment Project Exam Help

where  $q = (1 - p)$ . Email: [tutorcs@163.com](mailto:tutorcs@163.com)

NB

QQ: 749389476

This leads to a probability distribution on sequences of outcomes, known as the **binomial distribution**.

<https://tutorcs.com>

# Exercises

## Exercise

程序代写代做 CS编程辅导

RW: 5.2.11 Two dice, a red die and a black die, are rolled.

What is the probability

(a) the sum of the values is even?



WeChat: cstutorcs

(b) the number on the red die is bigger than on the black die?

Email: tutorcs@163.com

QQ: 749389476

(c) the number on the black die is twice the one on the red die?

<https://tutorcs.com>

# Exercises

## Exercise

程序代写代做 CS编程辅导

RW: 5.2.12 Two dice, a red die and a black die, are rolled.

What is the probabil

(a) the maximum o  
bers is 4?



WeChat: cstutorcs

(b) their minimum is 4?  
Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Exercise

程序代写代做 CS编程辅导

## Exercises

- RW: 5.2.5 An urn contains 3 red and 4 black balls. 3 balls are removed without replacement. What are the probabilities that
- (a) all 3 are red
  - (b) all 3 are black
  - (c) one is red, two are black



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Outline

程序代写代做 CS编程辅导

Elementary Discrete



Independence

Infinite Sample Spaces

WeChat: cstutorcs

Recursive Probability Computations

Assignment Project Exam Help

Conditional Probability

Email: tutorcs@163.com

Indpendence, revisited

QQ: 749389476

<https://tutorcs.com>

# Infinite sample spaces

程序代写代做 CS编程辅导

Probability distributions generalize to infinite sample spaces **with some provisos.**



- In continuous spaces (e.g.  $\mathbb{R}$ ):
  - Probability distributions are *measures*;
  - Sums are *integrals*;
  - Non-zero probabilities apply to ranges;
  - Probability of a single event is 0. Note: Probability 0 is not the same as impossible.
- In discrete spaces (e.g.  $\mathbb{N}$ ):
  - Probability 0 is the same as impossible.
  - No uniform distribution!
  - Non-uniform distributions exist, e.g.  $P(0) = 1$ ,  $P(n) = 0$  for  $n > 0$ ; or  $P(0) = 0$ ,  $P(n) = \frac{1}{2^n}$  for  $n > 0$ .
  - May consider limiting probabilities if that makes sense.

WeChat: estutores

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Asymptotic Estimate of Relative Probabilities

程序代写代做 CS编程辅导

## Example

Event  $A \stackrel{\text{def}}{=} \text{one die rolled } n \text{ times and you obtain two 6's}$

Event  $B \stackrel{\text{def}}{=} n \text{ dice rolled simultaneously and you obtain one 6}$



$$P(A) = \frac{\binom{n}{2} \cdot 5^{n-2}}{6^n} \quad P(B) = \frac{\binom{n}{1} \cdot 5^{n-1}}{6^n}$$

WeChat: estutorcs  
Assignment Project Exam Help

$$\text{Therefore } \frac{P(A)}{P(B)} = \frac{\binom{n}{2}}{\binom{n}{1}} \cdot \frac{5}{6} = \frac{n(n-1)}{2} \cdot \frac{5}{6} = \frac{n-1}{10} \in \Theta(n)$$

QQ: 749389476

$n$	1	2	3	4	...	11	...	20	...
$P(A)$	0	$\frac{1}{36}$	$\frac{5}{72}$	$\frac{25}{216}$	...	0.296	...	0.198	...
$P(B)$	$\frac{1}{6}$	$\frac{10}{36}$	$\frac{25}{72}$	$\frac{125}{324}$	...	0.296	...	0.104	...

<https://tutorcs.com>

# Outline

程序代写代做 CS编程辅导

Elementary Discrete



Independence

Infinite Sample Spaces

WeChat: cstutorcs

Recursive Probability Computations

Assignment Project Exam Help

Conditional Probability

Email: tutorcs@163.com

Indpendence, revisited

QQ: 749389476

<https://tutorcs.com>

# Use of Recursion in Probability Computations

程序代写代做 CS编程辅导

## Question

Given  $n$  tosses of a coin, what is the probability of two HEADS in a row?



## Answer

WeChat: cstutorcs

Recall  $N(n)$ : the number of sequences without HH.

Assignment Project Exam Help

$N(n) = N(n - 1) + N(n - 2)$ :  $N(n) = \text{FIB}(n + 1)$

Email: tutorcs@163.com

$$N(n) \approx \frac{1}{\sqrt{5}} \left( \frac{\sqrt{5}+1}{2} \right)^{n+1}$$

QQ: 749389476

$$p_n = 1 - \frac{\text{FIB}(n+1)}{2^n} \approx 1 - 0.72 \cdot (0.8)^n$$

# Example

程序代写代做 CS编程辅导

## Question

Given  $n$  tosses, what is the probability  $q_n$  of at least one HHH?

$$q_0 = q_1 = q_2 = 0; q_3 = ?$$



Then recursive computation:

WeChat: cstutorcs

$$\begin{aligned} q_n &= \frac{1}{2} q_{n-1} && \text{(initial: T)} \\ &+ \frac{1}{4} q_{n-2} && \text{(initial: HT)} \\ &+ \frac{1}{8} q_{n-3} && \text{(initial: HHT)} \\ &+ \frac{1}{8} && \text{(start with: HHH)} \end{aligned}$$

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Example

## Question

程序代写代做 CS编程辅导

Two dice are rolled repeatedly. What is the probability that '6–6' will occur before two consecutive (back-to-back) 'totals seven'?




NB

The probability of either occurring at a given roll is the same:  $\frac{1}{36}$ .

WeChat: cstutorcs

Let  $p = P(6-6 \text{ first})$  Assignment Project Exam Help



# Example

## Question

程序代写代做 CS编程辅导

A coin is tossed 'indefinitely'. Which pattern is more likely (and by how much) to appear - H or HHT?



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Difficult probability calculations

程序代写代做 CS编程辅导



NB

*The majority of problems in probability and statistics do not have such elegant solutions. Hence the use of computers for either precise calculations or approximate simulations is mandatory. However, it is the use of recursion that simplifies such computing or, quite often, makes it possible in the first place.*

WeChat: estutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Outline

程序代写代做 CS编程辅导

Elementary Discrete



Independence

Infinite Sample Spaces

WeChat: cstutorcs

Recursive Probability Computations

Assignment Project Exam Help

Conditional Probability

Email: tutorcs@163.com

Indpendence, revisited

QQ: 749389476

<https://tutorcs.com>

# Conditional probability

程序代写代做 CS编程辅导

## Definition

Conditional probability



given  $S$ :

$$P(E|S) = \frac{P(E \cap S)}{P(S)}, \quad E, S \subseteq \Omega$$

WeChat: cstutorcs

It is defined only when  $P(S) \neq 0$

Assignment Project Exam Help

NB

Email: tutorcs@163.com

$P(A|B)$  and  $P(B|A)$  are, in general, not related — one of these values predicts, by itself, essentially nothing about the other.

QQ: 749389476

The only exception, applicable when  $P(A), P(B) \neq 0$ , is that  
 $P(A|B) = 0$  iff  $P(B|A) = 0$  iff  $P(A \cap B) = 0$ .

<https://tutorcs.com>

# Conditional probability in uniform distributions

If  $P$  is the uniform distribution over a finite set  $\Omega$ , then

程序代写代做 CS编程辅导

$$P\left(\frac{\frac{|E \cap S|}{|\Omega|}}{\frac{|S|}{|\Omega|}} = \frac{|E \cap S|}{|S|}\right)$$

This observation can help calculations...



## Example

WeChat: cstutorcs

RW: 9.1.6 A coin is tossed four times. What is the probability of

(a) two consecutive HEADS

(b) two consecutive HEADS given that  $\geq 2$  tosses are HEADS

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Conditional probability in uniform distributions

If  $P$  is the uniform distribution over a finite set  $\Omega$ , then

程序代写代做 CS 编程辅导

$$P\left(\frac{\frac{|E \cap S|}{|\Omega|}}{\frac{|S|}{|\Omega|}} = \frac{|E \cap S|}{|S|}\right)$$

This observation can help calculations...



## Example

WeChat: cstutorcs

RW: 9.1.6 A coin is tossed four times. What is the probability of

(a) two consecutive HEADS

Assignment Project Exam Help

(b) two consecutive HEADS given that  $\geq 2$  tosses are HEADS

Email: tutorcs@163.com

T T T T      H T T T

T T T H      H T T H

T T H T      H T H T

T T H H      H H T H https://tutorcs.com

T H T T      H H T T

T H T H      H H T H

T H H T      H H H T

T H H H      H H H H

QQ: 749389476

# Conditional probability in uniform distributions

If  $P$  is the uniform distribution over a finite set  $\Omega$ , then

程序代写代做 CS 编程辅导

$$P\left(\frac{|\Omega|}{|\Omega|} = \frac{|E \cap S|}{|S|}\right)$$


This observation can help calculations...

## Example

WeChat: cstutorcs

RW: 9.1.6 A coin is tossed four times. What is the probability of

(a) two consecutive HEADS

(b) two consecutive HEADS given that  $\geq 2$  tosses are HEADS

Email: tutorcs@163.com

T T T T

T T T H

T T H T

T T H H

T H T T

T H T H

T H H T

T H H H

QQ: 749389476

<https://tutorcs.com> (a)  $\frac{8}{16}$

# Conditional probability in uniform distributions

If  $P$  is the uniform distribution over a finite set  $\Omega$ , then

程序代写代做 CS 编程辅导

$$P\left(\frac{\frac{|E \cap S|}{|\Omega|}}{\frac{|S|}{|\Omega|}} = \frac{|E \cap S|}{|S|}\right)$$


This observation can help calculations...

## Example

WeChat: cstutorcs

RW: 9.1.6 A coin is tossed four times. What is the probability of

(a) two consecutive HEADS

(b) two consecutive HEADS given that  $\geq 2$  tosses are HEADS

Email: tutorcs@163.com

T T T T      H T T T

T T T H      H T T H

T T H T      H T H T

T T H H      H H T H

T H T T      H H T T

T H T H      H H T H

T H H T      H H H T

T H H H      H H H H

QQ: 749389476

https://tutorcs.com

(a)  $\frac{8}{16}$     (b)  $\frac{8}{11}$

# Some General Rules

程序代写代做 CS编程辅导

## Fact

- $A \subseteq B \rightarrow P(A|B) = P(A)$
- $A \subseteq B \rightarrow P(B|A) = P(B)$
- $P(A \cap B|B) = P(A|B)$
- $P(\emptyset|A) = 0$  for  $A \neq \emptyset$
- $P(A|\Omega) = P(A)$
- $P(A^c|B) = 1 - P(A|B)$



cstutorcs

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutores@163.com](mailto:tutores@163.com)

## NB

QQ: 749389476

- $P(A|B)$  and  $P(A|B^c)$  are not related
- $P(A|B), P(B|A), P(A^c|B^c), P(B^c|A^c)$  are not related

<https://tutorcs.com>

# Example

## Example

程序代写代做 CS编程辅导

Two dice are rolled and the outcomes recorded as  $b$  for the black die,  $r$  for the red die. Define the events  $B = \{b \geq 3\}$ ,  $R = \{r \geq 3\}$ ,  $S = \{s \geq 6\}$ .



$$P(S|B) = \frac{4+5+6+6}{24} = \frac{21}{24} = \frac{7}{8} = 87.5\%$$

WeChat: cstutorcs

Assignment Project Exam Help

$$P(B|S) = \frac{4+5+6+6}{26} = \frac{21}{26} = 80.8\%$$

Email: tutorcs@163.com

The (common) numerator  $4+5+6+6 = 21$  represents the size of the  $B \cap S$  — the common part of  $B$  and  $S$ , that is, the number of rolls where  $b \geq 3$  and  $s \geq 6$ . It is obtained by considering the different cases:  $b = 3$  and  $s \geq 6$ , then  $b = 4$  and  $s \geq 6$  etc.

The denominators are  $|B| = 24$  and  $|S| = 26$

# Example

## Example (cont'd) 程序代写代做 CS编程辅导

Recall:  $B = \{b \geq 3\}$ ,  $R = \{r > 3\}$ ,  $S = \{s \geq 6\}$



$$P(S|B \cup R) = 2/3 = 66.7\%$$

$$P(S) = \frac{5+6+5+4+3+2+1}{36} = \frac{26}{36} = 72.22\%$$

Assignment Project Exam Help

$$P(S|B \cup R) = \frac{2+3+4+5+6+6}{32} = \frac{26}{32} = 81.25\%$$

QQ: 749389476

The set  $B \cup R$  represents the event 'b or r'.

It comprises all the rolls except for those with both the red and the black die coming up either 1 or 2.

$$P(S|B \cap R) = 1 = 100\% \text{ — because } S \supseteq B \cap R$$

# Exercise

程序代写代做 CS编程辅导

## Exercise

RW: 9.1.9 Consider there are eight black marbles; draw two without replacement.



$b_1$  — Black on the first draw,  
 $b_2$  — Black on the second draw  
 $r_1$  — Red on first draw,  
 $r_2$  — Red on second draw

Find the probabilities

WeChat: cstutorcs

(a) both Red:

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

## Exercise

(b) both Black:

程序代写代做 CS编程辅导



(c) one Red, one Bla

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Exercise

程序代写代做 CS编程辅导

## Exercise

RW: 9.1.12 What is the probability of a flush given that all five cards in a Poker hand are red?

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



# Exercise

## Exercise

程序代写代做 CS编程辅导

RW: 9.1.22 Prove the following:

If  $P(A|B) > P(A)$  (" correlation") then  $P(B|A) > P(B)$



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Outline

程序代写代做 CS编程辅导

Elementary Discrete



Independence

Infinite Sample Spaces

WeChat: cstutorcs

Recursive Probability Computations

Assignment Project Exam Help

Conditional Probability

Email: tutorcs@163.com

Indpendence, revisited

QQ: 749389476

<https://tutorcs.com>

# Stochastic Independence, again

## Definition

程序代写代做 CS编程辅导

$A$  and  $B$  are **(stochastic) independent** (notation:  $A \perp B$ ) if  
 $P(A \cap B) = P(A) \cdot P(B)$



If  $P(A) \neq 0$  and  $P(B) \neq 0$ , then all four of the following are *equivalent* definitions:

WeChat: cstutorcs

- $P(A \cap B) = P(A)P(B)$
- $P(A|B) = P(A)$
- $P(B|A) = P(B)$
- $P(A^c|B) = P(A^c)$  or  $P(A|B^c) = P(A)$  or  $P(A^c|B^c) = P(A^c)$

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

The last one claims that

<https://tutorcs.com>

$$A \perp B \leftrightarrow A^c \perp B \leftrightarrow A \perp B^c \leftrightarrow A^c \perp B^c$$

# Using independence to simplify calculations

程序代写代做 CS编程辅导



Independence of events, just pairwise independence, can greatly simplify computations and reasoning in AI applications. It is common for many expert systems to make an approximating assumption of independence, even if it is not completely satisfied.

WeChat: cstutorcs  
Assignment Project Exam Help



Email: [tutorcs@163.com](mailto:tutorcs@163.com) =  $P(\text{sense}_t | \text{loc}_t, \text{sense}_{t-1}, \text{loc}_{t-1}, \dots)$

QQ: 749389476

<https://tutorcs.com>

# Exercise

## Exercise

程序代写代做 CS编程辅导

RW: 9.1.7 Suppose the experiment leads to events  $A$ ,  $B$  and  $C$  with  $P(A) = 0.3$ ,  $P(B) = 0.4$  and  $P(A \cap B) = 0.1$

(a)  $P(A|B) = \frac{P(A \cap B)}{P(B)}$

(b)  $P(A^c) = 1 - P(A) = 0.7$

(c) Is  $A \perp B$ ? No.  $P(A) \cdot P(B) = 0.12 \neq P(A \cap B)$

(d) Is  $A^c \perp B$ ? No, as can be seen from (c).

QQ: 749389476

Note:  $P(A^c \cap B) = P(B) - P(A \cap B) = 0.4 - 0.1 = 0.3$

$$P(A^c) \cdot P(B) = 0.7 \cdot 0.4 = 0.28$$

# Exercise

程序代写代做 CS编程辅导



## Exercise

RW: 9.1.8 Given  $A \perp B$ ,  $P(A) = 0.4$ ,  $P(B) = 0.6$

$$P(A|B) =$$

WeChat: cstutorcs

$$P(A \cup B) =$$

Assignment Project Exam Help

$$P(A^c \cap B) =$$

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Supplementary Exercise

## Exercise

程序代写代做 CS编程辅导

RW: 9.5.5 (Supp) W  two events with

$$P(A) = \frac{1}{4}, P(B) =$$

True, false or could be



a  $P(A \cap B) = \frac{1}{12}$

b  $P(A \cup B) = \frac{7}{12}$

c  $P(B|A) = \frac{P(B)}{P(A)}$

Assignment Project Exam Help

Email: tutorcs@163.com

d  $P(A|B) \geq P(A)$

QQ: 749389476

e  $P(A^c) = \frac{3}{4}$

<https://tutorcs.com>

f  $P(A) = P(B)P(A|B) + P(B^c)P(A|B^c)$