

程序代写代做 CS编程辅导



COM

Foundations of Computer Science

WeChat: cstutorcs

Lecture 16: Statistics

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



UNSW
SYDNEY

Outline

程序代写代做 CS编程辅导



Random Variables and Expectation

Linearity of Expectation

Expected Time to Success

Standard Deviation and Variance

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Outline

程序代写代做 CS编程辅导



Random Variables and Expectation

Linearity of Expectation

Expected Time to Success

Standard Deviation and Variance

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Random Variables

程序代写代做 CS编程辅导

Definition

An (integer) **random variable** is a function from Ω to \mathbb{Z} .
In other words, it assigns a number value with every outcome.



Random variables are often denoted by X, Y, Z, \dots

We extend arithmetic to random variables in the natural way.

WeChat: cstutorcs

Definition

Given random variables X, Y and integer k :

Email: tutorcs@163.com

$$X + Y : \quad \omega \mapsto X(\omega) + Y(\omega)$$

$$XY : \quad \omega \mapsto X(\omega) \cdot Y(\omega)$$

$$X - k : \quad \omega \mapsto X(\omega) - k$$

$$kX : \quad \omega \mapsto k \cdot X(\omega)$$

<https://tutorcs.com>

Examples

Example

程序代写代做 CS编程辅导

Random variable X : value of rolling one die

$$\Omega = \{1, 2, 3, 4, 5, 6\}$$

$$X(i) = i$$



Example

WeChat: cstutorcs

Random variable X_s : sum of rolling two dice

Assignment Project Exam Help

$$\Omega = \{(1, 1), (1, 2), \dots, (6, 6)\}$$

$$X_s((1, 1)) = 2 \quad X_s((1, 2)) = 3 = X_s((2, 1)) \dots$$

Email: tutorcs@163.com

QQ: 749389476

Question

Is $X_s = X + X$? No. <https://tutorcs.com>

$X_s = X + Y$ where X and Y are independent and identically distributed (i.i.d)

Expectation

程序代写代做 CS编程辅导

Definition

The **expected value** (also called “expectation” or “average”) of a random variable X



$$E(X) = \sum_{k \in \mathbb{Z}} P(X = k) \cdot k$$

WeChat: [tutorcs](#)

Assignment Project Exam Help

NB

Email: tutorcs@163.com

Expectation is a truly universal concept; it is the basis of all decision making, of estimating gains and losses, in all actions under risk. Historically, a rudimentary concept of expected value arose long before the notion of probability.

QQ: [749389476](#)

<https://tutorcs.com>

Examples

程序代写代做 CS编程辅导

Example

The expected value when rolling one die is:



$$E(X) = \frac{1}{6} \cdot 1 + \frac{1}{6} \cdot 2 + \dots + \frac{1}{6} \cdot 6 = 3.5$$

WeChat: [cstutorcs](#)

Assignment Project Exam Help

Example

The expected sum when rolling two dice is

Email: tutorcs@163.com

QQ: [749389476](#)

$$E(X_s) = \frac{1}{36} \cdot 2 + \frac{2}{36} \cdot 3 + \dots + \frac{6}{36} \cdot 7 + \dots + \frac{1}{36} \cdot 12 = 7$$

<https://tutorcs.com>

Examples

程序代写代做 CS编程辅导



Example

RW: 9.3.3 Buy one lottery ticket for \$1. The only prize is \$1M. Each ticket has probability $6 \cdot 10^{-7}$ of winning.

$$\Omega = \{win, lose\} \quad X_L(win) = \$999,999 \quad X_L(lose) = -\$1$$
$$E(X_L) = 6 \cdot 10^{-7} \cdot \$999,999 + (1 - 6 \cdot 10^{-7}) \cdot -\$1 = -\$0.4$$

QQ: 749389476

<https://tutorcs.com>

Outline

程序代写代做 CS编程辅导



Random Variables and Expectation

Linearity of Expectation

Expected Time to Success

Standard Deviation and Variance

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Linearity of expectation

程序代写代做 CS编程辅导

Theorem (linearity of expected value)

For any random variables X and Y and integer k :

$$E(X + Y) = E(X) + E(Y) \quad E(k \cdot X) = k \cdot E(X)$$

WeChat: cstutorcs

Assignment Project Exam Help

Example

The expected sum when rolling two dice can be computed as

$$E(X_s) = E(X) + E(Y) = 3.5 + 3.5 = 7$$

<https://tutorcs.com>



Example

程序代写代做 CS编程辅导

Example

$E(S_n)$, where $S_n \stackrel{\text{def}}{=} |\text{HEADS in } n \text{ tosses}|$

- 'hard way'

$$E(S_n) = \sum_{k=0}^n \binom{n}{k} \cdot k = \sum_{k=0}^n \frac{1}{2^n} \binom{n}{k} \cdot k$$

since there are $\binom{n}{k}$ sequences of n tosses with k HEADS, and each sequence has the probability $\frac{1}{2^n}$

$$= \frac{1}{2^n} \sum_{k=1}^n \frac{n}{k} \binom{n-1}{k-1} k = \frac{n}{2^n} \sum_{k=0}^{n-1} \binom{n-1}{k} = \frac{n}{2^n} \cdot 2^{n-1} = \frac{n}{2}$$

Email: tutores@163.com

using the 'binomial identity' $\sum_{k=0}^n \binom{n}{k} = 2^n$

QQ: 749389476

- 'easy way'

$$E(S_n) = E(S_1^1 + \dots + S_1^n) = \sum_{i=1}^n E(S_1^i) = nE(S_1) = n \cdot \frac{1}{2}$$

Note: $S_n \stackrel{\text{def}}{=} |\text{HEADS in } n \text{ tosses}|$ while each $S_1^i \stackrel{\text{def}}{=} |\text{HEADS in 1 toss}|$



WeChat: tutores

Assignment Project Exam Help

Email: tutores@163.com

QQ: 749389476

https://tutores.com

Observations

程序代写代做 CS编程辅导



Fact

If X_1, X_2, \dots, X_n are independent, identically distributed random variables, then $E(X_1 + X_2 + \dots + X_n) = E(nX_1) = nE(X_1)$.

Assignment Project Exam Help

NB

$X_1 + X_2 + \dots + X_n$ and X_1 are very different random variables.

QQ: 749389476

<https://tutorcs.com>

Exercises

程序代写代做 CS编程辅导

Exercise



You face a quiz consisting of n true/false questions, and your plan is to guess the answer to each question (randomly, with probability 0.5 of being correct). There are no negative marks, and answering four or more questions correctly suffices to pass. What is the probability of passing and what is the expected score?

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Exercises

Exercise

程序代写代做 CS编程辅导

RW: 9.3.7



An urn has $m + n = 7$ marbles, $m \geq 0$ red and $n \geq 0$ blue. 7 marbles selected at random without replacement. What is the expected number of red marbles drawn?

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Outline

程序代写代做 CS编程辅导



Random Variables and Probability Distribution

Linearity of Expectation

Expected Time to Success

Standard Deviation and Variance

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example

Example

程序代写代做 CS编程辅导

Find the average waiting time for the first HEAD, with no upper bound on the 'duration' (allows for all possible sequences of tosses, regardless of how many times TAILS occur initially).



$$A = E(X_w) = \sum_{k=1}^{\infty} k \cdot P(X_w = k) = \sum_{k=1}^{\infty} k \frac{1}{2^k}$$

WeChat: tutores

Assignment Project Exam Help

This can be evaluated by breaking the sum into a sequence of geometric progressions

Email: tutores@163.com

QQ: 749389476

$$\frac{1}{2} + \frac{2}{2^2} + \frac{3}{2^3} + \dots$$

$$= \left(\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots \right) + \left(\frac{1}{2^2} + \frac{1}{2^3} + \dots \right) + \left(\frac{1}{2^3} + \dots \right) + \dots$$

$$= 1 + \frac{1}{2} + \frac{1}{2^2} + \dots$$

$$= 2$$

<https://tutores.com>

Expected time to success

There is also a recursive 'trick' for solving the sum

程序代写代做 CS编程辅导

$$A = \sum_{k=1}^{\infty} \frac{k}{2^k} = \sum_{k=1}^{\infty} \frac{k-1}{2^k} + \sum_{k=1}^{\infty} \frac{1}{2^k} = \frac{1}{2} \sum_{k=1}^{\infty} \frac{k-1}{2^{k-1}} + 1 = \frac{1}{2}A + 1$$

Now $A = \frac{A}{2} + 1$ all



NB

WeChat: cstutorcs

A much simpler but equally valid argument is that you expect 'half' a HEAD in 1 toss, so you ought to get a 'whole' HEAD in 2 tosses.

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

Theorem

If the probability of success is p then:

<https://tutorcs.com>

- The expected number of (indep.) trials before 1 success is $\frac{1}{p}$
- The expected number of (indep.) trials before k successes is $\frac{k}{p}$

Exercise

程序代写代做 CS编程辅导



Exercise

RW: 9.4.12 A die is rolled until the first 4 appears. What is the expected waiting time?

WeChat: tutormcs

Assignment Project Exam Help

Email: tutormcs@163.com

QQ: 749389476

<https://tutormcs.com>

Example

程序代写代做 CS编程辅导

To find an object \mathcal{X} in a sorted list L of elements, one needs to search linearly through the list. The probability of $\mathcal{X} \in L$ be p , hence there is $1 - p$ likelihood of \mathcal{X} being absent altogether. Find the expected number of operations.



If the element is in the list, then the number of comparisons averages to $\frac{1}{n}(1 + \dots + n)$; if absent we need n comparisons.

The first case has probability p , the second $1 - p$. Combining these we find

$$E_n = p \frac{1 + \dots + n}{n} + (1 - p)n = p \frac{n + 1}{2} + (1 - p)n = (1 - \frac{p}{2})n + \frac{p}{2}$$

As one would expect, increasing p leads to a lower E_n .

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

https://tutorcs.com

Success vs Expected value

程序代写代做 CS编程辅导



Question

Does high probability of success lead to a high expected value?

Generally, no.

WeChat: cstutorcs

Example

Assignment Project Exam Help

Buying more tickets in the lottery increases your chances of winning, but the expected value of winnings decreases.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example

程序代写代做 CS编程辅导



Example

Roulette (outcomes $0, 1, \dots, 36$). Win: $35 \times \text{bet}$

WeChat: cstutorcs

Strategy 1: Bet \$1 on a single number

Assignment Project Exam Help

- Probability of winning: $\frac{1}{37}$

- Expected winnings: $\frac{1}{37}(\$35) + \frac{36}{37}(-\$1) \approx -2.7c$

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example

Example

程序代写代做 CS编程辅导

Roulette (outcomes (0-36)). Win: $35 \times \text{bet}$

Strategy 2: Place \$1 on 24 numbers, selected from among 0 to 36.



- Probability of winning: $\frac{24}{37} \approx 65\%$
- Expected winnings:
 - If one of the numbers comes up, win \$35 from the bet on that number and lose \$23 from the bets on the remaining numbers, thus collecting \$12.
This happens with probability $p = \frac{24}{37}$.
 - With probability $q = \frac{13}{37}$ none of the numbers appear, leading to loss of \$24.

So expected winnings are:

$$p \cdot \$12 - q \cdot \$24 = \$12 \frac{24}{37} - \$24 \frac{13}{37} = -\$ \frac{24}{37} \approx -65c = 24 \times -2.7c$$

Gambler's ruin

程序代写代做 CS编程辅导

Many so-called 'winning systems' that purport to offer a winning strategy do something that they provide a scheme for frequent relatively moderate wins at the cost of an occasional very big loss.



It turns out (it is a formal theorem) that there can be *no system* that converts an 'unfair' game into a 'fair' one. In the language of decision theory, 'unfair' denotes a game whose individual bets have negative expectation.

It can be easily checked that any individual bets on roulette, on lottery tickets or on just about any commercially offered game have negative expected value.

WeChat: tutorms

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Outline

程序代写代做 CS编程辅导



Random Variables and Expectation

Linearity of Expectation

Expected Time to Success

Standard Deviation and Variance

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Standard Deviation and Variance

Definition

程序代写代做 CS编程辅导

For random variable X , the **standard deviation** is the expected value (or: **mean**) $\mu = E(X)$,



$$\sqrt{E((X - \mu)^2)}$$

and the **variance** of X is

WeChat: cstutorcs

Assignment σ^2 Project Exam Help

Standard deviation and variance measure how spread out the values of a random variable are. The smaller σ^2 the more confident we can be that $X(\omega)$ is close to $E(X)$, for a randomly selected ω .

Email: tutores@163.com

QQ: 749389476

<https://tutorcs.com>

NB

The variance can be calculated as $E((X - \mu)^2) = E(X^2) - \mu^2$

Example

程序代写代做 CS编程辅导

Example

Random variable X_d of a rolled die



WeChat: [tutorcs](https://tutorcs.com)

Assignment Project Exam Help

Email: tutorcs@163.com

$$E(X_d^2) = \frac{1}{6} \cdot 1 + \frac{1}{6} \cdot 4 + \frac{1}{6} \cdot 9 + \frac{1}{6} \cdot 16 + \frac{1}{6} \cdot 25 + \frac{1}{6} \cdot 36 = \frac{91}{6}$$

Hence, $\sigma^2 = E(X_d^2) - \mu^2 = \frac{35}{12} \rightarrow \sigma \approx 1.71$

<https://tutorcs.com>

Exercises

程序代写代做 CS编程辅导

Exercises



RW: 9.5.10 (Supp) Two independent experiments are performed.
 $P(\text{1st experiment succeeds}) = 0.7$
 $P(\text{2nd experiment succeeds}) = 0.2$
Random variable X counts the number of successful experiments.

- Assignment Project Exam Help**
Email: tutorcs@163.com
QQ: 749389476
- a Expected value of X ?
 - b Probability of exactly one success?
 - c Probability of at most one success?
 - e Variance of X ?
- <https://tutorcs.com>**