

程序代写代做 CS编程辅导



COM

Foundations of Computer Science

WeChat: cstutorcs

Lecture 6: Equivalence Relations and Partial Orders

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



UNSW  
SYDNEY

# Outline

程序代写代做 CS编程辅导

Equivalence Relation

Partial Orders

Feedback



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Outline

程序代写代做 CS编程辅导

Equivalence Relation

Partial Orders

Feedback



WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

# Equivalence relations

程序代写代做 CS编程辅导

Equivalence relations are a general notion of “equality”. They are relations which a



- Reflexive (R): Every element should be “equal” to itself
- Symmetric (S): If  $x$  is “equal” to  $y$ , then  $y$  should be “equal” to  $x$
- Transitive (T): If  $x$  is “equal” to  $y$  and  $y$  is “equal” to  $z$ , then  $x$  should be “equal” to  $z$ .

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

## Definition

A binary relation  $R \subseteq S \times S$  is *equivalence relation* if it satisfies (R), (S), (T).

QQ: 749389476

<https://tutorcs.com>

### Example

程序代写代做 CS编程辅导

Partition of  $\mathbb{Z}$  into classes of numbers with the same remainder on division by  $p$ ; it is particularly important for  $p$  prime

$$\mathbb{Z}(p) = \{0, 1, \dots, p-1\}$$

One can define all four arithmetic operations (with the usual properties) on  $\mathbb{Z}_p$  for a prime  $p$ ; division has to be restricted when  $p$  is not prime.

WeChat: cstutorcs

Assignment Project Exam Help

### NB

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

$(\mathbb{Z}_p, +, \cdot, 0, 1)$  are fundamental algebraic structures known as **rings**. These structures are very important in coding theory and cryptography.

QQ: 749389476

<https://tutorcs.com>

# Equivalence Classes and Partitions

程序代写代做 CS编程辅导



Suppose  $R \subseteq S \times S$  is an equivalence relation  
The **equivalence class** (w.r.t.  $R$ ) of an element  $s \in S$  is

$[s] = \{t : t \in S \text{ and } sRt\}$

Assignment Project Exam Help

**Fact**

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

$s R t$  if and only if  $[s] = [t]$ .

QQ: 749389476

<https://tutorcs.com>

# Equivalence classes: Proof example

程序代写代做 CS编程辅导



## Proof

Suppose  $[s] = [t]$ . Recall that  $[s] = \{x \in S : (s, x) \in R\}$ . We will show that  $(s, t) \in R$ .

WeChat: cstutorcs

Because  $R$  is reflexive,  $(t, t) \in R$ .

Assignment Project Exam Help

Therefore  $t \in [t]$ .

Email: tutorms@163.com

Because  $[t] = [s]$ , it follows that  $t \in [s]$ .

But then  $(s, t) \in R$  by the definition of  $[s]$ .

QQ: 749389476

<https://tutorcs.com>

# Equivalence classes: Proof example

程序代写代做 CS编程辅导

## Proof



Now suppose  $(s, t) \in R$ . We will show  $[s] = [t]$  by showing  $[s] \subseteq [t]$  and  $[t] \subseteq [s]$ .

Take any  $x \in [s]$ .

By the definition of  $[s]$ ,  $(s, x) \in R$ .

Since  $R$  is symmetric  $(x, s) \in R$ .

Since  $R$  is transitive and  $(s, t) \in R$  we have that  $(x, t) \in R$ .

Since  $R$  is symmetric  $(t, x) \in R$ .

Therefore,  $x \in [t]$ .

Therefore  $[s] \subseteq [t]$ .

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutors@163.com

QQ: 749389476

<https://tutorcs.com>



# Equivalence classes: Proof example

程序代写代做 CS编程辅导

## Proof



Now suppose  $(s, t) \in R$ . We will show  $[s] = [t]$  by showing  $[s] \subseteq [t]$  and  $[t] \subseteq [s]$ .

Take any  $x \in [t]$ . WeChat: cstutorcs

By the definition of  $[t]$ ,  $(t, x) \in R$ . Assignment Project Exam Help

Since  $R$  is transitive and  $(s, t) \in R$  we have that  $(s, x) \in R$ . Email: tutores@163.com

Therefore  $x \in [s]$ .

QQ: 749389476

Therefore  $[t] \subseteq [s]$ .

<https://tutores.com>



# Partitions

程序代写代做 CS编程辅导

## Definition

A **partition** of a set  $S$  is a collection of sets  $S_1, \dots, S_k$  such that

- $S_i$  and  $S_j$  are disjoint ( $i \neq j$ )
- $S = S_1 \cup S_2 \cup \dots \cup S_k = \bigcup_{i=1}^k S_i$



WeChat: cstutorcs

The collection of all equivalence classes  $\{[s] : s \in S\}$  forms a partition of  $S$ .

Assignment Project Exam Help

In the opposite direction, a partition of a set defines the

equivalence relation on that set. If  $S = S_1 \cup \dots \cup S_k$ , then we can

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

define  $\sim \subseteq S \times S$  as:

QQ: 749389476

$s \sim t$  exactly when  $s$  and  $t$  belong to the same  $S_i$ .

<https://tutorcs.com>

# Exercises

程序代写代做 CS编程辅导

## Exercises

RW: 3.6.6 (supp)

- (d) Show that  $m^2 =_{(5)} n^2$  is an equivalence on  $S = \{1, \dots, 7\}$ .



WeChat: cstutorcs

Assignment Project Exam Help

Find all the equivalence classes.

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

# Exercises

程序代写代做 CS编程辅导

## Exercises

RW: 3.6.6 (supp)



- (d) Show that  $m \equiv_{(5)} n^2$  is an equivalence on  $S = \{1, \dots, 7\}$ .

It just means that  $m \equiv_{(5)} n$  or  $m \equiv_{(5)} -n$ ,  
e.g.  $1 \equiv_{(5)} -4$ .

This satisfies (R), (S), (T).

Find all the equivalence classes.

We have

$$\begin{aligned}[1] &= \{1, 4, 6\} \\ [2] &= \{2, 3, 7\} \\ [5] &= \{5\}\end{aligned}$$

QQ: 749389476

<https://tutorcs.com>

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

Assignment Project Exam Help

WeChat: [tutorcs](https://tutorcs.com)

# Outline

程序代写代做 CS编程辅导



Equivalence Relation

Partial Orders

Feedback

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Partial Order

程序代写代做 CS编程辅导

A **partial order**  $\preceq$  on  $S$  satisfies (R), (AS), (T).

We call  $(S, \preceq)$  a **poset** or **partially ordered set**



## Examples

Posets:

- $(\mathbb{Z}, \leq)$

- $(\text{Pow}(X), \subseteq)$  for some set  $X$

- $(\mathbb{N}, |)$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

Not posets:

- $(\mathbb{Z}, <)$

- $(\mathbb{Z}, |)$

QQ: 749389476

<https://tutorcs.com>

# Hasse diagram

Every finite poset  $(S, \preceq)$  can be represented with a **Hasse diagram**:

- Nodes are elements of  $S$
- An edge is drawn from  $x$  to  $y$  if  $x \prec y$  and there is no  $z$  such that  $x \prec z \prec y$

WeChat: cstutorcs


## Example

Hasse diagram for positive divisors of 24 ordered by  $\mid$ :

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



# Ordering Concepts

## Definition

程序代写代做 CS编程辅导

Let  $(S, \preceq)$  be a poset



- **Minimal** element:  $x$  such that there is no  $y$  with  $y \preceq x$
- **Maximal** element:  $x$  such that there is no  $y$  with  $x \preceq y$
- **Minimum (least)** element:  $x$  such that  $x \preceq y$  for all  $y \in S$
- **Maximum (greatest)** element:  $x$  such that  $y \preceq x$  for all  $y \in S$

WeChat: cstutorcs

Assignment Project Exam Help

## NB

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

- *There may be multiple minimal/maximal elements.*
- *Minimum/maximum elements are the unique minimal/maximal elements if they exist.*
- *Minimal/maximal elements always exist in finite posets, but not necessarily in infinite posets.*

QQ: 749389476

<https://tutorcs.com>



# Examples

程序代写代做 CS编程辅导



## Examples

- $\text{Pow}(\{a, b, c\})$  with the order  $\subseteq$   
 $\emptyset$  is minimum;  $\{a, b, c\}$  is maximum
- $\text{Pow}(\{a, b, c\}) \setminus \{\{a, b, c\}\}$  (Proper subsets of  $\{a, b, c\}$ )  
Each two-element subset  $\{a, b\}, \{a, c\}, \{b, c\}$  is maximal.
  - But there is no maximum

QQ: 749389476

<https://tutorcs.com>

# Ordering Concepts

程序代写代做 CS编程辅导

## Definition

Let  $(S, \preceq)$  be a poset



- $x$  is an **upper bound** for  $A$  if  $a \preceq x$  for all  $a \in A$

- $x$  is a **lower bound** for  $A$  if  $x \preceq a$  for all  $a \in A$

- The **set of upper bounds** for  $A$  is defined as

$$\text{ub}(A) = \{x : a \preceq x \text{ for all } a \in A\}$$

- The **set of lower bounds** for  $A$  is defined as

$$\text{lb}(A) = \{x : x \preceq a \text{ for all } a \in A\}$$

- The **least upper bound** of  $A$ ,  $\text{lub}(A)$ , is the minimum of  $\text{ub}(A)$  (if it exists)

- The **greatest lower bound** of  $A$ ,  $\text{glb}(A)$  is the maximum of  $\text{lb}(A)$  (if it exists)

WeChat: tutores

Assignment Project Exam Help

Email: tutores@163.com

QQ: 749389476

<https://tutores.com>

# glb and lub

程序代写代做 CS编程辅导

To show  $x$  is  $\text{glb}(A)$  to show:

- $x$  is a lower bound for all  $a \in A$ .
- $x$  is the greatest of all lower bounds: If  $y \preceq a$  for all  $a \in A$  then  $y \preceq x$ .

WeChat: cstutorcs

## Example

Assignment Project Exam Help

$\text{Pow}(X)$  ordered by  $\subseteq$

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

- $\text{glb}(A, B) = A \cap B$

- $\text{lub}(A, B) = A \cup B$

QQ: 749389476

<https://tutorcs.com>

# Ordering Concepts

程序代写代做 CS编程辅导



## Definition

Let  $(S, \preceq)$  be a poset

- $(S, \preceq)$  is a **lattice** if  $\text{lub}(x, y)$  and  $\text{glb}(x, y)$  exist for every pair of elements  $x, y \in S$ .
- $(S, \preceq)$  is a **complete lattice** if  $\text{lub}(A)$  and  $\text{glb}(A)$  exist for every subset  $A \subseteq S$ .

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

## NB

*A finite lattice is always a complete lattice.*

<https://tutorcs.com>

# Examples

程序代写代做 CS编程辅导

## Examples



- $\{1, 2, 3, 4, 6, 8, 12\}$  partially ordered by divisibility is a lattice
  - e.g.  $\text{lub}(\{4, 6\}) = 12$ ;  $\text{glb}(\{4, 6\}) = 2$
- $\{1, 2, 3\}$  partially ordered by divisibility is not a lattice
  - $\{2, 3\}$  has no lub
- $\{2, 3, 6\}$  partially ordered by divisibility
  - $\{2, 3\}$  has no glb
- $\{1, 2, 3, 12, 18, 36\}$  partially ordered by divisibility
  - $\{2, 3\}$  has no lub (12, 18 are minimal upper bounds)

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749589476

<https://tutorcs.com>

## NB

*An infinite lattice need not have a lub (or no glb) for an arbitrary infinite subset of its elements. In particular no such bound may exist for **all** its elements.*



## Examples

- $(\mathbb{Z}, \leq)$ : neither  $\text{lub}(\mathbb{Z})$  nor  $\text{glb}(\mathbb{Z})$  exist
- $(\mathcal{F}(\mathbb{N}), \subseteq)$  [all finite subsets of  $\mathbb{N}$ ]:  $\text{lub}$  exists for pairs of elements but not generally for (infinite) sets of elements.  $\text{glb}$  exists for any set of elements: intersection of a set of finite sets is finite.
- $(\mathcal{I}(\mathbb{N}), \subseteq)$  [all infinite subsets of  $\mathbb{N}$ ]:  $\text{glb}$  does not exist for some pairs of elements (e.g. odds and evens).  $\text{lub}$  exists for any set of elements: union of a set of infinite sets is always infinite.

程序代写代做CS编程辅导

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Exercises

程序代写代做 CS编程辅导

## Exercises



RW: 11.1.5 Consider  $\mathbb{R} \leq$

- (a) Is this a lattice?
- (b) Give an example of a non-empty subset of  $\mathbb{R}$  that has no upper bound.
- (c) Find  $\text{lub}(\{x \in \mathbb{R} : x \leq 73\})$
- (d) Find  $\text{lub}(\{x \in \mathbb{R} : x \leq 73\})$
- (e) Find  $\text{lub}(\{x : x^2 \leq 73\})$
- (f) Find  $\text{glb}(\{x : x^2 \leq 73\})$

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

# Exercises

程序代写代做 CS编程辅导

## Exercises



RW: 11.1.5 Consider  $\leq$

- (a) Is this a lattice? Yes
- (b) Give an example of a non-empty subset of  $\mathbb{R}$  that has no upper bound.  $\{ r \in \mathbb{R} : r > 0 \}$   
 $= (0, \infty)$
- (c) Find  $\text{lub}(\{ x \in \mathbb{R} : x^2 < 73 \})$  73
- (d) Find  $\text{lub}(\{ x \in \mathbb{R} : x^2 \leq 73 \})$  73
- (e) Find  $\text{lub}(\{ x : x^2 < 73 \})$   $\sqrt{73}$
- (f) Find  $\text{glb}(\{ x : x^2 < 73 \})$   $-\sqrt{73}$

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>



# Total orders

程序代写代做 CS编程辅导

## Definition

A **total order** is a partial order that also satisfies:

(L) *Linearity* (any two elements are comparable):

For all  $x, y$  either:  $x \leq y$  or  $y \leq x$  (or both if  $x = y$ )

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

## NB

On a finite set all total orders are "isomorphic"  
On an infinite set there is quite a variety of possibilities.

QQ: 749389476

<https://tutorcs.com>



# Examples

程序代写代做 CS编程辅导



## Examples

- $\mathbb{Z}$  with  $\leq$ :  
(no minimum/maximum element)
- $\mathbb{Z}$  with  
 $\{(x, y) : (xy \leq 0 \text{ and } x \leq y) \text{ or } (xy > 0 \text{ and } |x| \leq |y|)\}$ :  
(no maximum element, minimum element is -1)
- $\mathbb{Z}$  with  $\{(x, y) : (xy \leq 0 \text{ and } x \geq y) \text{ or } (xy > 0 \text{ and } x \leq y)\}$ :  
(minimum element 1, maximum element -1)

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

# Ordering of a Poset — Topological Sort

## Definition

程序代写代做 CS编程辅导

For a poset  $(S, \preceq)$  any total order  $\leq$  that is consistent with  $\preceq$  (if  $a \preceq b$  then  $a \leq b$ ) is a **topological sort**.



## Example

Consider

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

The following all are topological sorts:

$a \leq b \leq e \leq c \leq f \leq d$

$a \leq e \leq b \leq f \leq c \leq d$

$a \leq e \leq f \leq b \leq c \leq d$

<https://tutorcs.com>

# Well-Ordered Sets

## Definition

A **well-ordered set** is a poset where every subset has a least element.



程序代写代做 CS编程辅导

## NB

The greatest element is not required.

## Examples

- $\mathbb{N} = \{0, 1, \dots\}$
- Disjoint union of copies of  $\mathbb{N}$ :

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

where each  $\mathbb{N}_i \cap \mathbb{N}_j = \emptyset$  and  $\mathbb{N}_0 \subset \mathbb{N}_1 \subset \mathbb{N}_2 \subset \mathbb{N}_3 \dots$

<https://tutorcs.com>

WeChat: [cstutorcs](https://tutorcs.com)

Assignment Project Exam Help

## NB

*Well-ordered sets are an important mathematical tool to prove termination of programs.*

# Orders for Cartesian products and languages

There are several practical ways of combining orders:

- **Product order:** Given posets  $(S, \preceq_S)$  and  $(T, \preceq_T)$ , define:

$$(s, t) \preceq_{S \times T} (s', t') \text{ if } s \preceq_S s' \text{ and } t \preceq_T t'$$

- **Lexicographic order:** Given posets  $(S, \preceq_S)$  and  $(T, \preceq_T)$ , define:

$$(s, t) \leq_{\text{lex}} (s', t') \text{ if } s \prec_S s' \text{ or } (s = s' \text{ and } t \preceq_T t')$$

Extension to words:  $\lambda \leq_{\text{lex}} w$  for all words

- **Lenlex order:** Lexicographic ordering, but order by length first.

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

## Notes

QQ: 749389476

<https://tutorcs.com>

- No implicit weighting.
- No bias toward any component.
- In general, it is only a partial order, even if combining total orders.
- No implicit weighting.

# Example

程序代写代做 CS编程辅导

## Example



RW: 11.2.5 Let  $\mathbb{B} = \{0, 1\}$  with the usual order  $0 < 1$ . List the elements  $101, 010, 110, 0010, 1000$  of  $\mathbb{B}^*$  in the

(a) Lexicographic order

WeChat: cstutorcs

(b) Lenlex order

Assignment Project Exam Help

Email: tutorcs@163.com

RW: 11.2.8 When are the lexicographic order and *lenlex* on  $\Sigma^*$  the same?

QQ: 749389476

<https://tutorcs.com>

# Example

程序代写代做 CS编程辅导

## Example



[RW: 11.2.5] Let  $\mathbb{B} = \{0, 1\}$  with the usual order  $0 < 1$ . List the elements  $101, 010, 110, 0010, 1000$  of  $\mathbb{B}^*$  in the

(a) Lexicographic order

000, 0010, 010, 10, 1000, 101, 11

(b) Lenlex order

10, 11, 000, 010, 101, 0010, 1000

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

[RW: 11.2.8] When are the lexicographic order and *lenlex* on  $\Sigma^*$  the same?

Only when  $|\Sigma| = 1$ . <https://tutorcs.com>

QQ: 749389476

# Exercises

程序代写代做 CS编程辅导

## Exercises

RW: 11.6.6 True or false



- (a) If a set  $\Sigma$  is totally ordered, then the corresponding lexicographic partial order on  $\Sigma^*$  also must be totally ordered.
- (b) If a set  $\Sigma$  is totally ordered, then the corresponding lenlex order on  $\Sigma^*$  also must be totally ordered.
- (c) Every finite poset has a Hasse diagram.
- (d) Every finite poset has a topological sorting.
- (e) Every finite poset has a minimum element.
- (f) Every finite totally ordered set has a maximum element.
- (g) An infinite poset cannot have a maximum element.

WeChat: tutorms

Assignment Project Exam Help

Email: tutorms@163.com

QQ: 749389476

https://tutorms.com



# Exercises

程序代写代做 CS编程辅导

## Exercises

RW: 11.6.6 True or false



- (a) If a set  $\Sigma$  is totally ordered, then the corresponding lexicographic partial order on  $\Sigma^*$  also must be totally ordered. True
- (b) If a set  $\Sigma$  is totally ordered, then the corresponding lexicographic partial order on  $\Sigma^*$  also must be totally ordered. True
- (c) Every finite poset has a Hasse diagram. True
- (d) Every finite poset has a topological sorting. True
- (e) Every finite poset has a minimum element. False
- (f) Every finite totally ordered set has a maximum element. True
- (g) An infinite poset cannot have a maximum element. False

WeChat: tutorms

Assignment Project Exam Help

Email: tutorms@163.com

QQ: 749389476

https://tutorms.com

# Outline

程序代写代做 CS编程辅导



Equivalence Relation

Partial Orders

Feedback

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Weekly Feedback

程序代写代做 CS编程辅导

I would appreciate any comments/suggestions/requests you have on this week's lecture



WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)



QQ: 749389476  
<https://forms.office.com/r/xKKrxYMRn9>

<https://tutorcs.com>