

程序代写代做 CS编程辅导



COM111-1

Foundations of Computer Science

WeChat: cstutorcs

Lecture 10: Induction

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



UNSW
SYDNEY

Outline

程序代写代做 CS编程辅导

Motivation

Basic Induction



Variations on Basic Induction

WeChat: cstutorcs

Structural Induction

Assignment Project Exam Help

Feedback

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Outline

程序代写代做 CS编程辅导

Motivation

Basic Induction

Variations on Basic Induction

Structural Induction

Feedback



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

Recursive datatype

Describe finitely large objects in a finite way



Recursive functions

Define behaviour for these objects in a finite way

Induction

Reason about these objects in a finite way

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example

Recall the recursive program:

程序代写代做 CS编程辅导

Example

Summing the first n numbers:



```
if( $n = 0$ ): 0
```

Website: tutorcs.com

Assignment Project Exam Help

Another attempt:

Email: tutorcs@163.com

Example

QQ: 749389476

```
sum2( $n$ ):
```

<http://tutorcs.com>

```
return  $n * (n + 1) / 2$ 
```

Induction proof **guarantees** that these programs will behave the same.

Inductive Reasoning

程序代写代做 CS编程辅导

Suppose we would like to reach a conclusion of the form

$P(x)$ (of some type)

Inductive reasoning (good in philosophy) proceeds from examples.

E.g. From "This swan is white, that swan is white, in fact every swan I have seen so far is white"

Conclude: "Every Swan is white"

WeChat: cstutorcs

Assignment Project Exam Help

NB

Email: tutorcs@163.com

This may be a good way to discover hypotheses.

But it is not a valid principle of reasoning!

QQ: 749389476

Mathematical induction is a variant that is valid.

<https://tutorcs.com>

Mathematical Induction

程序代写代做 CS编程辅导

Mathematical Induction is based not just on a set of examples, but also a rule for deriving new cases of $P(x)$ from cases for which P is known to hold.



General structure of $P(x)$ by mathematical induction:

Base Case [B]: $P(a_1), P(a_2), \dots, P(a_n)$ for some small set of examples $a_1 \dots a_n$ (often $n = 1$)

Inductive Step [I]: A general rule showing that if $P(x)$ holds for some cases $x = x_1, \dots, x_k$, then $P(y)$ holds for some new case y , constructed in some way from x_1, \dots, x_k .

Conclusion: Starting with $a_1 \dots a_n$ and repeatedly applying the construction of y from existing values, we can eventually construct all values in the domain of interest.

QQ: 749389476

<http://tutorcs.com>

Induction proof structure

Let $P(x)$ be the proposition that ...

We will show that $P(x)$ holds for all x by induction on x .

Base case: $x = \dots$

- $P(x)$: ...
-
- so $P(x)$ holds.



[Repeat for all base cases]

WeChat: cstutorcs

Inductive case:

Assignment Project Exam Help

- Assume $P(x)$ holds. That is, ...
- We will show $P(y)$ holds.
- ...
- So $P(x)$ implies $P(y)$.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

[Repeat for all inductive cases]

Therefore, by induction, $P(x)$ holds for all x .

Outline

程序代写代做 CS编程辅导

Motivation

Basic Induction



Variations on Basic Induction

WeChat: cstutorcs

Structural Induction

Assignment Project Exam Help

Feedback

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Basic induction

程序代写代做 CS编程辅导



Basic induction is the principle applied to the natural numbers.

Goal: Show $P(n)$ holds for all $n \in \mathbb{N}$.

Approach: Show that:

Base case (B): $P(0)$ holds; and

Inductive case (I): If $P(k)$ holds then $P(k + 1)$ holds.

QQ: 749389476

<https://tutorcs.com>

Email: tutorcs@163.com

Assignment Project Exam Help

WeChat: [tutorcs](https://tutorcs.com)

Example

Recall the recursive program:

程序代写代做 CS编程辅导

Example

Summing the first n numbers:



```
if( $n = 0$ ): 0
```

Website: tutorcs.com

Assignment Project Exam Help

Another attempt:

Email: tutorcs@163.com

Example

QQ: 749389476

```
sum2( $n$ ):
```

<http://tutorcs.com>

```
return  $n * (n + 1) / 2$ 
```

Induction proof **guarantees** that these programs will behave the same.

Example

Let $P(n)$ be the proposition that:

程序代写代做 CS编程辅导

$$P(n): \sum_{i=0}^n i = \frac{n(n+1)}{2}.$$

We will show that $P(n)$ is true for all $n \in \mathbb{N}$ by induction on n .



Proof.

[B] $P(0)$, i.e.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorscs@163.com

[I] $\forall k \geq 0 (P(k) \rightarrow P(k+1))$, i.e.

QQ: 749389476

$$\sum_{i=0}^k i = \frac{k(k+1)}{2} \rightarrow \sum_{i=0}^{k+1} i = \frac{(k+1)(k+2)}{2}$$

<https://tutorscs.com>

(proof?)



Example (cont'd)

程序代写代做 CS编程辅导

Proof.

Inductive step [I]:



$$\sum_{i=0}^{k+1} i = \left(\sum_{i=0}^k i \right) + (k+1)$$

WeChat: cstutorcs

$$= \frac{k(k+1)}{2} + (k+1) \quad (\text{by the inductive hypothesis})$$

Assignment Project Exam Help

$$= \frac{k(k+1) + 2(k+1)}{2}$$

Email: tutorcs@163.com

QQ: 749389476

$$= \frac{(k+1)(k+2)}{2}$$

<https://tutorcs.com>



Outline

程序代写代做 CS编程辅导

Motivation

Basic Induction

Variations on Basic Induction

Structural Induction

Feedback



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Variations

程序代写代做 CS编程辅导



① Induction from $n=1$

② Induction steps > 1

③ Strong induction

④ Backward induction

⑤ Forward-backward induction

⑥ Structural induction

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Induction From m Upwards

程序代写代做 CS编程辅导



If

[B] $P(m)$

[I] $\forall k \geq m (P(k) \rightarrow P(k+1))$

then

[C] $\forall n \geq m (P(n))$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example

程序代写代做 CS编程辅导



Theorem. For all n the number $8^n - 2^n$ is divisible by 6.

[B] $8^1 - 2^1$ is divisible by 6

[I] if $8^k - 2^k$ is divisible by 6, then so is $8^{k+1} - 2^{k+1}$, for all $k \geq 1$

Prove [I] using the “trick” to rewrite 8^{k+1} as $8 \cdot (8^k - 2^k + 2^k)$ which allows you to apply the IH on $8^k - 2^k$

QQ: 749389476

<https://tutorcs.com>

WhatsApp: [tutorcs](https://tutorcs.com)
Assignment Project Exam Help

Email: tutorcs@163.com

Induction Steps $\ell > 1$

程序代写代做 CS编程辅导



If

[B] $P(m)$

[I] $P(k) \rightarrow P(k + \ell)$ for all $k \geq m$

then

[C] $P(n)$ for every ℓ 'th $n \geq m$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example

程序代写代做 CS编程辅导



Every 4th Fibonacci number is divisible by 3.

[B] $F_4 = 3$ is divisible by 3

[I] if $3 \mid F_k$, then $3 \mid F_{k+4}$, for all $k \geq 4$

Prove [I] by rewriting F_{k+4} in such a way

that you can apply the IH on F_k

QQ: 749389476

<https://tutorcs.com>

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

Strong Induction

程序代写代做 CS编程辅导



This is a version in which the inductive hypothesis is stronger. Rather than using the fact that $P(k)$ holds for a single value, we use *all* values up to k .

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

If

[B] $P(m)$

[I] $[P(m) \wedge P(m+1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$ for all $k \geq m$

then

[C] $P(n)$, for all $n \geq m$

Example

程序代写代做 CS编程辅导



Claim: All integers ≥ 2 can be written as a product of primes.

[B] 2 is a product of primes

[I] If all x with $2 \leq x \leq k$ can be written as a product of primes, then $k + 1$ can be written as a product of primes, for all $k \geq 2$

Proof for [I]?

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Negative Integers, Backward Induction

NB

程序代写代做 CS编程辅导

Induction can be conducted over any subset of \mathbb{Z} with least element. Thus m can be negative; eg. base case $m = -10^6$.



NB

One can apply induction in the 'opposite' direction $p(m) \rightarrow p(m-1)$. It means considering the integers with the opposite ordering where the next number after n is $n-1$. Such induction would be used to prove some $p(n)$ for all $n \leq m$.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

NB

QQ: 749389476

Sometimes one needs to reason about all integers \mathbb{Z} . This requires two separate simple induction proofs: one for \mathbb{N} , another for $-\mathbb{N}$. They both would start from some initial values, which could be the same, e.g. zero. Then the first proof would proceed through positive integers; the second proof through negative integers.

<https://tutorcs.com>

Forward-Backward Induction

程序代写代做 CS编程辅导

Idea

To prove $P(n)$ for all

- verify $P(k_0)$
- prove $P(k_i)$ for infinitely many $k_0 < k_1 < k_2 < k_3 < \dots$
- fill the gaps

$$P(k_1) \rightarrow P(k_1 - 1) \rightarrow P(k_1 - 2) \rightarrow \dots \rightarrow P(k_0 + 1)$$

$$P(k_2) \rightarrow P(k_2 - 1) \rightarrow P(k_2 - 2) \rightarrow \dots \rightarrow P(k_1 + 1)$$

.....

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorms@163.com

NB

QQ: 749389476

This form of induction is extremely important for the analysis of algorithms.

<https://tutorms.com>



Outline

程序代写代做 CS编程辅导



Motivation

Basic Induction

Variations on Basic Induction

Structural Induction

Feedback

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Structural Induction

Basic induction allows us to assert properties over **all natural numbers**. The induction scheme (layout) uses the recursive definition of \mathbb{N} .

The induction scheme is applied not only to natural numbers (and integers) but to any partially ordered set in general – especially those defined recursively.

The basic approach is always the same – we need to verify that

- **[B]** the property holds for all minimal objects — objects that have no predecessors; they are usually very simple objects allowing immediate verification
- **[I]** for any given object, if the property in question holds for all its predecessors (‘smaller’ objects) then it holds for the object itself

程序代写代做 CS编程辅导



WeChat: tutormcs

Assignment Project Exam Help

Email: tutormcs@163.com

QQ: 749389476

https://tutormcs.com

Example: Induction on Σ^*

程序代写代做 CS编程辅导

Recall definition of Σ

$$\lambda \in \Sigma^*$$

If $w \in \Sigma^*$ for all $a \in \Sigma$



Structural induction on Σ^*

WeChat: cstutorcs

Goal: Show $P(w)$ holds for all $w \in \Sigma^*$

Assignment Project Exam Help

Approach: Show that

Email: tutorcs@163.com

Base case (B): $P(\lambda)$ holds; and

QQ: 749389476

Inductive case (I): If $P(w)$ holds then $P(aw)$ holds for all

$a \in \Sigma$. <https://tutorcs.com>

Example: Induction on Σ^*

Recall:

程序代写代做 CS编程辅导

Formal definition of Σ^* :

$$\lambda \in \Sigma^*$$

If $w \in \Sigma^*$ for all $a \in \Sigma$



Formal definition of concatenation:

(concat.B) $\lambda v = v$ **WeChat: cstutorcs**

(concat.I) $(aw)v = a(wv)$ **Assignment Project Exam Help**

Formal definition of length:

(length.B) $\text{length}(\lambda) = 0$

(length.I) $\text{length}(aw) = 1 + \text{length}(w)$ **Email: tutorcs@163.com**

QQ: 749389476

Prove: <https://tutorcs.com>

$$\text{length}(wv) = \text{length}(w) + \text{length}(v)$$

Example: Induction on Σ^*

程序代写代做 CS编程辅导

Let $P(w)$ be the proposition that, for all $v \in \Sigma^*$:

$$\text{length}(\lambda v) = \text{length}(w) + \text{length}(v).$$

We will show that $P(w)$ holds for all $w \in \Sigma^*$ by **structural induction on w** .

WeChat: cstutorcs

Proof:

Assignment Project Exam Help

Base case ($w = \lambda$):

Email: tutorcs@163.com

$$\begin{aligned} \text{length}(\lambda v) &= \text{length}(v) && \text{(concat.B)} \\ &= 0 + \text{length}(v) && \text{QQ: 749389476} \\ &= \text{length}(\lambda) + \text{length}(v) && \text{https://tutorcs.com} \quad \text{(length.B)} \end{aligned}$$

Example: Induction on Σ^*

Proof cont'd:

程序代写代做 CS编程辅导

Inductive case ($w = aw'$): Assume that $P(w')$ holds. That is, for all $v \in \Sigma^*$:



$$(IH): \quad \text{length}(w'v) = \text{length}(w') + \text{length}(v).$$

Then, for all $a \in \Sigma$, we have:

WeChat: cstutorcs

$$\begin{aligned} \text{length}((aw')v) &= \text{length}(a(w'v)) && (\text{concat. I}) \\ &= 1 + \text{length}(w'v) && (\text{length. I}) \\ &= 1 + \text{length}(w') + \text{length}(v) && (IH) \\ &= \text{length}(aw') + \text{length}(v) && (\text{length. I}) \end{aligned}$$

QQ: 749389476

So $P(aw')$ holds. <https://tutorcs.com>

We have $P(\lambda)$ and for all $w' \in \Sigma^*$ and $a \in \Sigma$: $P(w') \rightarrow P(aw')$.
Hence $P(w)$ holds for all $w \in \Sigma^*$.

Example 2: Induction on Σ^*

程序代写代做 CS编程辅导

Recall $\text{append} : \Sigma^* \times \Sigma^* \rightarrow \Sigma^*$ defined as:

- $\text{append}(\lambda, x) = x$
- $\text{append}(aw, x) = a(\text{append}(w, x))$

WeChat: cstutorcs

Prove:

For all $w, v \in \Sigma^*$ and $x \in \Sigma^*$:

$$\text{append}(wv, x) = w(\text{append}(v, x))$$

QQ: 749389476

<https://tutorcs.com>

Assignment Project Exam Help

Email: tutorcs@163.com

Example 2: Induction on Σ^*

程序代写代做 CS编程辅导



Theorem

For all $w, v \in \Sigma^*$ and x , $\text{append}(wv, x) = w(\text{append}(v, x))$.

Proof: By induction on w .

[B] $\text{append}(\lambda v, x) = \text{append}(v, x)$ (concat.B)

[I] $\text{append}((aw)v, x) = \text{append}(a(wv), x)$ (concat.I)

$= a \text{append}(wv, x)$ (append.I)

$= a(w \text{append}(v, x))$ (IH)

$= (aw) \text{append}(v, x)$ (concat.I)

<https://tutorcs.com>

WeChat: cstutorcs
Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

Example 3: Induction on Σ^*

程序代写代做 CS编程辅导



Define $\text{rev} : \Sigma^* \rightarrow \Sigma^*$:

(rev.B) $\text{rev}(\lambda) = \lambda$, **WeChat: cstutorcs**

(rev.I) $\text{rev}(a \cdot w) = \text{append}(\text{reverse}(w), a)$ **Assignment Project Exam Help**

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3: Induction on Σ^*

程序代写代做 CS编程辅导

Theorem

For all $w, v \in \Sigma^*$, $\text{rev}(v \cdot w) = \text{reverse}(v) \cdot \text{reverse}(w)$.



Proof: By induction

$$\begin{aligned} \text{[B]} \quad \text{rev}(\lambda v) &= \text{rev}(v) && \text{(concat.B)} \\ &= \text{rev}(v) \lambda && (*) \end{aligned}$$

WeChat: cstutorcs

$$= \text{rev}(v) \text{rev}(\lambda) \quad \text{(rev.B)}$$

Assignment Project Exam Help

$$\text{[I]} \quad \text{rev}((aw')v) = \text{rev}(a(w'v)) \quad \text{(concat.I)}$$

Email: tutors@163.com

$$= \text{append}(\text{rev}(w'v), a) \quad \text{(rev.I)}$$

QQ: 749389476

$$= \text{append}(\text{rev}(v) \text{rev}(w'), a) \quad \text{(IH)}$$

$$= \text{rev}(v) \text{append}(\text{rev}(w'), a) \quad \text{(Example 2)}$$

https://tutapps.com

$$= \text{rev}(v) \text{rev}(aw') \quad \text{(rev.I)}$$

Example 4: Induction on more complex structures

Recall expressions in the Proof assistant. 程序代写代做 CS编程辅导

- (B) A, B, \dots, Z  are expressions
- (B) \emptyset and \mathcal{U} are expressions
- (R) If E is an expression then so is (E) and E^c
- (R) If E_1 and E_2 are expressions then:

- $(E_1 \cup E_2)$, WeChat: cstutorcs
- $(E_1 \cap E_2)$, Assignment Project Exam Help
- $(E_1 \setminus E_2)$, and
- $(E_1 \oplus E_2)$ are expressions. Email: tutores@163.com

Theorem

QQ: 749389476

In any valid expression, the number of (equals the number of)

https://tutores.com

Proof: By induction on the structure of E ...

Exercise

程序代写代做 CS编程辅导

Exercise



RW: 4.4.2 Define $s_1 = 1$, $s_{n+1} = \frac{1}{1+s_n}$ for $n \geq 1$

Then $s_1 = 1$, $s_2 = \frac{1}{2}$, $s_3 = \frac{1}{3}$, $s_4 = \frac{3}{5}$, $s_5 = \frac{5}{8}$, ...

The numbers in numerator and denominator remind one of the Fibonacci sequence.

Prove by induction that

Email: tutores@163.com

QQ: 749389476

<https://tutores.com>

Outline

程序代写代做 CS编程辅导



Motivation

Basic Induction

Variations on Basic Induction

Structural Induction

Feedback

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Weekly Feedback

程序代写代做 CS编程辅导

I would appreciate any comments/suggestions/requests you have on this week's lecture



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com



QQ: 749389476
<https://forms.office.com/r/xKKrxYMRn9>

<https://tutorcs.com>