

# COMP9418: Advanced Topics in Statistical Machine Learning

Probability Calculus  
Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

Instructor: Gustavo Batista

University of New South Wales

# Introduction

---

- In this lecture, we introduce probability calculus
  - Framework for reasoning with uncertain beliefs
  - Each event is assigned a degree of belief which quantifies the belief in that event
- This topic covered in this lecture will be fundamental to build a framework for reasoning with uncertainty
  - Degree of belief updating
  - The notion of independence

Assignment Project Exam Help

<https://tutorcs.com>


WeChat: cstutorcs

# Degrees of Belief

- A propositional knowledge base  $\Delta$  classifies sentences into three categories
  - Sentences implied by  $\Delta$
  - Sentences whose negations are implied by  $\Delta$
  - All other sentences
- We can obtain a much finer classification of sentences through a finer classification of worlds
  - Assigning a *degree of belief* or *probability* in  $[0,1]$  to each world  $w$  ( $P(w)$ )
  - The belief in, or probability of, a sentence  $\alpha$  is defined as

$$P(\alpha) \stackrel{\text{def}}{=} \sum_{w \models \alpha} P(w)$$

- That is, the sum of probabilities assigned to worlds at which  $\alpha$  is true



# Degrees of Belief

- This table shows a *state of belief* or *joint probability distribution*
  - We require the degrees of belief assigned to all words sum up to 1
  - This is a normalization convention
  - It allows to directly compare the degrees of belief held by different states
- The joint probability distribution is usually too large for direct representation
  - 20 binary variables – 1,048,576 entries
  - 40 binary variables – 1,099,511,627,776 entries
  - We will deal with this issue by using Graphical Models

world	Earthquake	Burglary	Alarm	$P(\cdot)$
$w_1$	true	true	true	.0190
$w_2$	true	true	false	.0010
$w_3$	true	false	true	.0560
$w_4$	true	false	false	.0240
$w_5$	false	true	true	.1620
$w_6$	false	true	false	.0180
$w_7$	false	false	true	.0072
$w_8$	false	false	false	.7128

$$\begin{aligned}P(\text{Earthquake}) &= P(w_1) + P(w_2) + P(w_3) + P(w_4) = .1 \\P(\text{Burglary}) &= .2 \\P(\neg \text{Burglary}) &= .8 \\P(\text{Alarm}) &= .2442\end{aligned}$$

# Properties of Beliefs

- Some degrees of belief (or simply, beliefs) properties

- Beliefs are bounded for any sentence  $\alpha$  into the  $[0,1]$  interval

$$0 \leq P(\alpha) \leq 1$$

- An inconsistent sentence  $\alpha$  have belief equal to zero

$$P(\alpha) = 0 \text{ when } \alpha \text{ is inconsistent}$$

- A valid sentence  $\alpha$  has belief equal to one


$$P(\alpha) = 1 \text{ when } \alpha \text{ is valid}$$

- We can compute the belief of a sentence given the belief in its negation

$$P(\alpha) + P(\neg\alpha) = 1$$

- We can also compute the belief in a disjunction

$$P(\alpha \vee \beta) = P(\alpha) + P(\beta) - P(\alpha \wedge \beta)$$



# Quantifying Uncertainty

- This table summarises the beliefs associated with the alarm example

	Earthquake	Burglary	Alarm
true	.1	.2	.2442
false	.9	.8	.7558
$H(\cdot)$	.469	.722	.802

- The beliefs seem most certain that an Earthquake has occurred
- We are least certain if an alarm has triggered

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs


- We can quantify uncertainty using entropy

- Where  $0 \log 0 = 0$  by convention

$$H(X) = - \sum_x P(X) \log_2 P(X)$$

- This plot shows the entropy for different probabilities

- When  $p = 0$  or  $p = 1$ , the entropy is zero, indicating no uncertainty
- When  $p = \frac{1}{2}$ , the entropy is at a maximum value



# Updating Beliefs

- Suppose now that we know that the alarm has triggered
  - $Alarm = true$
- We need to accommodate the state of belief to this new piece of information we call *evidence*
  - Evidence is represented by an event, say  $\beta$
  - Our objective is to update the state of belief  $P(\cdot)$  into a new state of belief we will denote  $P(\cdot | \beta)$

world	Earthquake	Burglary	Alarm	$P(\cdot)$	$P(\cdot   Alarm)$
$w_1$	true	true	true	.0190	
$w_2$	true	true	false	.0010	
$w_3$	true	false	true	.0560	
$w_4$	true	false	false	.0240	
$w_5$	false	true	true	.1620	
$w_6$	false	true	false	.0180	
$w_7$	false	false	true	.0072	
$w_8$	false	false	false	.7128	

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

# Updating Beliefs

- Given that  $\beta$  is known for sure

- The new state of belief  $P(\cdot | \beta)$  should assign a belief 1 to  $\beta$
- $P(\beta | \beta) = 1$

- This implies that  $P(\neg\beta | \beta) = 0$

- This implies that every world  $w$  that satisfies  $\neg\beta$  must be assigned to belief 0

$$P(w | \beta) = 0 \quad \text{for all } w \models \neg\beta$$

world	Earthquake	Burglary	Alarm	$P(\cdot)$	$P(\cdot   Alarm)$
$w_1$	true	true	true	.0190	
$w_2$	true	true	false	.0010	0
$w_3$	true	false	true	.0560	
$w_4$	true	false	false	.0240	0
$w_5$	false	true	true	.1620	
$w_6$	false	true	false	.0180	0
$w_7$	false	false	true	.0072	
$w_8$	false	false	false	.7128	0

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs



# Updating Beliefs

- We know that

$$\sum_{w \models \beta} P(w|\beta) = 1$$

- But these leaves many options for the  $P(w|\beta)$  for  $w$ 's that satisfies  $\beta$
- It is reasonable to perturb the beliefs as little as possible, so
  - $P(w|\beta) = 0$  for all  $w$  where  $P(w) = 0$
  - $\frac{P(w)}{P(w')} = \frac{P(w|\beta)}{P(w'|\beta)}$  for all  $w, w' \models \beta, P(w) > 0, P(w') > 0$
- With these three constraints, the only options for the new beliefs is
  - $P(w|\beta) = \frac{P(w)}{P(\beta)}$  for all  $w \models \beta$

# Updating Beliefs

- The new beliefs are just the normalization of old beliefs

- The normalization constant is  $P(\beta)$

$$P(w|\beta) \stackrel{\text{def}}{=} \begin{cases} 0 & \text{if } w \models \neg\beta \\ \frac{P(w)}{P(\beta)} & \text{if } w \models \beta \end{cases}$$

- The new state of belief is referred as *conditioning* the old state  $P$  on evidence  $\beta$

world	Earthquake	Burglary	Alarm	$P(\cdot)$	$P(\cdot   Alarm)$
$w_1$	true	true	true	.0190	.0190/.2442
$w_2$	true	true	false	.0010	0
$w_3$	true	false	true	.0560	.0560/.2442
$w_4$	true	false	false	.0240	0
$w_5$	false	true	true	.1620	.1620/.2442
$w_6$	false	true	false	.0180	0
$w_7$	false	false	true	.0072	.0072/.2442
$w_8$	false	false	false	.7128	0

# Updating Beliefs: Example

- Let us suppose  $Alarm = true$  and compute some conditional probabilities
  - $P(Burglary) =$
  - $P(Burglary|Alarm) =$
- Also,
  - $P(Earthquake) =$
  - $P(Earthquake|Alarm) =$

world	Earthquake	Burglary	Alarm	$P(\cdot)$	$P(\cdot   Alarm)$
$w_1$	true	true	true	.0190	.0778
$w_2$	true	true	false	.0010	0
$w_3$	true	false	true	.0560	.2293
$w_4$	true	false	false	.0240	0
$w_5$	false	true	true	.1620	.6634
$w_6$	false	true	false	.0180	0
$w_7$	false	false	true	.0072	.0295
$w_8$	false	false	false	.7128	0

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

# Updating Beliefs: Example

- Let us suppose  $Alarm = true$  and compute some conditional probabilities

- $P(Burglary) = .2$
- $P(Burglary|Alarm) = .741$

- Also,

- $P(Earthquake) = .1$
- $P(Earthquake|Alarm) = .307$

world	Earthquake	Burglary	Alarm	$P(\cdot)$	$P(\cdot   Alarm)$
$w_1$	true	true	true	.0190	.0778
$w_2$	true	true	false	.0010	0
$w_3$	true	false	true	.0560	.2293
$w_4$	true	false	false	.0240	0
$w_5$	false	true	true	.1620	.6634
$w_6$	false	true	false	.0180	0
$w_7$	false	false	true	.0072	.0295
$w_8$	false	false	false	.7128	0

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

# Updating Beliefs: Closed Form

- There is a simple closed form to compute the updated belief

$$\begin{aligned}
 P(\alpha|\beta) &= \sum_{w \models \alpha} P(w|\beta) \\
 &= \sum_{w \models \alpha, w \models \beta} P(w|\beta) + \sum_{w \models \alpha, w \models \neg \beta} P(w|\beta) \\
 &= \sum_{w \models \alpha, w \models \beta} P(w|\beta) \\
 &= \sum_{w \models \alpha \wedge \beta} P(w|\beta) \\
 &= \sum_{w \models \alpha \wedge \beta} P(w)/P(\beta)
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{P(\beta)} \sum_{w \models \alpha \wedge \beta} P(w) \\
 &= \frac{P(\alpha, \beta)}{P(\beta)}
 \end{aligned}$$

$$P(\alpha|\beta) = \frac{P(\alpha \wedge \beta)}{P(\beta)}$$

This equation is known as *Bayes' conditioning*. It is only defined when  $P(\beta) \neq 0$

# Bayes Conditioning

- Bayes conditioning follows the following commitments:

- Worlds that contradict the evidence  $\beta$  will have zero probability
- Worlds that have zero probability continue to have zero probability
- Worlds that are consistent with evidence  $\beta$  and have positive probability will maintain their relative beliefs

Earthquake	Burglary	Alarm	$P(\cdot)$
true	true	true	.0190
true	true	false	.0010
true	false	true	.0560
true	false	false	.0240
false	true	true	.1620
false	true	false	.0180
false	false	true	.0072
false	false	false	.7128

	Earthquake	Burglary	Alarm
true	.1	.2	.2442
false	.9	.8	.7558

# Bayes Conditioning: Example

- Let us analyse how some beliefs would change given evidence *Earthquake*

- $P(\text{Burglary}) =$

- $P(\text{Burglary}|\text{Earthquake}) =$

- Also,

- $P(\text{Alarm}) =$

- $P(\text{Alarm}|\text{Earthquake}) =$

- Now, considering evidence *Burglary*

- $P(\text{Alarm}) =$

- $P(\text{Alarm}|\text{Burglary}) =$

- Also,

- $P(\text{Earthquake}) =$

- $P(\text{Earthquake}|\text{Burglary}) =$

Earthquake	Burglary	Alarm	$P(\cdot)$
true	true	true	.0190
true	true	false	.0010
true	false	true	.0560
true	false	false	.0240
false	true	true	.1620
false	true	false	.0180
false	false	true	.0072
false	false	false	.7128

	Earthquake	Burglary	Alarm
true	.1	.2	.2442
false	.9	.8	.7558

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

# Bayes Conditioning: Example

- Let us analyse how some beliefs would change given evidence *Earthquake*

- $P(\text{Burglary}) = .2$
- $P(\text{Burglary}|\text{Earthquake}) = .2$

- Also,

- $P(\text{Alarm}) = .2442$
- $P(\text{Alarm}|\text{Earthquake}) \approx .75$

- Now, considering evidence *Burglary*

- $P(\text{Alarm}) = .2442$
- $P(\text{Alarm}|\text{Burglary}) \approx .905$

- Also,

- $P(\text{Earthquake}) = .1$
- $P(\text{Earthquake}|\text{Burglary}) = .1$

Earthquake	Burglary	Alarm	$P(\cdot)$
true	true	true	.0190
true	true	false	.0010
true	false	true	.0560
true	false	false	.0240
false	true	true	.1620
false	true	false	.0180
false	false	true	.0072
false	false	false	.7128

	Earthquake	Burglary	Alarm
true	.1	.2	.2442
false	.9	.8	.7558



# Bayes Conditioning: Example

- These beliefs dynamics are a property of the state of belief in this table

- It may not hold for other states of belief
- For instance, we can conceive a state of belief in which evidence *Earthquake* would change the belief about *Burglary*

- A central question is synthesizing states of beliefs that are *faithful*

- For instance, those that correspond to the beliefs held by some human expert
- We will see this is a central issue of modelling a real problem

Earthquake	Burglary	Alarm	$P(\cdot)$
true	true	true	.0190
true	true	false	.0010
true	false	true	.0560
true	false	false	.0240
false	true	true	.1620
false	true	false	.0180
false	false	true	.0072
false	false	false	.7128

	Earthquake	Burglary	Alarm
true	.1	.2	.2442
false	.9	.8	.7558

# Bayes Conditioning: Example

- Let us look at one more example as we add more evidence

- $P(\text{Burglary}|\text{Alarm}) =$

- $P(\text{Burglary}|\text{Alarm} \wedge \text{Earthquake}) =$

- Also,

- $P(\text{Burglary}|\text{Alarm} \wedge \neg \text{Earthquake}) =$

Earthquake	Burglary	Alarm	$P(\cdot)$
true	true	true	.0190
true	true	false	.0010
true	false	true	.0560
true	false	false	.0240
false	true	true	.1620
false	true	false	.0180
false	false	true	.0072
false	false	false	.7128

	Earthquake	Burglary	Alarm
true	.1	.2	.2442
false	.9	.8	.7558

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

# Bayes Conditioning: Example

- Let us look at one more example as we add more evidence

- $P(\text{Burglary}|\text{Alarm}) \approx .741$
  - $P(\text{Burglary}|\text{Alarm} \wedge \text{Earthquake}) \approx .253$

- Also,

- $P(\text{Burglary}|\text{Alarm} \wedge \neg \text{Earthquake}) \approx .957$

Earthquake	Burglary	Alarm	$P(\cdot)$
true	true	true	.0190
true	true	false	.0010
true	false	true	.0560
true	false	false	.0240
false	true	true	.1620
false	true	false	.0180
false	false	true	.0072
false	false	false	.7128

	Earthquake	Burglary	Alarm
true	.1	.2	.2442
false	.9	.8	.7558

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

# Conditional Entropy

- The *conditional entropy* of a variable  $X$  given another variable  $Y$

- It quantifies the average uncertainty about  $X$  after observing  $Y$

- We can show that the entropy never increases after conditioning

$$H(X|Y) \leq H(X)$$

- Although for a specific value  $y$  we may have  $H(X|y) > H(X)$

$$H(X|Y) \stackrel{\text{def}}{=} \sum_y P(y)H(X|y)$$

$$H(X|y) \stackrel{\text{def}}{=} - \sum_x P(x|y) \log_2 P(x|y)$$

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

	$B$	$B a$	$B \neg a$
true	.2	.741	.025
false	.8	.259	.975
$H(\cdot)$	.722	.825	.169

# Conditional Entropy

- The *conditional entropy* of a variable  $X$  given another variable  $Y$

- It quantifies the average uncertainty about  $X$  after observing  $Y$

Assignment Project Exam Help

- We can show that the entropy never increases after conditioning

<https://tutorcs.com>

$$H(X|Y) \leq H(X)$$

- Although for a specific value  $y$  we may have  $H(X|y) > H(X)$

- $H(B|A) = H(B|a)P(a) + H(B|\neg a)P(\neg a)$   
 $= .329$

$$H(X|Y) \stackrel{\text{def}}{=} \sum_y P(y)H(X|y)$$

$$H(X|y) \stackrel{\text{def}}{=} - \sum_x P(x|y) \log_2 P(x|y)$$

WeChat: cstutorcs

	$B$	$B a$	$B \neg a$
true	.2	.741	.025
false	.8	.259	.975
$H(\cdot)$	.722	.825	.169

# Independence

- We observed that evidence *Burglary* does not change belief in *Earthquake*

$$P(\text{Earthquake}) = .1$$

- More generally, we say that event  $\alpha$  is independent of event  $\beta$  iff

$$P(\text{Earthquake}|\text{Burglary}) = .1$$

- $P(\alpha|\beta) = P(\alpha)$  or
- $P(\beta) = 0$

Assignment Project Exam Help

<https://tutorcs.com>

- We also found *Burglary* independent of *Earthquake*

$$P(\text{Burglary}) = .2$$

- It is indeed a general property

$$P(\text{Burglary}|\text{Earthquake}) = .2$$

- If  $\alpha$  is independent of  $\beta$  then  $\beta$  is independent of  $\alpha$

- $\alpha$  and  $\beta$  are independent iff

- $P(\alpha \wedge \beta) = P(\alpha)P(\beta)$
- This equation is often taken as the definition of independence

WeChat: csTutorcs

# Independence

- We observed that evidence *Burglary* does not change belief in *Earthquake*

$$P(\text{Earthquake}) = .1$$

- More generally, we say that event  $\alpha$  is independent of event  $\beta$  iff

$$P(\text{Earthquake}|\text{Burglary}) = .1$$

- $P(\alpha|\beta) = P(\alpha)$  or
- $P(\beta) = 0$

Assignment Project Exam Help

<https://tutorcs.com>

- We also found *Burglary* independent of *Earthquake*

$$P(\text{Burglary}) = .2$$

- It is indeed a general property

$$P(\text{Burglary}|\text{Earthquake}) = .2$$

- If  $\alpha$  is independent of  $\beta$  then  $\beta$  is independent of  $\alpha$

- $\alpha$  and  $\beta$  are independent iff

- $P(\alpha \wedge \beta) = P(\alpha)P(\beta)$
- This equation is often taken as the definition of independence

# Independence and Mutual Exclusiveness

- Independence and mutual exclusiveness are *not* the same notion
- Two events  $\alpha$  and  $\beta$  are mutually exclusive (logically disjoint) iff they do not share any models
  - $Mods(\alpha) \cap Mods(\beta) = \emptyset$
  - They cannot hold together at the same world
- Events  $\alpha$  and  $\beta$  are independent iff  $P(\alpha \wedge \beta) = P(\alpha)P(\beta)$

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs



# Conditional Independence

- Independence is a dynamic notion

- Two independent events may become dependent after some evidence

- For example, we saw that *Burglary* was independent of *Earthquake*. However, these events are dependent after accepting evidence *Alarm*

- This is expected since *Earthquake* and *Burglary* are competing explanations to *Alarm*

- Consider this table for another example

- We have two sensors that can detect the current state of temperature
- The sensors are noisy and have different reliabilities

$$P(\text{Burglary}|\text{Alarm}) \approx .741$$

$$P(\text{Burglary}|\text{Alarm} \wedge \text{Earthquake}) \approx .253$$

world	Temp	Sensor1	Sensor2	$P(\cdot)$
$w_1$	normal	normal	normal	.576
$w_2$	normal	normal	extreme	.144
$w_3$	normal	extreme	normal	.064
$w_4$	normal	extreme	extreme	.016
$w_5$	extreme	normal	normal	.008
$w_6$	extreme	normal	extreme	.032
$w_7$	extreme	extreme	normal	.032
$w_8$	extreme	extreme	extreme	.128

# Conditional Independence

- We have the following initial beliefs

- $P(Temp = normal) = .80$
- $P(Sensor1 = normal) = .76$
- $P(Sensor2 = normal) = .68$

- If the first sensor reads normal

- Our belief that the second sensor also reads normal increases
- $P(Sensor2 = normal | Sensor1 = normal) \approx .768$
- Therefore the sensors readings are dependent

- However, they will become independent if we observe the temperature is normal

- $P(Sensor2 = normal | Temp = normal) = .80$
- $P(Sensor2 = normal | Temp = normal, Sensor1 = normal) = .80$

world	Temp	Sensor1	Sensor2	$P(\cdot)$
$w_1$	normal	normal	normal	.576
$w_2$	normal	normal	extreme	.144
$w_3$	normal	extreme	normal	.064
$w_4$	normal	extreme	extreme	.016
$w_5$	extreme	normal	normal	.008
$w_6$	extreme	normal	extreme	.032
$w_7$	extreme	extreme	normal	.032
$w_8$	extreme	extreme	extreme	.128

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

# Conditional Independence

- In general,
  - Independent event may become dependent given new evidence
  - Dependent events may become independent given new evidence

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

- Event  $\alpha$  is conditionally independent of event  $\beta$  given event  $\gamma$  iff
- Conditional independence is also symmetric
  - $\alpha$  is conditionally independent of  $\beta$  given  $\gamma$  iff  $\beta$  is conditionally independent of  $\alpha$  given  $\gamma$
- This equation is often taken as definition of conditional independence

$$P(\alpha|\beta \wedge \gamma) = P(\alpha|\gamma) \text{ or}$$

$$P(\gamma) = 0$$

$$P(\alpha \wedge \beta|\gamma) = P(\alpha|\gamma)P(\beta|\gamma) \text{ or}$$

$$P(\gamma) = 0$$

# Variable Independence

- It is useful to talk about independence of sets of variables

- Let  $X$ ,  $Y$  and  $Z$  be three disjoint sets of variables

- $X$  is independent of  $Y$  given  $Z$  is denoted by  $X \perp Y | Z$

- It means that  $x$  is independent of  $y$  given  $z$  for all instantiations of  $x$ ,  $y$  and  $z$

- For example, suppose that  $X = \{A, B\}$ ,  $Y = \{C\}$  and  $Z = \{D, E\}$

- Where  $A, B, C, D$  and  $E$  are propositional variables
- The statement  $X \perp Y | Z$  is a compact notation for several statements about independence

$$I_P(X, Z, Y)$$

$A \wedge B$  is independent of  $C$  given  $D \wedge E$   
 $A \wedge \neg B$  is independent of  $C$  given  $D \wedge E$   
 $\neg A \wedge \neg B$  is independent of  $C$  given  $D \wedge E$

...

$\neg A \wedge \neg B$  is independent of  $\neg C$  given  $\neg D \wedge \neg E$

# Mutual Information

- Independence is a special case of a more general notion known as *mutual information*

- Mutual information quantifies the impact of observing one variable on the uncertainty in another

$$MI(X; Y) \stackrel{\text{def}}{=} \sum_{x,y} P(x, y) \log_2 \frac{P(x, y)}{P(x)P(y)}$$

Assignment Project Exam Help

<https://tutorcs.com>

- Mutual information is

- Non-negative
- Equal to zero only if  $X$  and  $Y$  are independent
- It measures the extent to which observing one variable will reduce the uncertainty in another

WeChat: cstutorcs

$$MI(X; Y) = H(X) - H(X|Y)$$

$$MI(X; Y) = H(Y) - H(Y|X)$$

# Conditional Mutual Information

- Conditional mutual information can be defined as

$$MI(X; Y|Z) \stackrel{\text{def}}{=} \sum_{x,y,z} P(x, y, z) \log_2 \frac{P(x, y|z)}{P(x|z)P(y|z)}$$

- It has the following properties

Assignment Project Exam Help

<https://tutorcs.com>

$$MI(X; Y|Z) = H(X|Z) - H(X|Y, Z)$$

$$MI(X; Y|Z) = H(Y|Z) - H(Y|X, Z)$$

WeChat: cstutorcs

- Entropy and mutual information can be extended to sets of variables
  - For instance, entropy can be generalized to a set of variables  $\mathbf{X}$  as follows

$$H(\mathbf{X}) = - \sum_{\mathbf{x}} P(\mathbf{x}) \log_2 P(\mathbf{x})$$

# Conditional Probability for Multiple Variables

- We can extend the definition for conditional probabilities for multiple variables

$$P(A|B) = \frac{P(A, B)}{P(B)}$$

Bayes' conditioning

$$P(A, B) = P(A|B)P(B)$$

Product rule

- For three variables  $A$ ,  $B$  and  $C$ , we have

$$P(A, B|C) = \frac{P(A, B, C)}{P(C)}$$

<https://tutorcs.com>

$$P(A|B, C) = \frac{P(A, B, C)}{P(B, C)}$$

WeChat: cstutorcs

# Case Analysis


- Also known as *law of total probability*

$$P(\alpha) = \sum_{i=1}^n P(\alpha \wedge \beta_i)$$

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs



The events  $\beta_1, \dots, \beta_n$  are mutually exclusive and exhaustive



# Chain Rule

---

- It is the repeated application of Bayes conditioning

$$P(\alpha_1 \wedge \alpha_2 \wedge \cdots \wedge \alpha_n) = P(\alpha_1 | \alpha_2 \wedge \cdots \wedge \alpha_n) P(\alpha_2 | \alpha_3 \wedge \cdots \wedge \alpha_n) \dots P(\alpha_n)$$

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

# Bayes Rule

- *Bayes rule or Bayes theorem*

- The classical usage of this rule is when event  $\alpha$  is perceived to be a cause of event  $\beta$
- For example,  $\alpha$  is a disease and  $\beta$  is a symptom
- Our goal is to assess our belief in the cause given the effect

$$P(\alpha|\beta) = \frac{P(\beta|\alpha)P(\alpha)}{P(\beta)}$$

Assignment Project Exam Help

<https://tutorcs.com>

- *Example*

WeChat: cstutorcs

- Suppose that we have a patient who was just tested for a particular disease and the test came out positive. We know that one in every thousand people has this disease. We also know that the test is not reliable: it has a false positive rate of 2% and a false negative rate of 5%. Our goal is then to assess our belief in the patient having the disease given that the test came out positive.

# Conclusion

---

- In this lecture, we discussed some fundamental aspects of probabilistic calculus
  - Belief update
  - Independence and Conditional Independence
  - Bayes conditioning, case analysis,
  - Chain rule, Bayes rule
- Tasks
  - Read Chapter 3, but Sections 3.6 and 3.7 from the textbook (Darwiche)

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs