

# 程序代写代做 CS编程辅导

CS861: Theoretical Foundations of Machine Learning

Lecture 5 - 09/15/2023

University of Wisconsin-Madison, Fall 2023

Lec

Function and VC Dimension

Lecturer: Kirthy

Scribed by: Chenghui Zheng, Yixuan Zhang



**Disclaimer:** These notes are subject to the usual scrutiny reserved for formal publications. They may be distributed outside this class only with the permission of the instructor.

In this lecture, we will first bound the Radamacher complexity using the growth function. Then, we will introduce the VC dimension and provide some examples.

WeChat: cstutorcs

## 1 Bounding Rademacher Complexity Using the Growth Function

Assignment Project Exam Help

First, we will prove Massart's lemma, which upper bounds the empirical Radamacher complexity.

**Lemma 1** (Massart's Lemma). Let  $S = \{(x_1, y_1), \dots, (x_n, y_n)\} \in \{\mathcal{X} \times \mathcal{Y}\}^n$ , and  $\mathcal{H}$  be a hypothesis class. Then,

Email: tutorcs@163.com

$$\widehat{\text{Rad}}(S, \mathcal{H}) \leq \frac{1}{n} \left( \max_{v \in \mathcal{L}(S, \mathcal{H})} \|v\|_2 \right) \sqrt{2 \log(|\mathcal{L}(S, \mathcal{H})|)},$$

where  $\|v\|_2^2 = \sum_{i=1}^n v_i^2$ .

**Proof** First, observe that we can write

QQ: 749389476

$$\widehat{\text{Rad}}(S, \mathcal{H}) = \mathbb{E}_\sigma \left[ \sup_{h \in \mathcal{H}} \frac{1}{n} \sum_{i=1}^n \sigma_i \ell(h(x_i), y_i) \right] = \frac{1}{n} \mathbb{E}_\sigma \left[ \max_{v \in \mathcal{L}(S, \mathcal{H})} \sum_{i=1}^n \sigma_i v_i \right]. \quad (1)$$

https://tutorcs.com

Next, let  $s > 0$ , whose value we will specify later.

$$\begin{aligned} \mathbb{E}_\sigma \left[ \max_{v \in \mathcal{L}(S, \mathcal{H})} \sum_{i=1}^n \sigma_i v_i \right] &= \frac{1}{s} \mathbb{E}_\sigma \left[ \max_{v \in \mathcal{L}(S, \mathcal{H})} s \sum_{i=1}^n \sigma_i v_i \right] \\ &= \frac{1}{s} \mathbb{E}_\sigma \left[ \log \left( \exp \left( \max_{v \in \mathcal{L}(S, \mathcal{H})} s \sum_{i=1}^n \sigma_i v_i \right) \right) \right] \\ &\leq \frac{1}{s} \log \left( \mathbb{E}_\sigma \left[ \exp \left( \max_{v \in \mathcal{L}(S, \mathcal{H})} s \sum_{i=1}^n \sigma_i v_i \right) \right] \right) \quad \text{by Jensen's Inequality} \\ &\leq \frac{1}{s} \log \left( \mathbb{E}_\sigma \left[ \sum_{v \in \mathcal{L}(S, \mathcal{H})} \exp \left( s \sum_{i=1}^n \sigma_i v_i \right) \right] \right) \\ &\leq \frac{1}{s} \log \left( \sum_{v \in \mathcal{L}(S, \mathcal{H})} \mathbb{E}_\sigma \left[ \exp \left( s \sum_{i=1}^n \sigma_i v_i \right) \right] \right) \\ &\stackrel{(i)}{\leq} \frac{1}{s} \log \left( \sum_{v \in \mathcal{L}(S, \mathcal{H})} \exp \left( \frac{s^2}{2} \sum_{i=1}^n v_i^2 \right) \right) \end{aligned}$$

# 程序代写代做 CS编程辅导

$$\leq \frac{1}{s} \log \left( |\mathcal{L}(S, \mathcal{H})| \max_{v \in \mathcal{L}(S, \mathcal{H})} \exp \left( \frac{s^2}{2} \sum_{i=1}^n v_i^2 \right) \right) + \frac{s}{2} \max_{v \in \mathcal{L}(S, \mathcal{H})} \|v\|_2^2. \quad (2)$$

The inequality (i) holds for all  $s$ . Let  $\sigma_1, \dots, \sigma_n$  and  $\sigma_i$  is 1-subgaussian. Then,

$$\mathbb{E}_\sigma \left[ \max_{v \in \mathcal{L}(S, \mathcal{H})} \sum_{i=1}^n v_i \sigma_i \right] \leq \prod_{i=1}^n \mathbb{E}_\sigma [\exp((sv_i)\sigma_i)] \leq \prod_{i=1}^n \exp \left( \frac{s^2 v_i^2}{2} \right).$$

Equation (2) holds for all  $s$ , we can choose

$$s = \sqrt{\frac{2 \log |\mathcal{L}(S, \mathcal{H})|}{\max_{v \in \mathcal{L}(S, \mathcal{H})} \|v\|_2^2}}. \quad (3)$$

Equation (1), (2) and (3) imply

$$\widehat{\text{Rad}}_n(S, \mathcal{H}) \leq \frac{1}{s} \left( \max_{v \in \mathcal{L}(S, \mathcal{H})} \|v\|_2 \right) \cdot \sqrt{2 \log |\mathcal{L}(S, \mathcal{H})|}.$$

□

Email: tutorcs@163.com

Corollary 1.  $\forall S$  such that  $|S| = n$ , we have

$$\widehat{\text{Rad}}_n(S, \mathcal{H}) \leq \sqrt{\frac{2 \log(g(n, \mathcal{H}))}{n}}.$$

Moreover,

$$\text{Rad}_n(\mathcal{H}) \leq \sqrt{\frac{2 \log(g(n, \mathcal{H}))}{n}}.$$

**Proof**  $\|v\|_2 \leq \sqrt{n}$  and  $|\mathcal{L}(S, \mathcal{H})| \leq g(n, \mathcal{H})$  by definition of  $g(n, \mathcal{H})$ . The second statement follows by taking the expectation over  $S$  of the LHS of the first statement. □

To motivate the ensuing discussion about the VC dimension, recall that with probability at least  $1 - 2e^{-2n\epsilon^2}$

$$R(\hat{h}) \leq \inf_{h \in \mathcal{H}} R(h) + c_1 \text{Rad}_n(\mathcal{H}) + c_2 \epsilon.$$

Then, with fixed  $n, \delta$ , where  $\epsilon \in O(\sqrt{\frac{1}{n} \log(\frac{1}{\delta})})$ . From the previous lecture, we obtained  $g(n, \mathcal{H}) \leq 2^n$ . However, when  $g(n, \mathcal{H}) = 2^n$ ,  $\text{Rad}_n(\mathcal{H})$  will never goes to 0. At the very least, we hope to have:  $g(n, \mathcal{H}) \in o(2^n)$ , but ideally we would like to have  $g(n, \mathcal{H}) \in \text{poly}(n)$  so that  $\sqrt{\frac{\log(g(n, \mathcal{H}))}{n}} \lesssim \sqrt{\frac{\log(n)}{n}}$ .

## 2 VC dimension

In this section, we begin with the definition of Shattering.

# 程序代写代做 CS编程辅导

**Definition 1.** Let  $S^X = \{x_1, \dots, x_n\} \in X^n$  be a set of  $n$  points in  $X$ . We say that  $S^X$  is shattered by a hypothesis class  $\mathcal{H}$  if  $\mathcal{H}$  “can realize any label on  $S^X$ ”. That is



$$|H(S^X)| = 2^n,$$

where  $H(S^X) = \{[h(x_1), \dots, h(x_n)] \mid h \in \mathcal{H}\}$ .

Then, we give two examples of hypothesis classes under the same hypothesis class  $\mathcal{H}$ , which is the two-sided threshold classifiers:

$$\mathcal{H} = \{h_a(x) = \mathbb{1}_{\{x < a\}} \mid a \in \mathbb{R}\} \cup \{h_a(x) = \mathbb{1}_{\{x < a\}} \mid a \in \mathbb{R}\}.$$

**Example 1.** Consider  $S^X = \{x_1, x_2\}$  and we can assume  $x_1 < x_2$  without loss of generality. Therefore, we can try different classifiers in  $\mathcal{H}$  to achieve different labels.

- When we use  $h_a(x) = \mathbb{1}_{\{x \geq x_1 - 1\}}$ , the label is  $[1, 1]$ .
- When we use  $h_a(x) = \mathbb{1}_{\{x \geq \frac{x_1 + x_2}{2}\}}$ , the label is  $[0, 1]$ .
- When we use  $h_a(x) = \mathbb{1}_{\{x \geq x_2 + 1\}}$ , the label is  $[0, 0]$ .
- When we use  $h_a(x) = \mathbb{1}_{\{x < \frac{x_1 + x_2}{2}\}}$ , the label is  $[1, 0]$ .

Then,  $|H(S^X)| = 2^2$  and we can say  $S^X$  is shattered by  $\mathcal{H}$ .

**Example 2.** Consider  $S^X = \{x_1, x_2, x_3\}$  and we can assume  $x_1 < x_2 < x_3$  without loss of generality. We can do the similar thing as Example 1.

- When we use  $h_a(x) = \mathbb{1}_{\{x \geq x_1 - 1\}}$ , the label is  $[1, 1, 1]$ .
- When we use  $h_a(x) = \mathbb{1}_{\{x \geq \frac{x_1 + x_2}{2}\}}$ , the label is  $[0, 1, 1]$ .
- When we use  $h_a(x) = \mathbb{1}_{\{x \geq \frac{x_2 + x_3}{2}\}}$ , the label is  $[0, 0, 1]$ .
- When we use  $h_a(x) = \mathbb{1}_{\{x \geq x_3 + 1\}}$ , the label is  $[0, 0, 0]$ .
- When we use  $h_a(x) = \mathbb{1}_{\{x < \frac{x_1 + x_2}{2}\}}$ , the label is  $[1, 0, 0]$ .
- When we use  $h_a(x) = \mathbb{1}_{\{x < \frac{x_2 + x_3}{2}\}}$ , the label is  $[1, 1, 0]$ .

However, the label  $[0, 1, 0]$  and  $[1, 0, 1]$  can't be achieved by any  $h \in \mathcal{H}$ . Then,  $|H(S^X)| = 6 < 2^3$  and we can say  $S^X$  can't be shattered by  $\mathcal{H}$ .

After introducing the shattering, we are ready to give the definition of VC-dimension. Here we use  $d_{\mathcal{H}}$  to denote VC-dimension of  $\mathcal{H}$  and we will use  $d$  when  $\mathcal{H}$  is clear from context.

**Definition 2.** The VC-dimension  $d_{\mathcal{H}}$  of a hypothesis class  $\mathcal{H}$  is the size of the largest set shattered by  $\mathcal{H}$ .

Below we introduce three examples of VC-dimension.

**Example 3. Two-sided threshold classifiers**

By Example 1, we can obtain  $d \geq 2$ . By Example 2, we have  $d < 3$ . Therefore, we can conclude that  $d = 2$ .

**Example 4. One-sided threshold classifiers**

The hypothesis class  $\mathcal{H}$  is defined as

$$\mathcal{H} = \{h_a(x) = \mathbb{1}_{\{x \geq a\}} \mid a \in \mathbb{R}\}.$$

Similarly, we can show  $d = 1$  by showing  $d \geq 1$  and  $d < 2$ .

# 程序代写代做 CS编程辅导

1. Consider  $S^X = \{x_1\}$ .

- When we use  $h_1(x) = \mathbb{1}_{\{x \geq 0\}}$ , the label is  $[1]$ .
- When we use  $h_2(x) = \mathbb{1}_{\{x < 0\}}$ , the label is  $[0]$ .

Then,  $|H(S^X)| = 2$  is shattered by  $\mathcal{H}$ , which implies  $d \geq 1$ .

2. Consider  $S^X = \{x_1, x_2\}$  where  $x_1 < x_2$  without loss of generality.

- When we use  $h_1(x) = \mathbb{1}_{\{x \geq 0\}}$ , the label is  $[1, 1]$ .
- When we use  $h_2(x) = \mathbb{1}_{\{x < 0\}}$ , the label is  $[0, 1]$ .
- When we use  $h_3(x) = \mathbb{1}_{\{x \geq x_2+1\}}$ , the label is  $[0, 0]$ .

However, the label  $[1, 0]$  can't be achieved by any  $h \in \mathcal{H}$ . Then,  $|H(S^X)| = 3 < 2^2$  and we can say  $S^X$  can't be shattered by  $\mathcal{H}$  which implies  $d < 2$ .


**Example 5. Two-dimensional linear classifiers.** Firstly, we consider three data points located at 2-dimensional space, which have the triangle shape. By Figure 1, we can say the dataset generated by three data distributed as Figure 1 can be shattered by  $\mathcal{H}$ , which implies  $d \geq 3$ .

Furthermore, the distribution of 4 points in 2-dimensional space can only have 4 different cases. By Figure 2, we give an counterexample for each of 4 cases to show all the dataset contained 4 data can't be shattered by  $\mathcal{H}$ , which implies  $d < 4$ .

Therefore, we have  $d = 3$ .



**Figure 1:** 8 different labels generated by linear classifier under 3 data in 2-dimensional space.



**Figure 2:** Unattainable labels by linear classifier under 4 different cases of 4 data in 2-dimensional space.

**Example 6. K-dimensional linear classifiers.** We directly give the result without proof here.  $d = K + 1$ . The proof of this result will appear on the next homework.