

Lecture 25: Online Convex Optimization

Lecturer: Kirthi

Scribed by: Xindi Lin, Tony Chang Wang



Disclaimer: These notes are not intended to the usual scrutiny reserved for formal publications. They may be distributed with the permission of the instructor.

In this lecture, we will introduce online convex optimization. We will first introduce two motivating examples, the online linear classification, and the expert problem, and give a **unified framework** for online convex optimization. Then, we will discuss two methods, **Follow the Leader (FTL)**, and **Follow the Regularized Leader (FTRL)**. Finally, we will use several examples to show how to choose the regularizer.

WeChat: cstutorcs

1 Examples and Unified Framework

We will first present two examples to show what is online convex optimization. The online linear classification, and the expert problem.

Example 1 (online linear classification). Let $\Theta = \{\theta \in \mathbb{R}^d : \|\theta\|_2 \leq 1\}$. On each round, the learner chooses some $\theta_t \in \Theta$. Simultaneously, the environment picks an instance $\{x_t, y_t\} \in \mathcal{X} \times \mathcal{Y}$ where the domain $\mathcal{X} \in \mathbb{R}^d, \mathcal{Y} = \{+1, -1\}$. Then, the learner incurs the hinge loss $\ell_t(\theta_t) = \max\{0, 1 - y_t \theta_t^\top x_t\}$. Finally, the learner observes the instance $\{x_t, y_t\}$, and hence knows the loss for all $\theta \in \Theta$. The regret is defined as follows

$$R_T(\pi, \{x_t, y_t\}_{t=1}^T) = \sum_{t=1}^T \ell_t(\theta_t) - \min_{\theta \in \Theta} \sum_{t=1}^T \ell_t(\theta)$$

Example 2 (The Expert Problem). Given K arms, and denote $\Delta^K = \{p \in \mathbb{R}_+^K : p^\top \mathbf{1} = 1\}$. On each round t , the learner chooses some $p_t \in \Delta^K$. Simultaneously, the environment picks a loss vector $\ell_t \in [0, 1]^K$. Then, the learner incurs the loss $p_t^\top \ell_t$. Finally, the learner observes the loss vector ℓ_t , and hence knows the loss for all $p \in \Delta^K$. The regret is defined as follows

$$R_T(\pi, \ell) = \sum_{t=1}^T p_t^\top \ell_t - \min_{a \in [K]} \sum_{t=1}^T \ell_t(a) = \sum_{t=1}^T p_t^\top \ell_t - \min_{p \in \Delta^K} \sum_{t=1}^T p^\top \ell_t$$

where $\min_{p \in \Delta^K} \sum_{t=1}^T p^\top \ell_t = \min_{a \in [K]} \sum_{t=1}^T \ell_t(a)$ is easy to see if we take derivative w.r.t. each coordinates of p in $\sum_{t=1}^T p^\top \ell_t$.

We will now present a unified framework for *online convex optimization*.

Definition 1 (Online convex optimization). Consider the following frame. A learner is given a weight space $\Omega \subset \mathbb{R}^d$. On each round t , the learner chooses a weight vector $w_t \in \Omega$. Simultaneously, the environment chooses a loss function $f_t : w \rightarrow \mathbb{R}$, a mapping from weight space to real line. Then the learner incurs the loss $f_t(w_t)$. Finally, the learner observes the loss function f_t , and hence knows the value of $f_t(w)$ for all $w \in \Omega$.

In the above framework, if (1) the weight space Ω is convex and compact, and (2) the loss function f_t at every round is convex, the framework is called *online convex optimization*.

Given a horizon T . The goal is to minimize the regret against the best-fixed weight vector in Ω w.r.t. the policy π of choosing the weight vector at each round.

$$R_T(\pi, f) = \sum_{t=1}^T f_t(w_t) - \min_{w \in [\Omega]} \sum_{t=1}^T f_t(w)$$

In example 1, the ℓ_2 -ball is convex and compact, and the hinge loss is convex. In 2, Δ^K is convex and compact, and the loss $p_t^T w$ is a linear function of w and thus convex.

2 Follow the Regularized Leader

A most straightforward leader(FTL). The weight w_t is chosen by



$$\arg \min_{w \in \Omega} \sum_{s=1}^{t-1} f_s(w)$$

which is the best weight for the observed loss function. However, this is often a bad idea, as the chosen weight could be forced to round. Therefore, we will stabilize the FTL by adding a regularized term $\Lambda(w)$

$$w_t \in \arg \min_{w \in \Omega} \left\{ \sum_{s=1}^{t-1} f_s(w) + \Lambda(w) \right\}$$

We call the above policy with the regularized term **Follow the Regularized Leader(FTRL)**, and we will give its regret upper bound.

Theorem 3 (Regret Upper Bound for FTRL). *For any $u \in \Omega$, FTRL satisfies*

$$\begin{aligned} R_T(\text{FTRL}, f) &\leq \sum_{t=1}^T f_t(w_t) - \sum_{t=1}^T f_t(u) \\ &\leq \sum_{t=1}^T (f_t(w_t) - f_t(w_{t+1})) + \Lambda(u) - \min_{w \in \Omega} \Lambda(w) \end{aligned}$$

N.B. We have not assumed convexity of Ω , f_t , or Λ in the theorem.

Proof The first inequality is by the definition of regret. For the proof of the second inequality, we denote

$$F_t(w) = \sum_{s=1}^t f_s(w) + \Lambda(w)$$

and let

$$\Phi_t = \min_{w \in \Omega} F_t(w) = F_t(w_{t+1})$$

Consider $\Phi_{t-1} - \Phi_t$, and we have

$$\begin{aligned} \Phi_{t-1} - \Phi_t &= F_{t-1}(w_t) - F_t(w_{t+1}) \\ &= F_{t-1}(w_t) - (F_{t-1}(w_{t+1}) + f_t(w_{t+1})) \\ &= (F_{t-1}(w_t) - F_{t-1}(w_{t+1})) - f_t(w_{t+1}) \\ &\leq -f_t(w_{t+1}) \end{aligned}$$

since $F_{t-1}(w_t) \leq F_{t-1}(w_{t+1})$, Then we will have

$$\Phi_{t-1} - \Phi_t + f_t(w_t) \leq f_t(w_t) - f_t(w_{t+1})$$

by adding $f_t(w_t)$ to both sides of the equation. Then we sum both sides from $t = 1, \dots, T$, and we will have

程序代写代做 CS编程辅导

We can compute the values of Φ_T, Φ_0 as follows:



$$\Phi_T - \Phi_0 = \sum_{t=1}^T f_t(w_t) \leq \sum_{t=1}^T (f_t(w_t) - f_t(w_{t+1})) + \sum_{s=1}^T f_s(u) + \Lambda(u)$$

$$\Phi_0 = \min_{w \in \Omega} \Lambda(w)$$

Therefore, we have

$$\sum_{t=1}^T f_t(w_t) - \sum_{s=1}^T f_s(u) - \Lambda(u) + \min_{w \in \Omega} \Lambda(w) \leq \sum_{t=1}^T (f_t(w_t) - f_t(w_{t+1}))$$

and thus

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

□

Remark:

- The above theorem implies that for follow the leader (FTL),

QQ: 749389476

$$R_T(\text{FTL}, \underline{f}) \leq \sum_{t=1}^T (f_t(w_t) - f_t(w_{t+1})).$$

- If w_t fluctuates frequently, the regret of FTL/FTL will be bad.
- The purpose of the regularized term $\Lambda(w)$ is to stabilize the chosen weight w_t .

https://tutorcs.com

3 Examples Analysis: How a regularizer is Chosen

To motivate how a regularizer is chosen, we will consider 3 examples for FTL with $\Omega = [0, 1]$ and $f_t : [0, 1] \rightarrow [0, 1]$

3.1 Example 1: FTL with linear losses

First, Let $\Omega = [0, 1]$. Then we define $f_t(w) \forall w \in \Omega$:

$$f_t(w) = \begin{cases} \frac{1}{2}w & \text{if } t = 1 \\ w & \text{if } t \text{ is odd, } t > 1 \\ 1 - w & \text{if } t \text{ is even} \end{cases}$$

We have:

$$F_t(w) = \sum_{s=1}^t f_s(w) = \begin{cases} \frac{1}{2}w + \frac{t-1}{2} & \text{if } t \text{ is odd} \\ -\frac{1}{2}w + \frac{t}{2} & \text{if } t \text{ is even} \end{cases}$$

Hence, we have the following:

$$w_t = \arg \min_{w \in [0,1]} F_{t-1}(w) = \begin{cases} 0 & \text{if } t \text{ is even} \\ 1 & \text{if } t \text{ is odd} \end{cases}$$

Therefore, we obtain the Upper Bound from the Thm 3:

$$R_T \leq \sum_{t \text{ s.t } t \text{ is odd}} (1 - 0) + \sum_{t \text{ s.t } t \text{ is even}} (1 - 0) \simeq T.$$

The bound given by the theorem is not tight. However, it is not hard to see that the actual regret is also large. The total loss of FTL is T . The best action in hindsight will have loss at most $\frac{T}{2}$. Therefore, we have **Regret** $\geq \frac{T}{2}$. The Bound on R_T is pretty tight. The linear losses are bad use case for FTL.



3.2 Example 2: FTL with quadratic losses

Let $\Omega = [0, 1]$, and we define $f_t(w), \forall w \in \Omega$ as following:

$$f_t(w) = \begin{cases} w^2 & \text{if } w \text{ is odd} \\ (1-w)^2 & \text{if } w \text{ is even} \end{cases}$$

Similar to the previous example, the best action for a given round oscillates between 0 and 1. However, we will see that the regret is not large.

First note that the sum of losses can be written as:

$$F_t(w) = \begin{cases} \frac{t+1}{2} w^2 + \frac{t-1}{2} (1-w)^2 & \text{if } t \text{ is odd} \\ \frac{t}{2} w^2 + \frac{t}{2} (1-w)^2 & \text{if } t \text{ is even} \end{cases}$$

Hence we have,

$$w_t = \arg \min_{w \in [0,1]} F_{t-1}(w) = \begin{cases} \frac{1}{2} & \text{if } t \text{ is odd} \\ \frac{1}{2} & \text{if } t \text{ is even} \end{cases}$$

We see that the choices made by FTL do not oscillate much, with $w_t \rightarrow \frac{1}{2}$ as $t \rightarrow \infty$. We have the following upper bound:

$$\begin{aligned} R_T &\leq \sum_{t=1}^T f_t(u_t) - f_t(u_{t-1}) \\ &= \sum_{t \text{ s.t } t \text{ is odd}} \left(\left(\frac{1}{2} \right)^2 - \left(\frac{1}{2} - \frac{1}{2t} \right)^2 \right) + \sum_{t \text{ s.t } t \text{ is even}} \left(\left(\frac{1}{2} + \frac{1}{2t} \right)^2 - \left(\frac{1}{2} \right)^2 \right) \\ &= \sum_{t=1}^T \frac{1}{2t} + \mathcal{O} \left(\frac{1}{t^2} \right) \\ &\in \mathcal{O}(\log T) \end{aligned}$$

3.3 Example 3: FTRL with Linear losses

For our final example, we will revisit the linear losses in the first example, but will add a regularizer to stabilize the fluctuations. Since quadratic losses achieved small regret, let us try $\Lambda(w) = \frac{1}{\eta} (w - \frac{1}{2})^2$ (η will be chosen later). We define f_t same as in example 1, namely: $\forall w \in \Omega = [0, 1]$:

$$f_t(w) = \begin{cases} 1/2w & \text{if } t = 1 \\ w & \text{if } t \text{ is odd, } t > 1 \\ 1-w & \text{if } t \text{ is even} \end{cases}$$

Then we have $F_t(w)$:

$$F_t(w) = \sum_{s=1}^t f_s(w) + \Lambda(w) = \begin{cases} \frac{1}{2}w + \frac{t-1}{2} + \frac{\eta}{4}(w-1)^2 & \text{if } t \text{ is odd} \\ \frac{1}{\eta}(w - \frac{1}{2})^2 - \frac{1}{2}w + \frac{t}{2} & \text{if } t \text{ is even} \end{cases}$$

Hence we got:



$$F_{t+1}(w) = \begin{cases} \frac{1}{2} + \frac{\eta}{4} & \text{if } t \text{ is odd} \\ \frac{1}{2} - \frac{\eta}{4} & \text{if } t \text{ is even} \end{cases}$$

Then we have the following regret. Define $B := \max_{w \in [0,1]} \frac{1}{\eta} (w - \frac{1}{2})^2 - \min_{w \in [0,1]} \frac{1}{\eta} (w - \frac{1}{2})^2 = \frac{1}{4\eta}$. We have,

$$\begin{aligned} R_T &\leq \left(\sum_{t=1}^T f_t(w_t) - f_t(w_{t+1}) \right) + B \\ &= \sum_{t \text{ s.t. } t \text{ is odd}} \left(\frac{1}{2} + \frac{\eta}{4} \right) - \sum_{t \text{ s.t. } t \text{ is even}} \left(\frac{1}{2} - \frac{\eta}{4} \right) + B \\ &= \sum_{t=1}^T \frac{\eta}{2} + \frac{1}{4\eta} \end{aligned}$$

Next, we decide to choose $\eta = \frac{1}{\sqrt{T}}$. Based on the regret's UB we just showed, we have:

$$R_T \in \mathcal{O}(\sqrt{T})$$

Some take-aways from the examples above:

- Linear functions have bad behaviour in FTL due to the instability of the chosen w_t
- We should add a "nice" regularizer to stabilize oscillations ("nice" means strong convexity here)

<https://tutorcs.com>