

# 程序代写代做 CS编程辅导

CS861: Theoretical Foundations of Machine Learning

Lecture 15 - 09/10/2023

University of Wisconsin-Madison, Fall 2023

Lecture 15: Nonparametric Density Estimation, Lower bounds for prediction problems

Lecturer: Kirthi Suresh

Scribed by: Haoyue Bai, Zexuan Sun



**Disclaimer:** These notes are subject to the usual scrutiny reserved for formal publications. They may be distributed outside this class only with the permission of the instructor.

In this lecture, we will first present a lower bound for **nonparametric density estimation**, and then study **kernel density estimation** to upper bound the risk. We will conclude with a framework for proving **lower bounds for prediction problems**.

## 1 Nonparametric Density Estimation (Cont'd)

### 1.1 Recap: Nonparametric Density Estimation

Consider the function space  $\mathcal{F} = \{f : [0, 1] \rightarrow [0, B], |f(x_1) - f(x_2)| \leq L|x_1 - x_2|\}$ . The constraint ensures the function is Lipschitz continuous with Lipschitz constant  $L$ . We are interested in the family of distributions  $\mathcal{P}$ , which consists of distributions whose densities are  $L$ -Lipschitz. That is,  $\mathcal{P} = \{p : \text{density } p \text{ of } P \text{ is in } \mathcal{F}\}$ .

We observe samples  $S = \{X_1, \dots, X_n\} \stackrel{\text{iid}}{\sim} p \in \mathcal{P}$ , where the set  $S$  represents a random sample of  $n$  data points that are independent and identically distributed (iid) according to some unknown density  $p$  that belongs to  $\mathcal{P}$ .

Given the sample  $S$ , we wish to estimate the density  $p$  of  $\mathcal{P}$  in the  $L_2$  loss. That is:

$$\Phi \circ \rho(p_1, p_2) = \int (p_1(t) - p_2(t))^2 dt$$

We will show that the minimal risk satisfies:

$$R_n^* = \inf_{\hat{p}} \sup_{p \in \mathcal{F}} \mathbb{E}_S[\|\hat{p} - p\|_2^2] \in \Theta(n^{-\frac{2}{3}}).$$

### 1.2 Lower bound


We will first prove a lower bound via Fano's method.

#### Step 1: Construct alternatives

Consider the function  $\psi$  illustrated in Figure 1. The following facts are straightforward to verify.

- $\psi$  is 1-Lipschitz, meaning that for any two inputs  $\lambda_1$  and  $\lambda_2$ :  $|\psi(\lambda_1) - \psi(\lambda_2)| \leq |\lambda_1 - \lambda_2|$ ;
- $\int \psi = 0$ , which indicates that the area under the curve of the function, over its entire domain, sums up to zero;
- $-\frac{1}{4} \leq \psi \leq \frac{1}{4}$ , which gives the range of the function;
- $\int \psi^2 = \frac{1}{48}$ , which is the squared integral of  $\psi$ .

# 程序代写代做 CS编程辅导



**Figure 1:** Illustrative figure for the function  $\psi(\lambda)$ .

**WeChat: cstutorcs**

To construct the alternatives let  $h$  is a positive number ( $h > 0$ ) that we will decide later. Let  $m = \frac{1}{n}$ . The alternative function space  $\mathcal{F}'$  is:

**Assignment Project Exam Help**

This space defines a set of functions  $P_\omega$  that are formed by a linear combination of basis function  $\Phi_j(t)$ . The vector  $\omega$  resides in some set  $\Omega_m$ . The basic function is defined as:

**Email: tutorcs@163.com**

$$\Phi_j(t) = Lh\Phi\left(\frac{t - (j - \frac{1}{2}) \cdot h}{h}\right),$$

where  $L$  denotes the Lipschitz constant, and  $h$  is the bandwidth. The illustrative example in Figure 2 would provide a visual representation of the approximation.

**QQ: 749389476**

**https://tutorcs.com**




**Figure 2:** Illustrative figure for the example  $p_\omega$ .

## Step 2: Lower bound the distance $\rho(p_\omega, p_{\omega'})$

The objective of this step is to determine a lower bound for the difference between  $p_\omega$  and  $p_{\omega'}$ . We can bound the density below:

# 程序代写代做 CS编程辅导



$$\begin{aligned}
 & (p_\omega - p_{\omega'})^2 \\
 & \int_{\frac{j-1}{m}}^{\frac{j}{m}} (1 + \omega_j \Phi_j - (1 + \omega'_j \Phi_j))^2 \\
 & \mathbb{I}(\omega_j \neq \omega'_j) \int_{\frac{j-1}{m}}^{\frac{j}{m}} \Phi_j^2 \\
 & = \frac{H(\omega, \omega') L^2 h^3}{48}
 \end{aligned}$$

By the Varshamov-Gilbert lemma, we have  $H(\omega, \omega') \geq \frac{m}{8} = \frac{1}{8h}$  as

WeChat: cstutorcs

$$\min_{\omega, \omega'} \|p_\omega - p_{\omega'}\| = \sqrt{\frac{L^2 h^3}{48}}$$

Assignment Project Exam Help

### Step 3: Upper bound KL

In this step, the goal is to determine an upper bound for the Kullback-Leibler (KL) divergence between two functions,  $p_\omega$  and  $p_{\omega'}$ . Based on the definition of KL divergence and expanding KL for  $p_\omega$  and  $p_{\omega'}$ :

Email: tutores@163.com

$$\begin{aligned}
 KL(p_\omega, p_{\omega'}) &= \int_0^1 p_\omega \log\left(\frac{p_\omega}{p_{\omega'}}\right) \\
 &= \sum_{j=1}^m \left( \int_{\frac{j-1}{m}}^{\frac{j}{m}} (1 + \omega_j \Phi_j) \log\left(\frac{1 + \omega_j \Phi_j}{1 + \omega'_j \Phi_j}\right) \mathbb{I}(\omega_j \neq \omega'_j) \right)
 \end{aligned}$$

QQ: 749389476

After some algebra (you will do this in the homework), we have the following upperbound:

https://tutorcs.com

$$KL(p_\omega, p_{\omega'}) \leq H(\omega, \omega') \frac{L^2 h^3}{48} \Rightarrow \forall \omega, \omega', KL(p_\omega, p_{\omega'}) \leq \frac{L^2 h^2}{48}$$

This suggests that regardless of the particular values of  $\omega$  and  $\omega'$ , the KL divergence between any two functions  $p_\omega$  and  $p_{\omega'}$  from the considered set is always less than or equal to  $\frac{L^2 h^2}{48}$ .

A formal proof of this statement is left as an exercise in a homework assignment.

### Step 4: Apply local Fano

Applying local Fano's inequality in this step, we derive conditions and constraints for the estimation problem. We want  $\max_{\omega, \omega'} KL(p_\omega, p_{\omega'}) \leq \frac{\log(\omega)}{4n}$ . This relation is upper bound the maximum KL divergence between any two functions in the set by a term that diminishes with increasing sample size  $n$ .

Sufficient if we have the equation  $\frac{L^2 h^2}{48} \leq \frac{\log(2^{\frac{m}{8}})}{4n} = \frac{\log(2)}{32nh}$ . Choose  $h = C, \frac{1}{n^{\frac{1}{3}} L^{\frac{2}{3}}}$ , which determines the choice of  $h$  as a function of  $n$  and  $L$ . Then, we have the equation:

$$R_n^* \geq \frac{1}{2} \Phi\left(\frac{Lh}{2 \times 8\sqrt{6}}\right) = C \frac{L^{\frac{2}{3}}}{n^{\frac{2}{3}}}.$$

This offers a lower bound on the risk,  $R_n^*$ , which quantifies the error in the estimation. The given requirements ensure the validity and applicability of the above relations:

- $m \geq \frac{1}{h} \geq 8$ . This is necessary for the Varshamov-Gilbert lemma to be applicable.

# 程序代写代做 CS编程辅导

- Cardinality of  $\mathcal{F}'$ :  $|\mathcal{F}'| \geq 16 \Leftrightarrow 2^{\frac{m}{8}} \geq 16 \Leftrightarrow h \leq \frac{1}{32} \Rightarrow$  satisfied if  $n \geq \frac{c}{L^2}$ . This ensures a sufficient number of observations given the Lipschitz constant  $L$ .
- KL bounding condition: imposes the KL divergence between functions remains bounded, and ties the bandwidth  $h$  to the Lipschitz constant  $L$ .



## 2 Upper Bound on Kernel Density Estimation

Kernel Density Estimation (KDE) is a non-parametric technique to estimate the probability density function of a continuous random variable. Kernel Density Estimation (KDE) has the following form:

$$\hat{p}(t) = \frac{1}{n} \sum_{i=1}^n \frac{1}{h} K\left(\frac{t - x_i}{h}\right),$$

where  $\hat{p}(t)$  is the estimated density at point  $t$ ,  $n$  is the number of data points, and  $x_i$  are the observed data points.  $h$  is bandwidth parameters, which plays a critical role in KDE.  $K$  is a (smoothing) kernel with the following properties:

(1) Normalization:

$$\int K(u) du = 1,$$

This ensures the result will integrate to 1 over its entire domain, maintaining the fundamental property of a probability density function.

(2) Symmetry:

$$K(u) = K(-u).$$

This property ensures that the kernel is symmetric around zero. As a result, the estimated density will not be biased towards any direction from the point of estimation.


For the problem at hand, the kernel selected is  $K(t) = \mathbb{I}(|t| \leq \frac{1}{2})$ , which is sufficient for Lipschitz functions. This is a simple uniform kernel.

We can bound the risk as follows:

$$\begin{aligned} \mathbb{E}[\|p - \hat{p}\|_2^2] &= \mathbb{E}\left[\int (p - \hat{p})^2 dt\right] \\ &= \mathbb{E}\left[\int (p - \mathbb{E}(\hat{p}))^2 dt + \int (\mathbb{E}(\hat{p}) - \hat{p})^2 dt + 2 \int (p - \mathbb{E}(\hat{p}))(\mathbb{E}(\hat{p}) - \hat{p}) dt\right] \\ &= \int_0^1 \underbrace{(p(t) - \mathbb{E}(\hat{p}(t)))^2}_{\text{bias}(t)} dt + \int_0^1 \underbrace{\mathbb{E}[(\hat{p}(t) - \mathbb{E}(\hat{p}(t)))^2]}_{\text{Var}(t)} dt + 2 \int_0^1 (p - \mathbb{E}(\hat{p})) \underbrace{\mathbb{E}[\mathbb{E}(\hat{p}) - \hat{p}]}_{=0} dt \end{aligned}$$

The bias and variance terms can be written and bounded below. The bias of an estimator indicates how far on average the estimate is from the true value. In our context, the bias term is derived from:

# 程序代写代做 CS编程辅导



$$\begin{aligned} \text{bias}(t) &= \int K(u) (p(t+uh) - p(t)) du \\ &= \int K(u) (p(t+uh) - p(t)) du \end{aligned}$$

(Apply  $\mathbb{E}[\frac{1}{n} \sum_i Z_i] = \mathbb{E}[Z_i]$ )

The bias depends on the choice of kernel and bandwidth  $h$ . Then we can bound the bias term:

$$\begin{aligned} |\text{bias}(t)| &\leq \int |K(u)| |p(t+uh) - p(t)| du \\ &\leq \int K(u) L |uh| du \\ &= Lh \int K(u) |u| du \\ &= C_1 Lh \end{aligned}$$

where  $C_1 = \int K(u) |u| du$  is a constant that essentially depends on the kernel, and  $L$  is the Lipschitz constant, signifying the bound on the rate of change of our function.

Variance quantifies the dispersion of an estimator around its expected value. In the context of KDE, variance arises from the randomness in the sample. For our setup, the variance term is captured by:

$$\begin{aligned} \text{Var}(t) &= \text{Var}\left(\frac{1}{n} \sum_{i=1}^n \frac{1}{h} K\left(\frac{t - X_i}{h}\right)\right) \\ &= \frac{1}{n} \text{Var}_{X \sim P}\left(\frac{1}{h} K\left(\frac{t - X}{h}\right)\right) \quad (\text{Apply } \text{Var}\left(\frac{1}{n} \sum_{i=1}^n Z_i\right) = \frac{1}{n} \text{Var}(Z_1)) \\ &\leq \frac{1}{n} \mathbb{E}_{X \sim P}\left[\frac{1}{h^2} K^2\left(\frac{t - X}{h}\right)\right] \quad (\text{Var}(Z) = \mathbb{E}[Z^2] - (\mathbb{E}[Z])^2 \leq \mathbb{E}[Z^2]) \\ &= \frac{1}{nh^2} \int K^2\left(\frac{x - t}{h}\right) p(x) dx \\ &= \frac{1}{nh} \int K^2(u) p(t + uh) du \\ &\leq \frac{B}{nh} \int K^2(u) du = \frac{B}{nh} \end{aligned}$$

Where  $B$  is a constant that bounds the product of the squared kernel and the true density. Combine these bounds for bias and variance terms, we have

$$\begin{aligned} \mathbb{E}_S[\|\rho - \hat{\rho}\|_2^2] &\leq \int_0^1 \text{bias}^2(t) dt + \int_0^1 \text{Var}(t) dt \\ &\leq C_1^2 L^2 h^2 + \frac{B}{nh} \\ &= (B + C_1^2) \frac{L^{2/3}}{n^{2/3}} \quad (\text{if } h = \frac{1}{n^{1/3} L^{2/3}}) \end{aligned}$$

# 程序代写代做 CS编程辅导

## 3 Lower Bounds for Prediction Problems

So far, we have talked about the metrics, for instance



- $|\theta_2|, \theta(P) \in \mathbb{R}$
- $\|\theta_2\|_{L_2}, \theta(P) \in \{f : \mathcal{X} \rightarrow \mathbb{R}\}$

Next, we will develop a framework for lower bounds for prediction problems (among others). The framework is comprised of the following elements:

1. Data space  $\mathcal{Z}$ . This is the space from which data samples arise.
2. A family of distribution  $\mathcal{P}$ , where  $\forall P \in \mathcal{P}, \text{supp}(P) \subseteq \mathcal{Z}$ . This represents a collection of probability distributions from which the data can be drawn.
3. A hypothesis/parameter space  $\mathcal{H}$ . This contains all the potential hypotheses or models that we might use to make predictions.
4. An “instance loss”,  $f : \mathcal{H} \times \mathcal{Z} \rightarrow \mathbb{R}$ , where  $f(h, Z)$  is the loss of hypothesis  $h$  on instance  $Z$ . This measures how well a particular hypothesis  $h$  from  $\mathcal{H}$  performs on an instance  $Z$  from  $\mathcal{Z}$ .
5. The *population loss*,  $F(h, P) = \mathbb{E}_{Z \sim P}[f(h, Z)]$ ; *Excess population loss*,  $L(h, P) = F(h, P) - \inf_{h' \in \mathcal{H}} F(h', P)$ . These terms indicate how well our hypothesis does on average (under distribution  $P$ ) and how it compares to the best possible hypothesis in  $\mathcal{H}$ , respectively.
6. A dataset  $S$  is drawn from some  $P \in \mathcal{P}$ .
7. An *estimator*  $\hat{h}$ , which maps the dataset  $S$  to a hypothesis in  $\mathcal{H}$ . Note that we overload notation here for  $\hat{h}$

QQ: 749389476

$h$  : a map from data to  $\mathcal{H}$  (estimator)

$\hat{h}$  : as the estimate ( $\hat{h} \in \mathcal{H}$ )

8. Risk of estimator  $R(\hat{h}, P)$

$$\begin{aligned}
 R(\hat{h}, P) &= \mathbb{E}[L(\hat{h}(S), P)] \\
 &= \mathbb{E}[F(\hat{h}(S), P)] - \inf_{h \in \mathcal{H}} F(h, P)
 \end{aligned}$$

9. Minimax risk:

$$R^* = \inf_{\hat{h}} \sup_{P \in \mathcal{P}} R(\hat{h}, P)$$

Next, let us see an example.

Example 1. Excess risk in classification/regression

- Data space  $\mathcal{Z} = \mathcal{X} \times \mathcal{Y}$ ,  $\mathcal{H} : \{h : \mathcal{X} \rightarrow \mathcal{Y}\}$ . This is the set of all possible prediction functions mapping from features  $\mathcal{X}$  to outcomes  $\mathcal{Y}$ .
- Instance loss (in classification),  $f(h, (X, Y)) = \mathbb{1}(h(X) \neq Y)$ . This measures the discrepancy between a predicted and actual class label.
- $F(h, P) = \mathbb{E}_{X, Y \sim P}[\mathbb{1}(h(X) \neq Y)]$  is the “risk”. This is the expected value of the instance loss over the joint distribution of  $\mathcal{X}$  and  $\mathcal{Y}$ , and can be understood as the overall error rate of the classifier.

# 程序代写代做 CS编程辅导

- $L(h, P) = F(h, P) - F(h^*, P)$ , where  $h^*$  is the Bayes optimal classifier, i.e.  $h^* = \arg \max_{y \in \mathcal{Y}} \mathbb{P}(Y = y|X = x)$ , and  $L(h, P)$  is the “excess risk” in classification. The excess risk quantifies how much worse our classifier  $h$  performs compared to the Bayes optimal classifier  $h^*$ .
- In regression we define  $L(h, P) = \mathbb{E}_{X \sim P} [(h(X) - Y)^2]$ . The typical loss function used is the squared loss.
- $L(h, P) = \mathbb{E}_{X \sim P} [(h(X) - \mathbb{E}[Y|X])^2]$ , where  $h^*(x) = \mathbb{E}[Y|X = x]$ ,  $L(h, P)$  is the “excess risk” in regression. It measures how much worse our regression function performs compared to the conditional expectation  $h^*(x)$  which is the conditional expectation.
- Note that this is also  $L(h_1, P) - L(h_2, P) = \|h_1 - h_2\|_2^2$ , which captures the squared difference between two hypotheses  $h_1$  and  $h_2$ , and  $\mathbb{E}_S[\Phi \circ \rho(\hat{h}, h^*)]$  measures how much estimator  $\hat{h}$  deviates from the optimal hypothesis  $h^*$ .

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>