

程序代写代做 CS编程辅导

Copyright © Copyright University of New South Wales 2022. All rights reserved.

Course materials subject to Copyright

UNSW Sydney owns copyright in these materials (including any associated content). The material is subject to copyright under Australian law and overseas under international treaties.

The materials are provided for use by enrolled UNSW students. These materials, or any part, may not be copied, shared or distributed, in print or digitally, outside the course without permission.

Students may only copy a reasonable portion of the materials for personal research or study or for criticism or review. Under no circumstances may these materials be copied or reproduced for sale or commercial purposes without prior written permission of UNSW Sydney.

Statement on class recording

To ensure the free and open discussion of ideas, students may not record, by any means, classroom lectures, discussion and/or activities without the advance written permission of the instructor, and any such recording properly approved in advance can be used solely for the student's own private use.

WARNING: Your failure to comply with these conditions may lead to disciplinary action and may give rise to a civil action or a criminal offence under the law.

THE ABOVE INFORMATION MUST NOT BE REMOVED FROM THIS MATERIAL.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



程序代写代做 CS编程辅导

ECON3



Review of CLT & LLN

Dr. Rachida Ouysse
WeChat: cstutorcs

School of Economics
UNSW
Assignment Project Exam Help

©Copyright University of New South Wales 2020. All rights reserved. This copyright notice must not be removed from this material

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

- LLN and CLT – pillars of all statistics

- Let $\{Z_1, Z_2, \dots, Z_T\}$ be a set of *independent* RVs with common mean μ and variance σ^2 .

- **Law of large numbers:** the probability that $\bar{Z} = \frac{1}{T} \sum_{t=1}^T Z_t$ is from μ converges to zero as T goes to infinity

- **Central limit theorem:** the distributions of $\frac{\bar{Z} - \mu}{\sqrt{\text{Var}(\bar{Z})}}$ converges to $N(0,1)$ as T goes to infinity.

- Note that $\text{Var}(Z) = \sigma^2/T$ (see Rule 8).

What happens if $\{Z_1, Z_2, \dots, Z_T\}$ are correlated?

程序代写代做 CS编程辅导



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Distribution of Sample Mean

程序代写代做 CS编程辅导



Consider N random variables X_1, \dots, X_N .

- Let's consider $\bar{X} = \frac{1}{N} \sum_{k=1}^N X_k$.
- \bar{X} is called the “sample mean” or the “empirical mean”.
- \bar{X} is a random variable.

WeChat: cstutorcs

Suppose we observe values for X_1, \dots, X_N and calculate the empirical mean of the observed values. That gives us one value for \bar{X} . But the value of \bar{X} changes depending on the observed values.

Assignment Project Exam Help

Email: tutorcs@163.com

- Suppose we toss a fair coin $N = 5$ times and get H, H, H, T, T . Let $X_k = 1$ when come up heads. Then $\frac{1}{N} \sum_{k=1}^N X_k = \frac{3}{5}$.
- Suppose we toss the coin another $N = 5$ times and get T, T, H, T, H . Now $\frac{1}{N} \sum_{k=1}^N X_k = \frac{2}{5}$.

QQ: 749389476


https://tutorcs.com

Distribution of Sample Mean

程序代写代做 CS编程辅导

Toss fair coin $N = 5$ times. Calculate $\sum_{k=1}^N X_k$. Repeat, and plot histogram of values. It is $\text{Bin}(N, \frac{1}{2})$.

- `X=[];for i=1:1000 X=[X sum(rand(1,5)<0.5)]]; end; hist(X,5)`



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Distribution of Sample Mean

程序代写代做 CS编程辅导



Random variable $\bar{X} =$

- Suppose the X_k are **independent and identically distributed (i.i.d)**
- Each X_k has mean $E(X_k) = \mu$ and variance $Var(X_k) = \sigma^2$.

Then we can calculate the mean of \bar{X} as:

$$E(\bar{X}) = E\left(\frac{1}{N} \sum_{k=1}^N X_k\right) = \frac{1}{N} \sum_{k=1}^N E(X_k) = \mu$$


NB: recall linearity of expectation: $E(X + Y) = E(X) + E(Y)$ and $E(aX) = aE[X]$

- We say \bar{X} is an **unbiased estimator** of μ since $E[\bar{X}] = \mu$

Distribution of Sample Mean

程序代写代做 CS编程辅导

We can calculate the variance of \bar{X} as:


$$\begin{aligned} \text{var}(\bar{X}) &= \text{var}\left(\frac{1}{N} \sum_{k=1}^N X_k\right) = \frac{1}{N^2} \text{var}\left(\sum_{k=1}^N X_k\right) \\ &= \frac{1}{N^2} \sum_{k=1}^N \text{var}(X_k) = \frac{N\sigma^2}{N^2} = \frac{\sigma^2}{N} \end{aligned}$$

WeChat: tutormcs

Assignment Project Exam Help

NB: recall $\text{Var}(aX) = a^2 \text{Var}(X)$ and $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y)$ when X, Y independent.

Email: tutormcs@163.com

- As N increases, the variance of \bar{X} falls.
- $\text{Var}(NX) = N^2 \text{Var}(X)$ for random variable X .
- But when add together independent random variables $X_1 + X_2 + \dots$ the variance is only $N\text{Var}(X)$ rather than $N^2 \text{Var}(X)$
- This is due to **statistical multiplexing**. Small and large values of X_i tend to cancel out for large N .

QQ: 749389476

<https://tutormcs.com>

Weak Law of Large Numbers¹

程序代写代做 CS编程辅导



Consider N independent and identically distributed (i.i.d) random variables X_1, \dots, X_N each with mean μ and variance σ^2 . Let $\bar{X} = \frac{1}{N} \sum_{k=1}^N X_k$. For any $\epsilon > 0$:

$$P(|\bar{X} - \mu| \geq \epsilon) \rightarrow 0 \text{ as } N \rightarrow \infty$$

That is, \bar{X} **concentrates** around the mean μ as N increases.

Proof:

- $E(\bar{X}) = E(\frac{1}{N} \sum_{k=1}^N X_k) = \frac{1}{N} \sum_{k=1}^N E(X_k) = \mu$
- $\text{var}(\bar{X}) = \text{var}(\frac{1}{N} \sum_{k=1}^N X_k) = \frac{1}{N^2} \sum_{k=1}^N \text{var}(X_k) = \frac{\sigma^2}{N}$
- By Chebyshev's inequality, $P(|\bar{X} - \mu| \geq \epsilon) \leq \frac{\sigma^2}{N\epsilon^2}$

¹There is also a **strong law of large numbers**, but we won't deal with that here.

Who cares ?

程序代写代做 CS编程辅导

- Suppose we have
- Define indicator random variable X_i equal to 1 when event E is observed in trial i otherwise
- Recall $E[X_i] = P(E)$ is the probability that event E occurs.
- $\bar{X} = \frac{1}{N} \sum_{k=1}^N X_k$ is then the relative frequency with which event E is observed over N experiments.
- And ...
 $P(|\bar{X} - \mu| \geq \epsilon) \rightarrow 0$ as $N \rightarrow \infty$
tells us that this observed relative frequency \bar{X} converges to the probability $P(E)$ of event E as N grows large.
- So the law of large numbers formalises the intuition of probability as frequency when an experiment can be repeated many times. But probability still makes sense even if cannot repeat an experiment many times – all our analysis still holds.



WeChat: tutores

Assignment Project Exam Help


Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Central Limit Theorem (CLT)

Histogram of $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$ as N increases, but now we normalise to keep the area under the curve fixed:



- See that (i) curve narrows as n increases, it concentrates as we already know from weak law of large numbers.
- Curve becomes more “bell-shaped” as N increases – this is the CLT


The Normal (or Gaussian) Distribution

程序代写代做 CS编程辅导

Define the following fu




$$\frac{1}{\sigma\sqrt{2\pi}}e^{-\frac{(y-\mu)^2}{2\sigma^2}}$$



Central Limit Theorem (CLT)

Overlay the Normal distribution, with parameter μ equal to the mean and σ^2 equal to the variance of the measured histograms:



- CLT says that as N increases the distribution of \bar{X} converges to a Normal (or Gaussian) distribution.
- Variance $\rightarrow 0$ as $N \rightarrow \infty$, i.e. distribution concentrates around its mean as N increases.