

程序代写代做 CS编程辅导

Copyright ©Copyright University of New South Wales 2020. All rights reserved.

Course materials subject to Copyright

UNSW Sydney owns copyright in these materials. The material is subject to copyright under Australian law and overseas under international treaties. The



otherwise). The material is subject to copyright under Australian law and overseas under international treaties. The

materials are provided for use by enrolled UNSW students. These materials, or any part, may not be copied, shared or distributed, in print or digitally, outside the course without permission. Students may only copy a reasonable portion of the material for personal research or study or for criticism or review. Under no circumstances may these materials be copied or reproduced for sale or commercial purposes without prior written permission of UNSW Sydney.

Statement on class recording

To ensure the free and open discussion of ideas, students may not record, by any means, classroom lectures, discussion and/or activities without the advance written permission of the instructor, and any such recording properly approved in advance can be used solely for the student's own private use.

WARNING: Your failure to comply with these conditions may lead to disciplinary action and may give rise to a civil action or a criminal offence under the law.

THE ABOVE INFORMATION MUST NOT BE REMOVED FROM THIS MATERIAL.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutors@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

Slides-10: Modeling

AR



Econometrics

Utility: Testing/Estimating/Forecasting

Introduction to GARCH

WeChat: cstutorcs

Dr. Rachida Ouyse

School of Economics¹

Assignment Project Exam Help

¹©Copyright University of New South Wales 2020. All rights reserved. This copyright notice must not be removed from this material.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



Lecture Plan

程序代写代做 CS编程辅导



- ARCH LM-test
 - Forecasting with ARCH
 - Generalised ARCH: why and how
 - Formulation of GARCH: parameter restrictions
 - Properties of GARCH
 - Mean, variance, ARMA(1,1) representation
 - ML estimation of GARCH
 - Forecasting with GARCH
- WeChat: cstutorcs
- Assignment Project Exam Help
- Email: tutors@163.com
- QQ: 749389476
- <https://tutorcs.com>

ARCH-LM TEST

LM test for ARCH effect

程序代写代做 CS编程辅导

Testing for ARCH



the ARCH LM test

- ▶ Obtain the residuals $\hat{\mu}_t$ from a regression, e.g.

$$y_t = \beta_1 + \beta_2 x_{2t} + \beta_3 x_{3t} + \beta_4 x_{4t} + \mu_t$$

- ▶ Obtain R^2 of the auxiliary regression

$$\hat{\mu}_t^2 = \gamma_0 + \gamma_1 \hat{\mu}_{t-1}^2 + \gamma_2 \hat{\mu}_{t-2}^2 + \dots + \gamma_q \hat{\mu}_{t-q}^2 + \nu_t$$

- ▶ Calculate test statistic $T'R^2$ with T' the number of observations in the auxiliary regression.
- ▶ Under the null hypothesis of no ARCH, $T'R^2 \sim \chi^2(q)$.

ARCH-LM TEST

LM test for ARCH effect: Example

程序代写代做 CS编程辅导



eg. NYSE composite returns

- 1 Estimate the model for mean (eg. AR(1)) and save the residual series $\hat{\mu}_t$.
- 2 OLS auxiliary regression: $\hat{\mu}_t^2 = \gamma_0 + \gamma_1 \hat{\mu}_{t-1}^2 + \cdots + \gamma_q \hat{\mu}_{t-q}^2 + error_t$
Save the R^2 . (q depends on T and data frequency)
- 3 $T' = T - q$, with $q = 5$ reject when $T R^2$ exceeds $\chi^2_{(5)}$

WeChat: estutories

Assignment Project Exam Help

eg. NYSE composite returns LM test with $q=5$

Email: tutors@163.com

E			V		
ARCH Test:			ARCH Test:		
F-statistic	37.43273	Probability	0.000000	F-statistic	1.318328
Obs*R-squared	171.0570	Probability	0.000000	Obs*R-squared	6.589613
				Probability	0.253338
					0.252993

QQ: 749389476

https://tutors.com

Performed on "V" to check the adequacy of variance equation

Forecasting with ARCH Models

程序代写代做 CS编程辅导



- Using repeated substitution we can make multi-step forecasts for the return and its volatility
- Example. AR(1)-ARCH(2)

WeChat: cstutorcs

$$y_t = c + \phi_1 y_{t-1} + \mu_t, \mu_t | \Omega_{t-1} \sim N(0, \sigma_t^2)$$

$$\sigma_t^2 = \alpha_0 + \alpha_1 \mu_{t-1}^2 + \alpha_2 \mu_{t-2}^2$$

Assignment Project Exam Help

$$y_{t+1|t} = c + \phi_1 y_t$$

Email: tutormcs@163.com

$$y_{t+2|t} = c + \phi_1 y_{t+1|t}, \dots$$

QQ: 749389476

$$\sigma_{t+1|t}^2 = \alpha_0 + \alpha_1 \mu_t^2 + \alpha_2 \mu_{t-1}^2,$$


https://tutormcs.com

$$\sigma_{t+2|t}^2 = \alpha_0 + \alpha_1 \mu_{t+1|t}^2 + \alpha_2 \mu_t^2,$$

$$\sigma_{t+3|t}^2 = \alpha_0 + \alpha_1 \sigma_{t+2|t}^2 + \alpha_2 \sigma_{t+1|t}^2, \dots$$



AR(1)-ARCH(5) forecasts
revert to unconditionals
(mean reverting)




WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

OO: 749389476

<https://tutors.com>



Remember the limitations of ARCH!

程序代写代做 CS编程辅导



Advantages of ARCH

- It is able to capture 'clustering' in return series or the autocorrelation in squared returns.
- It facilitates volatility forecasting.
- It explains, partially, non-normality in return series.

WeChat: cstutorcs

Assignment Project Exam Help

Limitations of ARCH

- In ARCH(q), the q may be selected by AIC, SIC or LP test. The correct value of q might be very large. The model might not be parsimonious. (eg. ARCH(1) would not work for the composite return).
- The conditional variance σ_t^2 cannot be negative: Requires non-negativity constraints on the coefficients. Sufficient (but not necessary) condition is: $\alpha_i \geq 0$ for all $i = 0, 1, 2, \dots, q$. Especially for large values of q this might be violated.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

GARCH Models: Introduction

程序代写代做 CS编程辅导



Generalised ARCH (GARCH) models allow the conditional variance to depend upon previous volatility.

- Let μ_t be the error term or shock in a model.

$$\text{ARCH}(q): \text{Var}(\mu_t | \Omega_{t-1}) = \sigma_t^2$$

$$\sigma_t^2 = \alpha_0 + \alpha_1 \mu_{t-1}^2 + \alpha_2 \mu_{t-2}^2 + \cdots + \alpha_q \mu_{t-q}^2,$$

is not parsimonious as a large q is often required.

- If σ_{t-1}^2 is a summary of volatility info in Ω_{t-2} , then $\Omega_{t-1} = \{\mu_{t-1}, \mu_{t-2}, \mu_{t-3}, \dots\} = \{\mu_{t-1}, \Omega_{t-2}\} \approx \{\mu_{t-1}, \sigma_{t-1}^2\}$ (volatility wise!)
- This leads to the GARCH(1,1) model:

$$\sigma_t^2 = \alpha_0 + \alpha_1 \mu_{t-1}^2 + \beta_1 \sigma_{t-1}^2$$

GARCH: Introduction

程序代写代做 CS编程辅导



- More generally, GARCH(p, q) model

$$\text{Var}(\mu_t | \Omega_{t-1}) = \sigma_t^2, \text{WeChat: cstutorcs}$$

$$\sigma_t^2 = \alpha_0 + \alpha_1 \mu_{t-1}^2 + \cdots + \alpha_p \mu_{t-p}^2 + \beta_1 \sigma_{t-1}^2 + \cdots + \beta_p \sigma_{t-p}^2,$$

where the parameters should satisfy:

- (1) Positivity constraint: $\alpha_0 > 0, \alpha_i \geq 0, \beta_j \geq 0$ for all $i = 1, \dots, q$ and $j = 1, \dots, p$
- (2) Finite Variance $\sum_{i=1}^q \alpha_i + \sum_{j=1}^p \beta_j < 1$

- In practice, the models for asset returns rarely go beyond GARCH(1,1).

Properties of GARCH

程序代写代做 CS编程辅导



The generalisation implied by GARCH can be seen from backward iterating the GARCH(1,1) model: **WeChat: cstutorcs**

$$\sigma_t^2 = \frac{\alpha_0}{1 - \beta_1} + \alpha_1 \sum_{j=1}^{\infty} \beta_1^j \mu_{t-j}^2.$$

Assignment Project Exam Help
Email: tutorscs@163.com

This shows that the GARCH model is an ARCH(∞) with geometrically declining coefficients (for $|\beta_1| < 1$): **QQ: 749389476**

<https://tutorscs.com>

Properties of GARCH(1,1)

程序代写代做 CS编程辅导



Alternatively, if we define the surprise in the squared innovations as $\omega_t = \mu_t^2 - \sigma_t^2$, the GARCH(1,1) model can be rewritten as

$$\begin{aligned}\mu_t^2 - \omega_t &= \alpha_0 + \alpha_1 \mu_{t-1}^2 + \beta_1 (\mu_{t-1}^2 - \omega_{t-1}) \\ \mu_t^2 &= \alpha_0 + (\alpha_1 + \beta_1) \mu_{t-1}^2 + \omega_t - \beta_1 \omega_{t-1}\end{aligned}$$

which shows that the squared errors follow an ARMA(1,1) model. As the root of the autoregressive part is $\alpha_1 + \beta_1$, the squared residuals are stationary provided $|\alpha_1 + \beta_1| < 1$.

Under stationarity, $E(\mu_t^2) = E(\mu_{t-1}^2) = E(\sigma_{t-1}^2) = \sigma^2$, the unconditional variance of μ_t is given by

$$\begin{aligned}\sigma^2 &= \alpha_0 + \alpha_1 \sigma^2 + \beta_1 \sigma^2 \\ &= \frac{\alpha_0}{1 - (\alpha_1 + \beta_1)}\end{aligned}$$

Properties of GARCH(1,1)

程序代写代做 CS编程辅导



Two general cases distinguished

- ▶ $\alpha_1 + \beta_1 < 1$ Conditional variance is defined, i.e. finite
- ▶ $\alpha_1 + \beta_1 \geq 1$ → the unconditional variance is not defined, i.e. infinite

WeChat: cstutorcs

Assignment Project Exam Help

The latter case is denoted **non-stationarity in variance**

Email: tutorcs@163.com

- ▶ Variance does not converge to an unconditional mean
- ▶ The special case where $\alpha_1 + \beta_1 = 1$ is known as a unit root in variance or **integrated GARCH** (IGARCH)

QQ: 749389476

<https://tutorcs.com>

Properties of GARCH(1,1)

程序代写代做 CS编程辅导



- GARCH(1,1): $\mu_t | \Omega_{t-1} \sim \mathcal{N}(\mu_t, \sigma_t^2)$,

$$\sigma_t^2 = \alpha_0 + \alpha_1 \mu_{t-1}^2 + \beta_1 \sigma_{t-1}^2$$

$$\alpha_0 > 0, \alpha_1 \geq 0, \beta_1 \geq 0, \alpha_1 + \beta_1 < 1$$

- Its conditional variance is time varying.

$$E(\mu_t | \Omega_{t-1}) = 0, \text{Var}(\mu_t | \Omega_{t-1}) = \sigma_t^2,$$

$$\text{CI}(95\%) = E(y_{t+1} | \Omega_{t-1}) \pm 2\sigma_t$$

- μ_t is a White Noise: $E(\mu_t) = 0$, $\text{Var}(\mu_t) = \frac{\alpha_0}{1 - (\alpha_1 + \beta_1)}$, $\text{Cov}(\mu_t, \mu_{t-j}) = 0$
- But it is NOT an independent WN or iid WN. It is NOT unconditionally Normally distributed: $\text{kurt}(\mu_t) > 3$

Properties of GARCH(1,1)

程序代写代做 CS编程辅导



- GARCH(1,1) can be expressed in terms of **standardised shocks** ν_t :

$$\mu_t = \sigma_t \nu_t \text{ and } \nu_t \sim \text{iid } N(0, 1)$$

- When model is correct, ν_t^2 should have no autocorrelation.

Advantages of the GARCH model (compared to ARCH)

- Avoids overfitting, i.e. a higher order ARCH model may have a more parsimonious GARCH representation
- Due to less estimated parameters, violations of the non-negativity constraint are less likely

<https://tutorcs.com>

GARCH(1,1) Estimation

程序代写代做 CS编程辅导



Estimating GARCH model

For instance, estimate the following AR(1)-GARCH(1,1) model

$$y_t = \mu + \phi(y_{t-1} - \mu)$$

$$\mu_t = \nu_t \sigma_t \quad \nu_t \sim N(0, 1)$$

$$\sigma_t^2 = \alpha_0 + \alpha_1 y_{t-1}^2 + \beta_1 \sigma_{t-1}^2$$

WeChat: cstutorcs

Assignment Project Exam Help

OLS is inappropriate

- ▶ OLS minimises the RSS, $\sum \mu_t^2 = \sum (y_t - \mu + \phi y_{t-1})^2$, which is a function of the parameters in the conditional mean equation only and not in the conditional variance equation
- ▶ In fact, OLS assumes that the residuals are homoscedastic, i.e. all slope coefficients in the conditional variance equation are set to zero

Email: tutors@163.com

QQ: 749389476

https://tutorcs.com

GARCH(1,1) Estimation

程序代写代做 CS编程辅导



Maximum Likelihood

- ▶ Make assumptions about the distribution of μ_t , e.g.

$$\nu_t \sim N(0, 1) \text{ such that } \mu_t \sim N(0, \sigma_t^2)$$

This means that conditional on information available at $t-1$, μ_t is normally distributed with mean zero and variance σ_t^2 with the latter being known at time $t-1$. Note that this does not imply that the **unconditional distribution** of μ_t is normal, as σ_t becomes a random variable if we do not condition on all information available on $t-1$.

- ▶ The conditional distribution of y_t is then also normal, given by

$$f(y_t | y_{t-1}, \dots, \mu_{t-1}, \dots) = \frac{1}{\sqrt{2\pi\sigma_t^2}} \exp\left(-\frac{1}{2} \frac{\mu_t^2}{\sigma_t^2}\right)$$

with $\mu_t = y_t - \mu - \phi y_{t-1}$ and $\sigma_t^2 = \alpha_0 + \alpha_1 \mu_{t-1}^2 + \beta_1 \sigma_{t-1}^2$.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorscs@163.com

QQ: 749389476

https://tutorcs.com

GARCH(1,1) Estimation

程序代写代做 CS编程辅导



- ▶ The **loglikelihood function** is the sum over all t of the log of the conditional density of y_t

$$L = -\frac{T}{2} \log(2\pi) - \frac{T}{2} \sum_{t=1}^T \log(\sigma_t^2) - \frac{1}{2} \sum_{t=1}^T \frac{\mu_t^2}{\sigma_t^2}$$

- ▶ The ML estimator is obtained by **maximising** the loglikelihood with respect to the unknown parameters $(\mu, \alpha_0, \alpha_1, \beta_1)$
- ▶ Analytical solution not possible: use **numerical procedures**

- ▶ These algorithms 'search' over the parameter space, from an **initial guess**, until a maximum for the loglikelihood function is found
- ▶ Potential problem: the loglikelihood function may have several **local maxima** such that alternative initial guesses may yield different results.
- ▶ In practice: use linear regression to get initial estimates of the parameters in the conditional mean equation and choose some (alternative) parameter value for the parameters in the conditional variance equation $\neq 0$.

WeChat: cstutors

Assignment Project Exam Help

Email: tutors@163.com

QQ: 749389476

https://tutors.com

GARCH(1,1) Estimation 程序代写代做 CS编程辅导



GARCH(1,1) Estimation

程序代写代做 CS编程辅导



- ▶ Fortunately, first order moments are, under some weak assumptions, valid even if ϵ_t is not normally distributed.
 - ▶ The parameter estimates are still consistent
 - ▶ Adjustments have to be made to the standard errors, i.e. use Bollerslev-Woodward large variance-covariance matrix, also known as **Quasi Maximum Likelihood Estimation**, which is robust for non-normality.

WeChat: estutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 1

Example 1: GARCH(1,1) Estimation

程序代写代做 CS编程辅导

— ML Estimation (1,1)

eg. NYSE comp



Dependent Variable: RC
Method: ML - ARCH (Marquardt) - Normal distribution

Sample (adjusted): 3 1931
Included observations: 1929 after adjustments
Convergence achieved after 14 iterations
Bollerslev-Wooldridge robust standard errors & covariance
Variance backcast: OFF
GARCH = C(3) + C(4)*RESID(-1)^2 + C(5)*GARCH(-1)

	Coefficient	Std. Error	z-Statistic	Prob.
C	0.072383	0.018494	3.91396	0.0001
AR(1)	0.102468	0.025514	4.016133	0.0001

Variance Equation				
C	0.013305	0.004529	2.93775	0.0033
RESID(-1)^2	0.119181	0.024748	4.81583	0.0000
GARCH(-1)	0.877098	0.021027	41.71362	0.0000

R-squared	0.002416	Mean dependent var	0.035168
Adjusted R-squared	0.000342	S.D. dependent var	1.005452
S.E. of regression	1.005280	Akaike info criterion	3.521965
Sum squared resid	1949.240	Schwarz criterion	3.545689
Log likelihood	-2523.598	F-statistic	1.185523
Durbin-Watson stat	2.055220	Prob(F-statistic)	0.324436

Inverted AR Roots	10
-------------------	----

ARCH Test: q = 5

F-statistic	0.764181	Probability	0.575599
Obs*R-squared	3.825239	Probability	0.574842

Test Equation:
Dependent Variable: STD_RESID*2

Correlogram of Standardized Residuals Squared

Sample: 3 1931
Included observations: 1929
Q-statistic probabilities adjusted for 1 ARMA term(s)

Autocorrelation	Partial Correlation	AC	PAC	Q-Stat	Prob
1	1	-0.001	-0.001	0.0005	
2	0.036	0.036	2.4991	0.114	
3	-0.014	-0.014	2.8675	0.238	
4	-0.011	-0.012	3.1058	0.376	
5	-0.020	-0.019	3.8964	0.420	
6	0.009	0.010	4.0531	0.542	
7	0.013	0.014	4.0658	0.668	
8	0.010	0.010	4.0670	0.772	
9	-0.011	-0.012	4.3009	0.829	
10	-0.010	-0.010	4.4826	0.877	
11	-0.008	-0.006	4.5929	0.917	
12	-0.016	-0.016	5.1005	0.926	
13	-0.004	-0.004	5.1268	0.954	
14	-0.011	-0.011	5.3589	0.966	
15	-0.019	-0.019	6.0452	0.965	
16	-0.009	-0.009	6.2090	0.976	


WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

https://tutorcs.com



Series: Standardized Residuals Sample 3 1931 Observations 1929	
Mean	-0.048341
Median	-0.039867
Maximum	2.850528
Minimum	-6.601836
Std. Dev.	0.996820
Skewness	-0.547486
Kurtosis	4.973199
Jarque-Bera	409.3080
Probability	0.000000

Example 1

Example 1: GARCH(1,1) Estimation

程序代写代做 CS编程辅导



– ML Estimation of GARCH(1,1)

eg. NYSE closing returns (continued)

Large β_1 estimate: about 0.9Small α_1 estimate: about 0.1 $\alpha_1 + \beta_1$ estimate: very close to 1

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutors@163.com

GARCH(1,1) is preferred by AIC and SIC.

	AIC	SIC
AR(1)-ARCH(5)	2.664	2.687
AR(1)-GARCH(1,1)	2.622	2.636

QQ: 749389476

https://tutorcs.com

Example 1

Example 1: GARCH(1,1) Estimation

程序代写代做 CS编程辅导



– ML Estimation of GARCH(1,1)

eg. NYSE closing returns (continued)

GARCH(1,1) σ_t plot is smoother than ARCH(5).Large β_1 estimate implies **persistence**: σ_t tends to continue at the current level.

Assignment Project Exam Help

$$\sigma_t^2 = \alpha_0 + \alpha_1 \varepsilon_{t-1}^2 + \beta_1 \sigma_{t-1}^2$$

Email: tutorcs@163.com

QQ: 749389476

<http://tutorcs.com>

Summary facts about GARCH models



- GARCH(1,1) is usually preferred to ARCH or higher order GARCH, because of its parsimony.
- Usually, GARCH β_1 estimate is about 0.9 or more and $\alpha_1 + \beta_1$ estimate is very close to 1, for daily returns.
- Standardised residuals are usually non-normal, with negative skewness and excessive kurtosis.
- GARCH(1,1) is able to capture clustering in returns but unable to account for
Asymmetry: negative returns tend to cause more volatility;
Non-normality; Structural change
- Coefficient restrictions are hard to impose in MLE