

程序代写代做 CS编程辅导

Copyright © Copyright University of New South Wales 2020. All rights reserved.



Course materials copyright

UNSW Sydney owns copyright in the materials (unless stated otherwise). The material is subject to copyright under Australian law and under international treaties. The materials are provided for use by enrolled UNSW students only. No part, or any part, may not be copied, shared or distributed, in print or digitally, outside the course without permission. Students may only copy a reasonable portion of the material for personal research or study or for criticism or review. Under no circumstances may these materials be copied or reproduced for sale or commercial purposes without prior written permission of UNSW Sydney.

WeChat: cstutorcs

Statement on class recording

To ensure the free and open discussion of ideas, students may not record by any means, classroom lectures, discussion and/or activities without the advance written permission of the instructor, and any such recording properly approved in advance can be used solely for the students own private use.

WARNING: Your failure to comply with these conditions may lead to disciplinary action, and may give rise to a civil action or a criminal offence under the law.

THE ABOVE INFORMATION MUST NOT BE REMOVED FROM THIS MATERIAL.

QQ: 749389476

<https://tutorcs.com>

University of New South Wales School of Economics

Financial Econometrics

Tutorial 6



1. (ARCH model)

Consider the following model,

$$y_t = c + \phi_1 y_{t-1} + \varepsilon_t, \quad |\phi_1| < 1, \quad \varepsilon_t | \Omega_{t-1} \sim N(0, \sigma_t^2),$$

$$\sigma_t^2 = \alpha_0 + \alpha_1 \varepsilon_{t-1}^2 + \alpha_2 \varepsilon_{t-2}^2, \quad \alpha_0 > 0, \quad \alpha_1 \geq 0, \quad \alpha_2 \geq 0, \quad \alpha_1 + \alpha_2 < 1,$$

where Ω_t is the information set at the end of period t .

- Find $E(\varepsilon_t | \Omega_{t-1})$, $E(y_t | \Omega_{t-1})$ and their unconditional counterparts.
- Find $\text{Var}(\varepsilon_t | \Omega_{t-1})$, $\text{Var}(y_t | \Omega_{t-1})$ and their unconditional counterparts.
- Is ε_t a white noise process? Is it an independent (or iid) WN process? Verify your answer.
- In what fundamental way does the ARCH model differ from the standard (homoscedastic) ARMA models? What is the purpose of the variance equation?
- Ceteris paribus, what is the change in σ_t^2 caused by a one-unit change in ε_{t-1}^2 ?
- Suppose $\alpha_2 = 0$, $\alpha_1^2 < 1/3$ and ε_t is strictly stationary. Find $E(\varepsilon_t^4)$ and the unconditional kurtosis of ε_t . Comment on its implication for the tails of the unconditional distribution of ε_t .

[Hint: for a zero-mean normal random variable $Z \sim N(0, \omega^2)$, $E(Z^4) = 3\omega^4$.]

2. (GARCH model characteristics)

Consider the following AR(1)-GARCH(1,1) model,

$$y_t = c + \phi_1 y_{t-1} + \varepsilon_t, \quad |\phi_1| < 1, \quad \varepsilon_t | \Omega_{t-1} \sim N(0, \sigma_t^2),$$

$$\sigma_t^2 = \alpha_0 + \alpha_1 \varepsilon_{t-1}^2 + \beta_1 \sigma_{t-1}^2, \quad \alpha_0 > 0, \quad \alpha_1 \geq 0, \quad \beta_1 \geq 0, \quad \alpha_1 + \beta_1 < 1,$$

where Ω_t is the information set at the end of period t .

- Find $E(\varepsilon_t | \Omega_{t-1})$, $E(y_t | \Omega_{t-1})$ and their unconditional counterparts.
- Find $\text{Var}(\varepsilon_t | \Omega_{t-1})$, $\text{Var}(y_t | \Omega_{t-1})$ and their unconditional counterparts.
- Is ε_t a white noise process? Is it an independent (or iid) WN process? Verify your answer.
- Ceteris paribus, what is the change in σ_t^2 caused by a one-unit change in ε_{t-1}^2 ?

- (e) Let $w_t = \varepsilon_t^2 - \sigma_t^2$. Show (i) w_t has no autocorrelation; (ii) ε_t^2 has an ARMA(1,1) representation with w_t being the shock.
- (f) Some researchers prefer to write $\alpha_0 = \omega(1 - \alpha_1 - \beta_1)$, where ω is a free parameter (the unconditional variance). Let σ_t^2 be an integrated GARCH(1,1), where $\alpha_1 + \beta_1 = 1$, show that the integrated series is in fact an EWMA of ε_t^2 .



COMPUTING EXERCISE

3. (Estimation of ARCH)

This question is based on the data contained in the Excel file *SHARE.XLS*. The file contains daily data on the S&P500 from the 2nd of January, 1998 to the 30th of December, 2001 comprising a total of 994 observations. The S&P500 index is designated *PRICE* in the file. Generate the series for the percentage log return as $R_t = 100 * (\log(PRICE_t) - \log(PRICE_{t-1}))$.

- (a) Perform the Jarque-Bera test for normality and show the empirical histogram for the returns. Also show the correlogram for the returns and interpret your results.
- (b) Generate the series for the squared return as $R2_t = R_t^2$ and create time series plot of $R2$. Also show the correlogram for squared returns and interpret your results.
- (c) Assume the mean equation for returns is $R_t = c + \varepsilon_t$. Perform an LM test for ARCH effects on the residuals from the regression. Interpret the testing results.
- (d) Assume the mean equation for returns is $R_t = c + \varepsilon_t$. Estimate an ARCH(5) model given the mean equation specified above. Interpret your results. Are the restrictions for the ARCH parameters satisfied? Extract and plot σ_t^2 from the estimated equation. Inspect and comment on the plot. Perform an LM test for ARCH effect on the standardised residual series and comment.
- (e) Compare the histograms of residuals and standardised residuals; and the correlograms of squared residuals and squared standardised-residuals from the model in (d) and comment.
- (f) How would you choose the lags in the variance equation [why ARCH(5)?]? Would a more sophisticated mean equation help? Try some of your suggestions and comment on your results.

程序代写代做 CS编程辅导

4. (Estimation of GARCH)

This question is based on data obtained in the Excel file *SHARE.XLS*. The file contains daily data on the S&P500 index from the 2nd of January, 1998 to the 10th of December, 2001 comprising a total of 2500 observations. The S&P500 index is designated *PRICE* in the file. Generate the series of log returns as: $R = 100 * (\log(PRICE) - \log(PRICE(-1)))$

- (a) Assume the log returns is

$$R_t = c + \varepsilon_t$$

and that the variance equation for returns is a GARCH(1,1). Estimate the model and interpret your results. Are the sign restrictions for the GARCH specification satisfied?

- (b) Extract and plot σ_t^2 from the estimated equation, and make a comparison to the same plot from ARCH(5).

- (c) Perform an LM test on the standardized residuals from this GARCH(1,1) model. Interpret your results.

- (d) Report the Jarque-Bera test for normality on the standardized residuals. Report the correlogram of the squared standardised residuals. Comment on your results.

- (e) Re-estimate the GARCH(1,1) model but do not select heteroscedastic-consistent standard errors. Compare the results with those in (a) and comment.

- (f) Estimate GARCH(2,1), GARCH(1,2) and GARCH(2,2) models. Inspect and comment on the estimation results.