

程序代写代做 CS编程辅导



Copyright of New South Wales 2020. All rights reserved.

Course materials subject to Copyright

UNSW Sydney owns copyright in these materials (unless stated otherwise). The material is subject to copyright under Australian law and overseas under international treaties. The materials are provided for use by enrolled UNSW students. The materials, or any part, may not be copied, shared or distributed, in print or digitally, outside the course without permission. Students may only copy a reasonable portion of the material for personal research or study or for criticism or review. Under no circumstances may these materials be copied or reproduced for sale or commercial purposes without prior written permission of UNSW Sydney.

Statement on class recording

To ensure the free and open discussion of ideas, students may not record, by any means, classroom lectures, discussion and/or activities without the advance written permission of the instructor, and any such recording properly approved in advance can be used solely for the students own private use.

WARNING: Your failure to comply with these conditions may lead to disciplinary action, and may give rise to a civil action or a criminal offence under the law.

THE ABOVE INFORMATION MUST NOT BE REMOVED FROM THIS MATERIAL.

程序代写代做 CS编程辅导

University of New South Wales

School of Economics

Financial Econometrics

Tutorial 7



1. (Value at Risk)

The following GARCH(1,1) model has been estimated (such that the parameters are treated as known), using the historical return series $\{y_t\}$ of a portfolio up to date T :

$$y_t = c + \varepsilon_t, \quad \varepsilon_t | \Omega_{t-1} \sim ?(0, \sigma_t^2),$$

$$\sigma_t^2 = \alpha_0 + \alpha_1 \varepsilon_{t-1}^2 + \beta_1 \sigma_{t-1}^2, \quad t = 1, \dots, T,$$

where Ω_{t-1} is the information set at $t - 1$ and “ $?(0, \sigma_t^2)$ ” is an unknown distribution with mean zero and variance σ_t^2 . Suppose that the portfolio's market value at T was \$10m. How would you calculate the 99% value-at-risk for the period from T to $T+1$? Assume that parameters, ε_T and σ_T^2 are known or already estimated.

2. (GARCH-in-mean model)

Consider the following GARCH-M model

$$y_t = c + \delta \sigma_t^2 + \varepsilon_t, \quad \varepsilon_t | \Omega_{t-1} \sim N(0, \sigma_t^2),$$

$$\sigma_t^2 = \alpha_0 + \alpha_1 \varepsilon_{t-1}^2 + \beta_1 \sigma_{t-1}^2, \quad \alpha_0 > 0, \quad \alpha_1 \geq 0, \quad \beta_1 \geq 0, \quad \alpha_1 + \beta_1 < 1,$$

where y_t is the return of a portfolio and Ω_{t-1} is the information set at $t - 1$.

- Why would the conditional variance σ_t^2 be included in the mean equation? What would be the sign of the parameter δ ?
- Find $E(y_t | \Omega_{t-1})$ and $E(y_t)$.

3. (EGARCH-model)

Consider the constant conditional mean - EGARCH model

$$y_t = c + \varepsilon_t, \quad \varepsilon_t | \Omega_{t-1} \sim N(0, \sigma_t^2)$$

$$\ln(\sigma_t^2) = \alpha_0 + \alpha_1 |v_{t-1}| + \gamma v_{t-1} + \beta_1 \ln(\sigma_{t-1}^2), \quad v_{t-1} = \varepsilon_{t-1} / \sigma_{t-1}$$



- What are the advantages of this model in comparison to GARCH(1,1) model.
- Which sign condition is satisfied and why?
- Compute one-period ahead optimal forecast of y and form 95% confidence bounds.

4. Computing Exercise

(GARCH, extensions and VaR, maybe useful for the project)

This question is based on the data contained in the Excel file *S&P500.XLS*. The file contains daily data on the S&P500 from the 2nd of January, 1998 to the 10th of December, 2001 comprising a total of 994 observations. The S&P500 index is designated *PRICE* in the file. Generate the series for the percentage log return as: $R = 100 * (\log(PRICE) - \log(PRICE(-1)))$.

- Show a graph of the empirical distribution or CDF for R . What is the percentage daily return which cuts off 1% of the left-tail of the empirical distribution? (Or, what is the 1% quantile of the return distribution?).

- Assume the mean equation for returns is $R = c + \varepsilon_{t-1}$ and the variance equation for returns is a GARCH(1,1). Estimate the model using the first 900 observations. (We leave the last 94 observations for use in an out-of-sample forecast exercise). Are the sign restrictions for the GARCH specification satisfied? Present a graph of the conditional standard deviation. Comment on the estimation results.

- Repeat (b) for GJR model [Use the same clicks as (b) but select **1** for Threshold order], and EGARCH model [Use the same clicks as (b) but select **EGARCH** for Model]. Compare the results and decide which model you prefer, which will be used for the questions below.

- For the preferred model, present a histogram and summary statistics of the standardized residuals. Compare and comment the distributions of the residuals and the standardized residuals. What is the 1% quantile of the standardized residuals?

- (e) With the preferred model, generate forecasts for returns and for the conditional standard deviation of returns for the out-of-sample observations 901-994.

Do the forecasts of σ_{t+1}^2 once show “mean-reverting” behavior?

- (f) Suppose a portfolio holds a position of ten million dollars (\$10m) in the market portfolio given by M_t . Calculate the daily empirical 99% Conditional-Value-at-Risk of this portfolio at observation 901. (To do this, you will need to look at the entries for series RF and VF at observation 901. You will also need to use your answers to parts (d) and (e)).

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>