

程序代写代做 CS编程辅导

Copyright © Copyright University of New South
Course materials subject to Copyright
UNSW Sydney owns copyright in these materials (including any text, images, diagrams, etc.). The material is subject to copyright under Australian law and overseas under international treaties. The materials are provided for your personal use only. You may not copy, share or distribute the materials, in print or digitally, outside the course without permission. You may only copy a reasonable portion of the material for personal research or study or for criticism or review. Under no circumstances may the material be copied or reproduced for sale or commercial purposes without prior written permission of UNSW Sydney.

Statement on class recording

To ensure the free and open discussion of ideas, students may not record, by any means, classroom lectures, discussion and/or activities without the advance written permission of the instructor, and any such recording properly approved in advance can be used solely for the student's own private use. **WARNING:** Your failure to comply with these conditions may lead to disciplinary action, and may give rise to a civil action or a criminal offence under the law.

THE ABOVE INFORMATION MUST NOT BE REMOVED FROM THIS MATERIAL



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

F Econometrics
Slides-06: G to ARMA and Forecasting



WeChat: cstutorcs

Dr. Rachida Ouyse

Assignment Project Exam Help

Email: tutorcs@163.com

¹©Copyright University of New South Wales 2020. All rights reserved. This copyright notice must not be removed from this material.

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导



- General $AR(p)$

- Wold Decomposition
- AF and PACF patterns
- Impulse response function

WeChat: cstutorcs

- Yule-Walker equations

- AR & MA mix- ARMA models

- AF and PACF patterns
- Impulse response function

Assignment Project Exam Help

Email: tutorcs@163.com

- Estimation of ARMA

QQ: 749389476

- Forecasting in ARMA

<https://tutorcs.com>

Stationarity of $AR(2)$

程序代写代做 CS编程辅导

The conditions for stationarity/invertibility of an $AR(1)$ process can be extended to higher processes.

- First consider an $AR(1)$

$$(1 - \alpha_1 L) y_t = \alpha_0 + \varepsilon_t.$$

In general, the polynomial $\alpha(L)$ can be rewritten as

$$(1 - \alpha_1 L - \alpha_2 L^2) = (1 - \phi_1 L)(1 - \phi_2 L)$$

where ϕ_1 and ϕ_2 can be solved from $\phi_1 + \phi_2 = \alpha_1$ and $-\phi_1\phi_2 = \alpha_2$

The conditions for invertibility of the second order polynomial are just the conditions that both the first order polynomials

$(1 - \phi_1 L)$ and $(1 - \phi_2 L)$ are invertible, i.e. $|\phi_1| < 1$ and $|\phi_2| < 1$.

Stationarity of $AR(2)$

程序代写代做 CS编程辅导

A more common way of presenting these conditions is in terms of the so-called **characteristic equation**



$$z^2 - \alpha_2 z - \alpha_1 = 0,$$

or $(1 - \phi_2 z)(1 - \phi_1 z) = 0.$

This equation has two solutions, denoted z_1 and z_2

WeChat: tutores

Assignment Project Exam Help

$$z_1, z_2 = \frac{\alpha_1 \pm \sqrt{\alpha_1^2 + 4\alpha_2}}{2\alpha_2},$$

Email: tutores@163.com

referred to as the **characteristic roots** of the $\alpha(L)$ polynomial.

QQ: 749389476

The requirement $|\phi_i| < 1$ corresponds to $|z_i| > 1$. If any solution satisfies $|z_i| \leq 1$, the polynomial $\alpha(L)$ is non-invertible. A solution that equals unity is referred to as a unit root.

https://tutores.com

General Conditions for Stationarity for an $AR(p)$

程序代写代做 CS编程辅导



Calculating the roots of a higher order AR process is computationally not a trivial job. In most circumstances there is little need to directly calculate the characteristic roots, though, as there are some useful **simple rules** for checking stationarity/invertibility of higher order processes

WeChat: cstutorcs

Assignment Project Exam Help

- ▶ **Necessary condition:** $\sum_{i=1}^p \alpha_i < 1$

Email: tutorcs@163.com

- ▶ **Sufficient condition:** $\sum_{i=1}^p |\alpha_i| < 1$

<https://tutorcs.com>

Useful representations

程序代写代做 CS编程辅导



As, under appropriate conditions, an $AR(p)$ process has an $MA(\infty)$ representation and an $MA(q)$ process has an $AR(\infty)$ representation, there is no fundamental difference between AR and MA models.

- ▶ The MA representation is convenient to derive the properties (mean, variance, ...) of a series
- ▶ The AR representation is convenient for making predictions conditional upon the past

When estimating time series models (cf. below), the choice is simply a matter of parsimony.

<https://tutorcs.com>

What is the AR process is stationary?

程序代写代做 CS编程辅导

For a stationary AR(p) process, it is more convenient to derive the properties from imposing $E(y_t) = \mu$, mean, variance and autocovariances do not change with time.

For computational convenience, consider an AR(2) process.

- ▶ The unconditional mean of y_t can be solved from

$$E(y_t) = \alpha_0 + \alpha_1 E(y_{t-1}) + \alpha_2 E(y_{t-2})$$

which, assuming that $E(y_t)$ does not depend on time allows us to write

$$E(y_t) = \alpha_0 / (1 - \alpha_1 - \alpha_2)$$


- ▶ The variance of y_t can be solved by defining $x_t = y_t - E(y_t)$ which yields

$$x_t = \alpha_1 x_{t-1} + \alpha_2 x_{t-2} + \varepsilon_t \quad (18)$$

What is the AR process is stationary?

程序代写代做 CS编程辅导

The variance of y_t can be obtained by multiplying both sides by x_t and taking expectations


$$\begin{aligned} V(y_t) = \gamma_0 &= \alpha_1 \gamma_0 + \alpha_2 \gamma_0 + \alpha_2 x_t x_{t-2} + x_t \varepsilon_t \\ &= \alpha_1 \gamma_0 + \alpha_2 \gamma_0 + E(x_t \varepsilon_t) \\ &= \alpha_1 \gamma_0 + \alpha_2 \gamma_0 + \sigma^2 \end{aligned} \quad (19)$$

where $E(x_t \varepsilon_t) = \sigma^2$ is obtained from multiplying both sides of (18) by ε_t and taking expectations. Multiplying both sides by x_{t-1} and x_{t-2} and taking expectations we obtain

$$\gamma_1 = \alpha_1 \gamma_0 + \alpha_2 \gamma_1 \quad (20)$$

$$\gamma_2 = \alpha_1 \gamma_1 + \alpha_2 \gamma_0 \quad (21)$$

These equations can be solved for γ_0 to obtain

$$\gamma_0 = \frac{\sigma^2}{(1 + \alpha_2)(1 - \alpha_1 - \alpha_2)(1 + \alpha_1 - \alpha_2)}$$

What is the AR process is stationary?

程序代写代做 CS编程辅导

- ▶ The autocorrelation coefficients ρ_1 and ρ_2 can be obtained by dividing (20) and (21) by $1 - \alpha_2$



$$\rho_1 = \alpha_1 + \alpha_2 \rho_1$$

$$\rho_2 = \alpha_1 \rho_1 + \alpha_2$$

and solving to obtain WeChat: cstutorcs

Assignment Project Exam Help

$$\rho_1 = \alpha_1 / (1 - \alpha_2)$$

Email: tutorcs@163.com

It is easily verified that the higher order autocorrelation coefficients are given by

<https://tutorcs.com>

$$\rho_k = \alpha_1 \rho_{k-1} + \alpha_2 \rho_{k-2}$$

Yule Walker Equations

程序代写代做 CS编程辅导

■ The beauty of the Yule Walker Equations!

$$x_t = \alpha_1 x_{t-1} + \cdots + \alpha_p x_{t-p} + \epsilon_t$$

$$x_t x_{t-1} = \alpha_1 x_{t-1} x_{t-1} + \cdots + \alpha_p x_{t-p} x_{t-1} + \epsilon_t x_{t-1}$$

$$E(x_t x_{t-1}) = \alpha_1 E(x_{t-1} x_{t-1}) + \cdots + \alpha_p E(x_{t-p} x_{t-1}) + E(\epsilon_t x_{t-1})$$

$$\gamma_1 = \alpha_1 \gamma_0 + \alpha_2 \gamma_{-1} + \cdots + \alpha_p \gamma_{p-1}$$

...

$$x_t x_{t-j} = \alpha_1 x_{t-1} x_{t-j} + \cdots + \alpha_p x_{t-p} x_{t-j} + \epsilon_t x_{t-j}$$

$$E(x_t x_{t-j}) = \alpha_1 E(x_{t-1} x_{t-j}) + \cdots + \alpha_p E(x_{t-p} x_{t-j}) + E(\epsilon_t x_{t-j})$$

$$\gamma_{|j|} = \alpha_1 \gamma_{|j|-1} + \alpha_2 \gamma_{|j|-2} + \cdots + \alpha_p \gamma_{|p-j|},$$

...

$$\gamma = \Gamma \alpha$$



WeChat: estutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Defining an ARMA Process

程序代写代做 CS编程辅导



Let ε_t be a white noise

Then:

$$\alpha(L)y_t = \alpha_0 + \beta(L)\varepsilon_t \quad (22)$$

with $\alpha(L)$ an AR polynomial of order p and $\beta(L)$ an MA polynomial of order q , is an **autoregressive moving average process** with orders p and q , denoted ARMA(p, q).

→ y_t depends on its own lagged values and on current and past values of a white noise disturbance term ε_t .


QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

■ Dynamic Behaviour and Impulse Response

If the AR polynomial $\alpha(L)$ is stable, the ARMA(p, q) process can be written as a stable MA process of the form


$$\begin{aligned}y_t &= \alpha(L)^{-1} \beta(L) \varepsilon_t \\ &= \alpha'_0 + \theta(L) \varepsilon_t\end{aligned}$$

WeChat: cstutorcs

where $\alpha'_0 = \alpha_0 / 1 - \sum_{i=1}^p \alpha_i$ and $\theta(L) = \alpha(L)^{-1} \beta(L)$
= $1 + \sum_{i=1}^{\infty} \theta_i L^i$, with θ_i = undetermined coefficients.

Assignment Project Exam Help

Even if the AR polynomial is non-invertible, we can still solve for the ε sequence but this solution will not be a stable MA process, i.e.

Email: tutorcs@163.com

QQ: 749389476

$y_t = f(t) + \theta(L) \varepsilon_t$
<https://tutorcs.com>

where $f(t)$ indicates that the mean is a function of time.

程序代写代做 CS编程辅导



■ Dynamic Behaviour Impulse Response

The impulse response function is obtained from the MA representation.

Note that as a finite order MA process is stationary by construction, an ARMA process is stationary if the AR component is stationary (i.e. if the AR polynomial is invertible).

- ▶ In the stationary case the impact of shocks gradually dies out (i.e. $\sum_{i=1}^{\infty} \theta_i$ is finite)
- ▶ In the non-stationary case the impact of a shock never vanishes (i.e. $\sum_{i=1}^{\infty} \theta_i$ is infinite)

<https://tutorcs.com>

General Properties of an ARMA(p,q)

程序代写代做 CS编程辅导

■ Unconditional Moments of an ARMA(p,q)

If the AR polynomial $\alpha(L)$ is invertible the mean, variance and covariances are time-varying.

If the AR polynomial $\alpha(L)$ is invertible, the AR process can be rewritten as the stable infinite MA process. The properties of a stationary AR process can easily be derived from this MA representation.

- ▶ Unconditional mean: $E(y_t) = \alpha_0 / (1 - \sum_{i=1}^p \alpha_i)$
- ▶ Unconditional variance: $V(y_t) = \sigma^2 \sum_{i=0}^{\infty} \theta_i^2$
- ▶ Covariances: $\gamma_k = (\theta_k + \theta_1 \theta_{k+1} + \theta_2 \theta_{k+2} + \dots) \sigma^2$

As an ARMA(p,q) process includes both an AR and an MA component, both the ACF and the PACF do not cut off at some point. As such, it is difficult to determine the order of an ARMA model from the ACF and PACF.

◀ ◻ ▶ ◻ ▶ ◻ ▶ ◻ ▶ ◻ ▶ ◻ ▶ ◻ ▶

Maximum Likelihood Estimation: Intuitive Illustration

程序代写代做 CS编程辅导

This illustration shows a sample of n independent observations, and two continuous distributions $f_1(x)$



Email: tutorcs@163.com

Of these two distributions, which one is the most likely to have generated the sample ?

Although it is not impossible, we don't believe that $f_2(x)$ generated the sample. Why?

On the other hand, the values taken by $f_1(x)$ are substantial for all the observations, which are then where one would expect them to be, would the sample be actually generated by $f_1(x)$.

Maximum Likelihood Estimation

程序代写代做 CS编程辅导



- Maximum Likelihood Estimation is a general method of estimation that can be used for many different types of data and economic models. It has very wide applicability.

WeChat: cstutorcs

- The Maximum Likelihood Estimator (MLE) answers the following question: What are the parameters that are most likely to have generated the observed data given the assumed model.

Email: tutorcs@163.com

- Begin by assuming a model for the outcome variable including a distribution function for the underlying population error term (and hence a distribution for the outcome variable in the population.)

Estimation of ARMA: Maximum Likelihood

程序代写代做 CS编程辅导

- Consider $AR(1)$ model: $y_t = \alpha_0 + \alpha_1 y_{t-1} + \epsilon_t$ where $\epsilon_t \sim \text{i.i.d } N(0, \sigma^2)$.

- it follows: $y_t | \Omega_{t-1} \sim N$

$$y_t | \Omega_{t-1} \sim N[\alpha_0 + \alpha_1 y_{t-1}, [1 - \alpha_1^2]^{-1} \sigma^2]$$

- Conditional pdf:

$$f(y_t | \Omega_{t-1}) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left\{ -\frac{(y_t - \alpha_0 - \alpha_1 y_{t-1})^2}{2\sigma^2} \right\}$$

- Information sets: $\Omega_1 = \{y_1\}, \Omega_2 = \{y_2, \Omega_1\}, \dots, \Omega_t = \{y_t, \Omega_{t-1}\}$.

- Joint pdf for a time series $\{y_1, \dots, y_T\}$ can be factorised:

$$\begin{aligned} f(y_T, y_{T-1}, \dots, y_1) &= f(y_T | \Omega_{T-1}) f(y_{T-1} | \Omega_{T-2}) \cdots f(y_2 | \Omega_1) f(y_1) \\ &= f(y_T | \Omega_{T-1}) f(y_{T-1} | \Omega_{T-2}) \cdots f(y_2 | \Omega_1) f(y_1) \\ &= f(y_T | \Omega_{T-1}) f(y_{T-1} | \Omega_{T-2}) \cdots f(y_2 | \Omega_1) f(y_1) \end{aligned}$$

程序代写代做 CS编程辅导

– Properties of ML Estimators

- When the pdf (likelihood) is correctly specified, the ML estimators have \sqrt{T} sampling properties:
 - consistent,
 - asymptotically normally distributed, and
 - asymptotically efficient.

WeChat: cstutorcs

Allow us to draw inference based on reported SEs.

Assignment Project Exam Help

- When the pdf (likelihood) is incorrect, the “ML” procedure is called quasi (or pseudo) ML.
 - When the normal pdf is used, which may be incorrect, the quasi ML estimators are still consistent and asymptotically normal, as long as the model is defined by the conditional mean and variance that are correctly specified.

Email: tutors@163.com

QQ: 749389476

https://tutors.com

Must use
“robust” SEs

ARMA Process: Identification

程序代写代做 CS编程辅导

The Box-Jenkins App



The so-called Box-Jenkins approach toward fitting ARMA models comprises three stages:

- ▶ **Identification:** determine tentative model(s)
 - ▶ Plot the time series to have a first idea on the DGP (stationary/non-stationary, structural break, ...)
 - ▶ Plot the ACF and the PACF to have a first idea on the order of the ARMA model
- ▶ **Estimation:** estimate the various tentative models
 - ▶ Compare the estimated models using information criteria
 - ▶ Select parsimonious model
- ▶ **Diagnostic checking:** check the selected model's diagnostics

<https://tutorcs.com>

AR Process: Estimation

程序代写代做 CS编程辅导

- ▶ Consider the **AR**(p)

$$y_t = \alpha_0 + \alpha_1 y_{t-1} + \dots + \alpha_p y_{t-p} + \varepsilon_t$$

$$\alpha(L)y_t = \varepsilon_t$$



with ε_t a zero-mean white noise process.

WeChat: cstutorcs

As y_{t-1}, \dots, y_{t-p} are observed in the data, the model can be estimated using **OLS**. **Assignment Project Exam Help**

The OLS estimator is

Email: tutorcs@163.com

- ▶ **biased**, because: $E(y_{t-j}\varepsilon_{t-j}) \neq 0$
- ▶ **consistent**, because: $E(y_{t-j}\varepsilon_t) = 0 \quad \forall j > 0$
- ▶ **asymptotically normal**

QQ: 749389476

Intuition: the error terms and the explanatory variables are not completely independent but contemporaneously uncorrelated.

https://tutorcs.com

MA Process: Estimation

程序代写代做 CS编程辅导

- Consider the **MA**(q) model

$$y_t = \alpha_0 + \beta_1 \varepsilon_{t-1} + \beta_2 \varepsilon_{t-2} + \dots + \beta_q \varepsilon_{t-q} + \varepsilon_t$$

$$y_t = \alpha_0 + \beta$$



with ε_t a zero-mean white noise process.

WeChat: cstutorcs

As $\varepsilon_{t-1}, \dots, \varepsilon_{t-q}$ are NOT observed in the data, the model cannot be directly estimated using OLS.

Assignment Project Exam Help

A possible solution is to estimate the coefficients in $\beta(L)$ from the AR representation of the MA model. For an invertible MA(1) model, this is given by (cf. above)

Email: tutors@163.com

QQ: 749389476

$$y_t = \alpha_0 / (1 + \beta) + \sum_{i=1}^{\infty} (\beta / (1 + \beta))^i y_{t-i} + \varepsilon_t$$

https://tutors.com

Model Selection: order of the ARMA(p,q)

程序代写代做 CS编程辅导

Information criteria

A fundamental idea in the Jenkins approach is the principle of **parsimony** (meaning simplicity)

- ▶ A parsimonious model fits the data well without incorporating any needless coefficients
- ▶ In general, parsimonious models produce better forecasts than over-parametrized models

Increasing the lag orders p and q will:

- ▶ Increase the goodness-of-fit of the model, i.e. reduce the RSS
- ▶ Reduce the degrees of freedom

→ **information criteria** provide a **trade-off** between the goodness-of-fit of the model and the number of parameters used to obtain that fit.



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutores@163.com

QQ: 749389476

https://tutores.com

Model Selection: order of the ARMA(p,q)

程序代写代做 CS编程辅导

The two most commonly used selection criteria are:

- ▶ **Akaike Information Criterion (AIC)**

$$AIC = \ln(\text{RSS}) + 2k$$

- ▶ **Schwarz Bayesian Criterion (SBC)**

$$SIC = \ln(\text{RSS}) + k \ln(T)$$

with $k = p + q + 1$ the number of estimated parameters.

The most appropriate model is the one that minimises AIC and/or SBC.

Model Selection: order of the ARMA(p,q)

程序代写代做 CS编程辅导

Note that:

- ▶ When you estimate models with lagged variables, some initial observations are lost. To compare models using information criteria, **keep T fixed!** Otherwise you will be comparing the performance of the models over different sample periods. Moreover, decreasing T has the direct effect of reducing the AIC and SBC.
- ▶ The SBC embodies a much stiffer penalty for the loss of degrees of freedom than the AIC. The main difference between the two in terms of performance is that SBC is consistent (i.e. asymptotically it delivers the correct model) while the AIC is biased toward selecting an over-parametrised model. However, in small samples, the AIC can work better than the SBC.



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutores@163.com

QQ: 749389476

https://tutores.com

Information Criteria: Example

程序代写代做 CS编程辅导

- If data are generated by an ARMA, the probability that SIC selects the correct model converges to one as $T \rightarrow \infty$.



- In finite samples, SIC may select a too-small model.

WeChat: cstutors

Assignment Project Exam Help

eg. unanticipated

US monthly inflation:

- SIC selects AR(1)
- AIC selects ARMA(1,1)

Email: tutorscs@163.com

QQ: 749389476

<https://tutorscs.com>

Sample (adjusted): 1978M02 1987M12
Included observations: 119 after adjustments
Convergence achieved after 2 iterations

Variable	Coefficient	Std. Error	t-Statistic	Prob.
AR(1)	0.574656	0.076201	7.541340	0.0000
R-squared	0.325217	Mean dependent var		0.000121
Adjusted R-squared	0.325217	S.D. dependent var		0.061977
S.E. of regression	0.050911	Akaike info criterion		-3.109100
Sum squared resid	0.305850	Schwarz criterion		-3.085746
Log likelihood	185.9914	Durbin-Watson stat		1.821211
Inverted AR Roots	.57			

Sample (adjusted): 1978M02 1987M12
Included observations: 119 after adjustments
Convergence achieved after 6 iterations
Backcast: OFF

Variable	Coefficient	Std. Error	t-Statistic	Prob.
AR(1)	0.315395	0.141540	2.228308	0.0278
MA(1)	0.386570	0.136971	2.822271	0.0056
R-squared	0.350263	Mean dependent var		0.000121
Adjusted R-squared	0.344710	S.D. dependent var		0.061977
S.E. of regression	0.050170	Akaike info criterion		-3.130116
Sum squared resid	0.294498	Schwarz criterion		-3.083408
Log likelihood	188.2419	Durbin-Watson stat		2.004409
Inverted AR Roots	.32			
Inverted MA Roots	-.39			

程序代写代做 CS编程辅导

Forecasting using ARMA models



- ▶ In-sample versus out-of-sample forecasts
 - ▶ **In-sample forecasts** are those generated for the same set of data that was used to estimate the model's parameters. Good performance may be due to fitting a spurious model to the noise in the sample, though!
 - ▶ **Out-of-sample forecasts** are those generated for a set of data that was not used to estimate the model, i.e. do not use all observations in estimating the model and evaluate the model from the forecasting accuracy in the holdout sample.
- ▶ Static versus dynamic forecasts
 - ▶ **Static forecasts** are a sequence of one-step-ahead forecasts, using actual, rather than forecasted values for lagged dependent variables.
 - ▶ **Dynamic forecasts** are a sequence of multi-step-ahead forecasts starting from the first period in the forecast sample, using forecasted values for lagged dependent variables.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutormcs@163.com

QQ: 749389476

http://tutormcs.com

程序代写代做 CS编程辅导

Forecasting Accuracy



In addition to the prediction, it is important to know how accurate this prediction is. To judge forecasting accuracy, define the **prediction error** as

WeChat: cstutorcs

Assignment Project Exam Help

and the **variance of the forecasting error** by

Email: tutorescs@163.com

$$\text{var}(f_{t,s}) = E(y_{t,s} - f_{t,s})^2$$

QQ: 749389476

<https://tutorcs.com>

Forecasting MA(q)

程序代写代做 CS编程辅导

For the **MA(q) model** we



$$fe_{t,1} = \varepsilon_{t+1}$$

$$fe_{t,2} = \varepsilon_{t+2} + \beta_1 \varepsilon_{t+1}$$

$$fe_{t,3} = \varepsilon_{t+3} + \beta_1 \varepsilon_{t+2} + \beta_2 \varepsilon_{t+1}$$

\vdots

$$fe_{t,q} = \varepsilon_{t+q} + \beta_1 \varepsilon_{t+q-1} + \dots + \beta_{q-1} \varepsilon_{t+1}$$

$$fe_{t,q+1} = \varepsilon_{t+q+1} + \beta_1 \varepsilon_{t+q} + \dots + \beta_q \varepsilon_{t+1}$$

$$fe_{t,q+2} = \varepsilon_{t+q+2} + \beta_1 \varepsilon_{t+q+1} + \dots + \beta_q \varepsilon_{t+1}$$

\vdots

<https://tutorcs.com>

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749589476

程序代写代做 CS编程辅导

such that



$$\text{var}(fe_{t,1}) = E(\varepsilon_{t+1})^2$$

$$\text{var}(fe_{t,2}) = E(\varepsilon_{t+2} + \beta_1 \varepsilon_{t+1})^2 = (1 + \beta_1^2) \sigma^2$$

$$\text{var}(fe_{t,3}) = E(\varepsilon_{t+3} + \beta_1 \varepsilon_{t+2} + \beta_2 \varepsilon_{t+1})^2 = (1 + \beta_1^2 + \beta_2^2) \sigma^2$$

⋮

$$\begin{aligned} \text{var}(fe_{t,q}) &= E(\varepsilon_{t+q} + \beta_1 \varepsilon_{t+q-1} + \dots + \beta_{q-1} \varepsilon_{t+1})^2 \\ &= (1 + \beta_1^2 + \dots + \beta_{q-1}^2) \sigma^2 \end{aligned}$$

$$\begin{aligned} \text{var}(fe_{t,q+1}) &= E(\varepsilon_{t+q+1} + \beta_1 \varepsilon_{t+q} + \dots + \beta_q \varepsilon_{t+1})^2 \\ &= (1 + \beta_1^2 + \dots + \beta_q^2) \sigma^2 \end{aligned}$$

<https://tutorcs.com>

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

程序代写代做 CS编程辅导



The accuracy of the pred

- ▶ decreases as we predict further into the future
- ▶ does not decrease any further from $s = q + 1$ onward as the variance of the prediction error stabilises at the unconditional variance. This is the upper bound on the inaccuracy of the predictor.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Forecasting AR(p)

程序代写代做 CS编程辅导

For a **stationary AR(p) model**, the prediction errors are most easily obtained from the representation


$$y_t = \alpha_0 + \alpha_1 y_{t-1} + \dots + \alpha_p y_{t-p} + \sum_{i=0}^{\infty} \beta_i \varepsilon_{t-i}$$

with β_i undetermined coefficients

WeChat: cstutorcs

Consequently, the s-period-ahead prediction error is given by

Assignment Project Exam Help

$$fe_{t,s} = \sum_{i=0}^{s-1} \beta_i \varepsilon_{t+s-i}$$

Email: tutorcs@163.com

QQ: 749389476

with variance

$$\text{var}(fe_{t,s}) = \sigma^2 \sum_{i=0}^{s-1} \beta_i^2$$

Forecasting AR(q)

程序代写代做 CS编程辅导

The accuracy of the prediction

- ▶ decreases as we predict into the future as $\beta_i^2 > 0$
- ▶ converges to the stationary conditional variance $\sigma^2 \sum_{i=0}^{\infty} \beta_i^2$ as $t \rightarrow \infty$. This is the end on the inaccuracy of the predictor.

WeChat: cstutorcs

As an illustration, consider an AR(1) model where $\beta_i = \alpha_1^i$. The forecasting errors are given by

Assignment Project Exam Help

$$fe_{t,1} = \varepsilon_{t+1}$$

$$fe_{t,2} = \varepsilon_{t+2} + \alpha_1 \varepsilon_{t+1}$$

QQ: 749389476

$$fe_{t,s} = \sum_{i=0}^{s-1} \alpha_1^i \varepsilon_{t+s-i}$$