

Assignment Project Exam Help

ECON6300/7320/8300

Advanced Microeconometrics

Introduction and Review

<https://tutorcs.com>

Christiern Rose

¹University of Queensland

WeChat: cstutorcs

Lecture 1

Assignment Project Exam Help

- ▶ Christien Rose (Coordinator & Lecturer)
Colin Clark Building (39) Room 537
Email: christiern.rose@uq.edu.au
Consultation: Tuesdays 15:00-17:00
- ▶ <https://tutorcs.com>
Lectures: Fridays, 08:00-10:00
Practicals: Fridays 10:00-12:00 or 12:00-14:00 (starting week 1)
- ▶ Lectures: mainly theoretical "which methods and why"
- ▶ Practical: how to work with data and implement methods (STATA)

WeChat: [cstutorcs](#)

Course Information

- ▶ The course will focus on estimation and inference methods that are widely used in applied microeconomics.
- ▶ The course has a topics-based structure, and theory and applications are closely integrated.
- ▶ Whenever you learn a method theoretically (lecture) there will be a Stata exercise with data for the method (practical)
- ▶ The course will provide econometric skills that could be used in quantitative research at the Graduate level.
- ▶ The course has a strong practical focus, but we will use some basic mathematical/statistical concepts.

Assignment Project Exam Help

- ▶ Required Textbook:
Cameron, A.C. and P.K. Trivedi (2009) Microeconometrics Using Stata
Revised Edition, Stata Press, College Station: TX. **CT-Stata**
- ▶ <https://tutorcs.com>
There are many useful texts, e.g., Greene, Wooldridge, etc. see ECP.
- ▶ As the semester progresses, lecture notes, slides, datasets, problem sets, etc. will be provided. It is however strongly encouraged that you read the relevant part of the textbook and other references

WeChat: cstutorcs

Assignment Project Exam Help

- ▶ You are **EXPECTED** to have a basic grasp of mathematics/statistics. If most of the concepts we cover today are unfamiliar to you, this course may be unsuitable.
- ▶ Random Variables: Expectation, Variance, Covariance
- ▶ Point Estimation, Hypothesis Testing, Interval Estimation
- ▶ Linear Algebra: vector and matrix and their operations

<https://tutorcs.com>
WeChat: cstutorcs

- Informally, a **Random Variable (RV)** takes on numerical values determined by an experiment.

- **Example:** Consider an experiment in which a fair coin is tossed. Then, the possible outcomes are Head and Tail, i.e., $\{H, T\}$. Then, we define a random variable X as follows:

$$X = \begin{cases} 0 & \text{if the outcome is } T \\ 1 & \text{if the outcome is } H \end{cases}$$

The X could be either 0 or 1 whenever the coin is tossed. Each time, the realisation of X would be different.

The uncertainty can be summarised as $\Pr(X = 0) = 0.5$.

- ▶ The uncertainty of a random variable, say X , is represented by its cumulative distribution function (CDF), defined as

Assignment Project Exam Help

- ▶ When $F_X(x)$ is a continuous function, X is said to be a continuous random variable. For a continuous random variable, the probability density function (PDF), denoted by $f_X(x)$, is a function that satisfies

<https://tutorcs.com>

$$\underbrace{\Pr(a < X < b)}_{=F_X(b)-F_X(a)} = \int_a^b f_X(t) dt$$

- ▶ If $F_X(x)$ is differentiable, we have

$$f_X(x) = \frac{d}{dx} F_X(x)$$

- ▶ If $F_X(x)$ is a step function, X is discrete, which we do not cover today.

Math Review expectation

- ▶ A function of a random variable is also a random variable. For example, $\exp(X)$, $\log(X)$, etc. More generally, $g(X)$ with some function $g(\cdot)$.
- ▶ Suppose X have the PDF $f_X(x)$. The expectation of $g(X)$ is

$$E[g(X)] := \int_{-\infty}^{\infty} g(t) \cdot f_X(t) dt$$

which represents the central tendency of the random variable, $g(X)$.

- ▶ As a special case with $g(X) = X$,

$$E[X] := \int_{-\infty}^{\infty} t \cdot f_X(t) dt$$

which is the expectation of X or the expected value or the mean

- ▶ Consider another case with $g(X) = (X - E[X])^2$.

$$V(X) := E[(X - E[X])^2]$$

which is called the variance that represents the variability of X . Note that $\sqrt{V(X)}$ is the standard deviation of X .

- ▶ $E[X]$ and $V(X)$ are the population parameters, or theoretical moments.

You need to know the distribution to compute $E[X]$ and $V(X)$.

- ▶ $E[X]$ must be distinguished from the sample mean (average). For example, when you observe some numbers x_1, \dots, x_n , its average is

$$\bar{x}_n := \frac{1}{n} \sum_{i=1}^n x_i$$

which is NOT the expectation of the distribution

- ▶ Similarly, you must distinguish $V(X)$ and the sample variance

$$\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x}_n)^2$$

1. $E[c] = c$, for any constant c
2. $E[aX + b] = aE[X] + b$ for any constants a and b
3. a_1, \dots, a_n are constants, and X_1, \dots, X_n are RV's. Then,

$$E \left[\sum_{i=1}^n a_i X_i \right] = \sum_{i=1}^n a_i E[X_i].$$

<https://tutorcs.com>

As a special case, we have

$$E \left[\sum_{i=1}^n X_i \right] = \sum_{i=1}^n E[X_i].$$

4. $V(c) = 0$, for any constant c .
5. $V(aX + b) = a^2 \text{Var}(X)$ for any constants a and c .

- ▶ Suppose X and Y are **jointly** distributed with the **joint PDF** $f_{XY}(x, y)$.

- ▶ The marginal PDF of X can be obtained by integrating Y out

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy$$

- ▶ X and Y are **independent** if and only if for all x and y

$$f_{XY}(x, y) = f_X(x)f_Y(y)$$

- ▶ The expectation of $g(X, Y)$ for some function $g(X, Y)$ is

$$E[g(X, Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) f_{XY}(x, y) dx dy$$

- ▶ If $g(X, Y) = (X - E[X])(Y - E[Y])$, then we have the covariance,

$$C(X, Y) = E[(X - E[X])(Y - E[Y])]$$

- ▶ If $C(X, Y) > 0$, the X and Y move in the same direction.
If $C(X, Y) < 0$, they move in the opposite direction.

- ▶ $|C(X, Y)| \leq \sqrt{V(X)}\sqrt{V(Y)}$. So, the correlation coefficient

$$\rho_{XY} := \frac{C(X, Y)}{\sqrt{V(X)}\sqrt{V(Y)}}$$

is always between -1 and 1.

- ▶ If X and Y are independent, $C(X, Y) = 0$. But, the converse is not generally true

Assignment Project Exam Help

- Suppose X and Y are jointly distributed with the joint PDF $f_{XY}(x, y)$. Then, the conditional PDF of Y given $X = x$ is

$$f_{Y|X}(y|x) := \frac{f_{XY}(x, y)}{f_X(x)}$$

which summarises the distribution of Y when X takes a value x .

- If X and Y are independent, $f_{Y|X}(y|x) = f_Y(y)$ and $f_{X|Y}(x|y) = f_X(x)$

WeChat: cstutorcs

- ▶ The conditional expectation of Y given $X = x$ is

$$E[Y|X = x] = \int_{-\infty}^{\infty} y f_{Y|X}(y|x) dy$$

- ▶ The conditional variance of Y given $X = x$ is

$$V(Y|X = x) = E[(Y - E[Y|X = x])^2 | X = x]$$

- ▶ $E[Y|X = x]$ and $V(Y|X = x)$ are functions of x .
- ▶ When x is not specified, $E[Y|X]$ and $V(Y|X)$ are random.

Assignment Project Exam Help

1. $E[c(X)|X] = c(X)$ for any function $c(\cdot)$
2. $E[a(X)Y + b(X)|X] = a(X)E[Y|X] + b(X)$ for any functions $a(\cdot), b(\cdot)$
3. If X and Y are independent, $E[Y|X] = E[Y]$
4. $E[E[Y|X]] = E[Y]$. This is called the **law of iterated expectations**.
5. If X and Y are independent, $V(Y|X) = V(Y)$

<https://tutorcs.com>
WeChat: cstutorcs

Assignment Project Exam Help

- Suppose a random variable Y follows a normal distribution with mean μ and variance σ^2 . Or simply we can write

$$Y \sim \mathcal{N}(\mu, \sigma^2)$$

- The Y takes on a value from $\mathbb{R} = (-\infty, \infty)$. Whenever it is drawn from $\mathcal{N}(\mu, \sigma^2)$, the realisation of Y will be different. The uncertainty is summarised as its probability density function (PDF),

$$f(y|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2\sigma^2}(y - \mu)^2\right)$$


Math Review normal distribution

- ▶ The shape of $f_Y(y|\mu, \sigma^2)$ is given as follows when $\mu = 0$ and $\sigma^2 = 1$.

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs



- ▶ It is symmetric about μ and the scale is determined by $\sigma := \sqrt{\sigma^2}$
- ▶ The whole shape (distribution) is completely determined by (μ, σ^2)

- ▶ $E[Y] = \mu$ and $V(Y) = \sigma^2$ for $Y \sim \mathcal{N}(\mu, \sigma^2)$,

- ▶ $\mathcal{N}(0, 1)$ is called the standard normal distribution (the graph above), often used for statistical inference

- ▶ I simulated three number from $\mathcal{N}(0, 1)$ using my computer and I obtained
 $\{0.5377, 1.8339, -2.2588\}$

- ▶ These numbers are **realisations** of $\mathcal{N}(0, 1)$. Their average is NOT the expectation of $\mathcal{N}(0, 1)$. In fact the average is 0.0376 while the expectation is 0.

- ▶ But, the average seems quite close to the mean

- Suppose Y_1, \dots, Y_n are identically and independently distributed as $\mathcal{N}(\mu, \sigma^2)$, i.e.

$$Y_1, \dots, Y_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2),$$

and also assume that μ is unknown but σ^2 is known. That is, if we knew the **parameter** μ , we would know all.

- We wish to learn μ from the **random sample** of size n , i.e., Y_1, \dots, Y_n . Especially, we consider the **(point) estimator**

$$\hat{\mu}_n := \frac{1}{n} \sum_{i=1}^n Y_i$$

- The estimator $\hat{\mu}_n$ is random because Y_1, \dots, Y_n are all random. (Its realisation is an **estimate** of $\hat{\mu}_n$.)

Assignment Project Exam Help

- ▶ **Result:** a linear transformation of normal variables is a normal.

- ▶ This implies that

$$\hat{\mu}_n = \frac{1}{n} \sum_{i=1}^n Y_i$$

<https://tutorcs.com>

- ▶ That is, we know that

$$\hat{\mu}_n \sim \mathcal{N}(E[\hat{\mu}_n], V(\hat{\mu}_n))$$

- ▶ Then, what are the mean and the variance of the estimator $\hat{\mu}_n$?

[WeChat: cstutorcs](#)

Assignment Project Exam Help

► **Result 2:** $E[X + Z] = E[X] + E[Z]$ for any two RVs X and Z

► This implies that

$$E[\hat{\mu}_n] = \frac{1}{n} \sum_{i=1}^n E[Y_i] = \frac{1}{n} \sum_{i=1}^n \mu = \frac{1}{n} (n\mu) = \mu$$

which means

$$\hat{\mu}_n \sim \mathcal{N}(\mu, V(\hat{\mu}_n))$$

► The estimator $\hat{\mu}_n$ is unbiased: $E[\hat{\mu}_n] = \mu$

► **Result:** $V(aX) = a^2 V(X)$ when X is random and a is a constant

► **Result:** $V(X + Z) = V(X) + V(Z)$ if X and Z are independent

► Therefore,

$$V[\bar{X}_n] = V\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n^2} \sum_{i=1}^n V(X_i) = \frac{1}{n^2} \sum_{i=1}^n \sigma^2 = \frac{1}{n^2} (n\sigma^2) = \frac{\sigma^2}{n}$$

► Finally, we have

WeChat: $\bar{X}_n \sim N\left(\mu, \frac{\sigma^2}{n}\right)$ tutors

where μ is the unknown parameter and σ^2 is assumed to be known.

Assignment Project Exam Help

- ▶ Recall that

$$\hat{\mu}_n \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right)$$

especially, $E[\hat{\mu}_n] = \mu$ and $V(\hat{\mu}_n) = \frac{\sigma^2}{n}$

- ▶ The estimator is distributed centred around μ and its variance gets smaller when n grows,
- ▶ This suggests that $\hat{\mu}_n$ gets 'very' close to μ as $n \rightarrow \infty$.
- ▶ This quality is referred to as the consistency of the estimator.

Assignment Project Exam Help

- Suppose X_1, X_2, \dots , are sequence of random variables such that

$$\lim_{n \rightarrow \infty} \Pr(|X_n - c| > \varepsilon) = 0,$$

for any $\varepsilon > 0$. Then X_n converges in probability to c , and we often write

$$X_n \xrightarrow{p} c$$

- $\hat{\mu}_n$ is consistent to μ if $\hat{\mu}_n \xrightarrow{p} \mu$

WeChat: cstutorcs

- ▶ When

$$Y \sim \mathcal{N}(\mu, \sigma^2),$$

the properties of normal random variables implies

$$\frac{Y - \mu}{\sigma} \sim \mathcal{N}(0, 1)$$

Exercise: verify this.

- ▶ Back to our estimator $\hat{\mu}_n$: since

$$\hat{\mu}_n \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right)$$

we have

$$\frac{\hat{\mu}_n - \mu}{\sqrt{\sigma^2/n}} \sim \mathcal{N}(0, 1)$$

Assignment Project Exam Help

- ▶ Recall that when $Z \sim \mathcal{N}(0, 1)$

$\Pr(|Z| \leq 1.96) = 0.95$ or, equivalently, $\Pr(|Z| > 1.96) = 0.05$

- ▶ That is, the event that $|Z| \leq 1.96$ happens with 95% probability.
But, the event that $|Z| > 1.96$ happens with 5% probability.

- ▶ So, the event that $|Z| \leq 1.96$ is likely to happen.
But, the event that $|Z| > 1.96$ is unlikely to happen.

- ▶ Conversely, when the event that $|Z| > 1.96$ happens, you might want to suspect whether Z really follows $\mathcal{N}(0, 1)$

Assignment Project Exam Help

- ▶ We know

$$\frac{\hat{\mu}_n - \mu}{\sqrt{\sigma^2/n}} \sim \mathcal{N}(0, 1)$$

but we do not know the true value of μ .

- ▶ So let's assume that $\mu = \mu_0$ for some number μ_0 and study whether this value is reasonable with respect to our distributional knowledge of $\hat{\mu}_n$.
- ▶ We test the null hypothesis

$H_0: \mu = \mu_0$
WeChat: cstutorcs

- Under the null hypothesis H_0 , we know $\frac{\hat{\mu}_n - \mu_0}{\sqrt{\sigma^2/n}} \sim \mathcal{N}(0, 1)$ and the event

Assignment Project Exam Help

happens with 5% of probability (low prob).

- Since μ_0 is a hypothesised value, we consider this unlikely event as a statistical evidence against H_0 . So, we reject H_0 .
- We admit that even if H_0 is correct, we could mistakenly reject H_0 with 5% of probability. This error rate is called the size (level) of the test.
- You could have a different size. For example, you could use

$$P(|Z| > 2.5758) = 0.01$$

Assignment Project Exam Help

- ▶ A confidence interval (CI) can be obtained by inverting the test
- ▶ For example, 95% CI is the set of all that values for μ_0 that would not be rejected by the test of 5% size. After some algebra, we construct

$$95\% \text{ CI} = \left[\hat{\mu}_n - 1.96\sqrt{\sigma^2/n}, \hat{\mu}_n + 1.96\sqrt{\sigma^2/n} \right]$$

or simply we may write $\hat{\mu}_n \pm 1.96\sqrt{\sigma^2/n}$

- ▶ CI = interval estimator

WeChat: cstutorcs

Assignment Project Exam Help

- ▶ We relax the normality assumption. That is, Y_1, \dots, Y_n are i.i.d from some distribution (may not be normal) with mean μ and variance σ^2

- ▶ Two useful asymptotic results:

<https://tutorcs.com>

$$\bar{\mu}_n \xrightarrow{p} \mu$$

Weak Law of Large Numbers (WLLN)

$$\sqrt{n}(\hat{\mu}_n - \mu)/\sigma \xrightarrow{d} \mathcal{N}(0, 1)$$

Central Limit Theorem (CLT)

- ▶ WLLN directly implies consistency of $\hat{\mu}_n$

[WeChat: cstutorcs](#)

Assignment Project Exam Help

• The CLT

$$\sqrt{n}(\hat{\mu}_n - \mu)/\sigma \xrightarrow{d} \mathcal{N}(0, 1)$$

implies that when n is large

<https://tutorcs.com>

where $\stackrel{a}{\sim}$ is 'approximately distributed,' which can be rewritten as

WeChat: $\frac{\hat{\mu}_n - \mu}{\sqrt{\sigma^2/n}} \stackrel{a}{\sim} \mathcal{N}(0, 1)$ [estutorcs](#)

- ▶ So, the hypothesis testing and confidence interval are asymptotically valid even without the normality assumption.
- ▶ For example, we reject $H_0 : \mu = \mu_0$ at 5% level if

$$\left| \frac{\hat{\mu}_n - \mu_0}{\sqrt{\sigma^2/n}} \right| > 1.96$$

<https://tutorcs.com>
and the asymptotic 95% confidence interval is

$$\hat{\mu}_n \pm 1.96 \sqrt{\sigma^2/n}$$

- ▶ Under normality, $\frac{\hat{\mu}_n - \mu_0}{\sqrt{\sigma^2/n}} \sim \mathcal{N}(0, 1)$ but
without normality $\frac{\hat{\mu}_n - \mu_0}{\sqrt{\sigma^2/n}} \stackrel{a}{\sim} \mathcal{N}(0, 1)$

WeChat: cstutorcs

- ▶ So far, we have assumed that σ^2 is known. Now, we relax this assumption.

- ▶ If σ^2 is unknown, we cannot compute the test statistic and the confidence interval.

- ▶ Suppose we have a consistent estimator for σ^2 , i.e.,

$$\hat{\sigma}_n^2 \xrightarrow{p} \sigma^2,$$

- ▶ Exercise: show that the CLT and Slutsky lemma implies

$$\sqrt{n}(\hat{\mu}_n - \mu)/\hat{\sigma}_n \xrightarrow{d} \mathcal{N}(0, 1)$$

Assignment Project Exam Help

Then, this result

$$\sqrt{n}(\hat{\mu}_n - \mu)/\hat{\sigma}_n \xrightarrow{d} \mathcal{N}(0, 1)$$

implies that when n is large

$$\sqrt{n}(\hat{\mu}_n - \mu)/\hat{\sigma}_n \stackrel{a}{\sim} \mathcal{N}(0, 1)$$

where $\stackrel{a}{\sim}$ is 'approximately distributed,' which can be rewritten as

$$\frac{\hat{\mu}_n - \mu}{\hat{\sigma}_n/\sqrt{n}} \stackrel{a}{\sim} \mathcal{N}(0, 1)$$

WeChat: cstutorcs

Assignment Project Exam Help

- ▶ Therefore, we can construct the asymptotic testing and confidence interval as before.

- ▶ Specifically, we reject $H_0 : \mu = \mu_0$ at 5% level if

$$\frac{|\hat{\mu}_n - \mu_0|}{\sqrt{\hat{\sigma}_n^2/n}} > 1.96$$

and the asymptotic 95% confidence interval is

$$\hat{\mu}_n \pm 1.96 \sqrt{\hat{\sigma}_n^2/n}$$

- ▶ Note that $\sqrt{\hat{\sigma}_n^2/n}$ is called the standard error, $se(\hat{\mu}_n) = \sqrt{\hat{\sigma}_n^2/n}$.

- ▶ Recall that $Z \sim \mathcal{N}(0, 1)$. Then, $Z^2 \sim \chi^2(1)$.

- ▶ Also, $Z_1, \dots, Z_K \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$. Then, $\sum_{j=1}^K Z_j^2 \sim \chi^2(K)$

- ▶ Under $H_0 : \mu = \mu_0$, since

$$\frac{\hat{\mu}_n - \mu_0}{\sqrt{\hat{\sigma}_n^2/n}} \stackrel{a}{\sim} \mathcal{N}(0, 1)$$

we have

$$W_n := \left(\frac{\hat{\mu}_n - \mu_0}{\sqrt{\hat{\sigma}_n^2/n}} \right)^2 \stackrel{a}{\sim} \chi^2(1)$$

- ▶ The LHS is the Wald statistic. At level 5%, we reject H_0 if $W_n > (1.96)^2$.

- We started from the result

$\sqrt{n}(\hat{\mu}_n - \mu)/\sigma \xrightarrow{d} \mathcal{N}(0, 1)$ or $\sqrt{n}(\hat{\mu}_n - \mu) \xrightarrow{d} \mathcal{N}(0, \sigma^2)$
and by squaring both sides and using $\hat{\sigma}_n^2 \xrightarrow{p} \sigma^2$, we obtain

$$W_n = n(\hat{\mu}_n - \mu) (\hat{\sigma}_n^2)^{-1} (\hat{\mu}_n - \mu) \stackrel{a}{\sim} \chi^2(1)$$

- More generally, for a parameter $\theta \in \mathbb{R}^k$ and its estimator $\hat{\theta}_n$, if

$$\sqrt{n}(\hat{\theta}_n - \theta) \xrightarrow{d} \mathcal{N}(0, V),$$

we can construct the Wald statistic

$$n(\hat{\theta}_n - \theta)' \hat{V}^{-1} (\hat{\theta}_n - \theta) \stackrel{a}{\sim} \chi^2(k)$$

with a consistent estimator $\hat{V} \xrightarrow{p} V$, the asymptotic variance covariance matrix of $\hat{\theta}$. You will see this sandwich form often.

- ▶ As mentioned briefly, we often use a vector and matrix notation in econometrics. Vectors and matrices are collections of numbers in a certain form.

- ▶ Let $\mathbf{a} := \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix}$ and $\mathbf{b} := \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix}$ be n dimensional column vectors.

- ▶ Then, $\mathbf{a}' = (a_1 \quad \dots \quad a_n)$, the transpose of \mathbf{a} , is an n -dim row vector.

- ▶ Addition is defined when two vectors are in the same form.

$$\mathbf{a} + \mathbf{b} = \begin{pmatrix} a_1 + b_1 \\ \vdots \\ a_n + b_n \end{pmatrix} \text{ and } \mathbf{a}' + \mathbf{b}' = (\mathbf{a} + \mathbf{b})'. \text{ But } \mathbf{a}' + \mathbf{b} \text{ is not defined.}$$

- ▶ Inner product

$$(\mathbf{a}, \mathbf{b}) := \mathbf{a}' \mathbf{b} = \begin{pmatrix} a_1 & \dots & a_n \end{pmatrix} \cdot \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix} = \sum_{i=1}^n a_i b_i$$

- ▶ Consider an $n \times k$ dimensional matrix:

$$\mathbf{A} := \begin{pmatrix} a_{11} & \dots & a_{1k} \\ \vdots & & \vdots \\ a_{n1} & \dots & a_{nk} \end{pmatrix} = \begin{pmatrix} \mathbf{a}_1 & \dots & \mathbf{a}_k \end{pmatrix}$$

where $\mathbf{a}_j = \begin{pmatrix} a_{1j} \\ \vdots \\ a_{nj} \end{pmatrix}$ for $j = 1, \dots, k$.

- ▶ a vector is a special case of a matrix (e.g., $k = 1$)

- ▶ Transpose:

$$\mathbf{A}' := \begin{pmatrix} a_{11} & \dots & a_{n1} \\ \vdots & & \vdots \\ a_{1k} & \dots & a_{nk} \end{pmatrix} = \begin{pmatrix} \mathbf{a}'_1 \\ \vdots \\ \mathbf{a}'_k \end{pmatrix}$$

- ▶ property: $(AB)' = B'A'$

- ▶ Addition is allowed between two matrices in the same dimension. Let

$$\mathbf{B} := \begin{pmatrix} b_{11} & \dots & b_{1k} \\ \vdots & & \vdots \\ b_{n1} & \dots & b_{nk} \end{pmatrix} = (\mathbf{b}_1 \quad \dots \quad \mathbf{b}_k)$$

where \mathbf{b}_i denotes the i^{th} column of \mathbf{B} . Then

$$\mathbf{A} + \mathbf{B} = \begin{pmatrix} a_{11} + b_{11} & \dots & a_{1k} + b_{1k} \\ \vdots & & \vdots \\ a_{n1} + b_{n1} & \dots & a_{nk} + b_{nk} \end{pmatrix}$$

But, $\mathbf{A}' + \mathbf{B}$ is not defined unless $k = n$, i.e., \mathbf{A} and \mathbf{B} are square

- To explain multiplication, we define a $k \times m$ matrix

Assignment Project Exam Help

$$\mathbf{C} := \begin{pmatrix} c_{11} & \dots & c_{1m} \\ \vdots & & \vdots \\ c_{k1} & \dots & c_{km} \end{pmatrix}$$

Then,

$$\mathbf{A} \times \mathbf{C} := \begin{pmatrix} a_{11} & \dots & a_{1k} \\ \vdots & & \vdots \\ a_{n1} & \dots & a_{nk} \end{pmatrix} \begin{pmatrix} c_{11} & \dots & c_{1m} \\ \vdots & & \vdots \\ c_{k1} & \dots & c_{km} \end{pmatrix}$$

$$:= \begin{pmatrix} \sum_{\ell=1}^k a_{1\ell} c_{\ell 1} & \sum_{\ell=1}^k a_{1\ell} c_{\ell 2} & \dots & \sum_{\ell=1}^k a_{1\ell} c_{\ell m} \\ \sum_{\ell=1}^k a_{2\ell} c_{\ell 1} & \sum_{\ell=1}^k a_{2\ell} c_{\ell 2} & \dots & \sum_{\ell=1}^k a_{2\ell} c_{\ell m} \\ \vdots & \vdots & & \vdots \\ \sum_{\ell=1}^k a_{n\ell} c_{\ell 1} & \sum_{\ell=1}^k a_{n\ell} c_{\ell 2} & \dots & \sum_{\ell=1}^k a_{n\ell} c_{\ell m} \end{pmatrix}$$

- ▶ The $(i, j)^{\text{th}}$ entry of $\mathbf{A} \times \mathbf{C}$ is $\sum_{\ell=1}^k a_{i\ell} c_{\ell j}$

- ▶ Multiplication is defined only when the number of columns of the first matrix is equal to the number of rows of the second matrix.

- ▶ For example, $\underbrace{\mathbf{A}}_{n \times k} \times \underbrace{\mathbf{C}}_{k \times m}$ is well defined and the resulting matrix is $n \times m$ dimensional but $\underbrace{\mathbf{C}}_{k \times m} \times \underbrace{\mathbf{A}}_{n \times k}$ is not defined unless $m = n$.

- ▶ Suppose \mathbf{A} is a square matrix, i.e. $n = k$.
 a_{ij} is called a diagonal element if $i = j$. Otherwise, it is off-diagonal.
If off-diagonal elements are all zero, \mathbf{A} is said to be a diagonal matrix.
If $\mathbf{A}' = \mathbf{A}$, it is symmetric ($a_{ij} = a_{ji}$ for all (i, j))

- ▶ If \mathbf{A} is diagonal and all diagonal elements are ones, \mathbf{A} is an n dimensional identity matrix, for which a reserved notation is \mathbf{I}_n . So,

$$\mathbf{I}_n := \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix}$$

- ▶ A multiplication with the identity does not change.
That is, $\mathbf{A} \times \mathbf{I} = \mathbf{A}$ and $\mathbf{I} \times \mathbf{B} = \mathbf{B}$.

- ▶ For a square matrix \mathbf{A} , if there is another matrix \mathbf{B} such that $\mathbf{I} := \mathbf{AB} = \mathbf{BA}$, then \mathbf{B} is called the inverse of \mathbf{A} and denoted by \mathbf{A}^{-1} , and \mathbf{A} is said to be invertible or nonsingular.

- ▶ (i) $(\alpha \mathbf{A})^{-1} = \frac{1}{\alpha} \mathbf{A}^{-1}$ for a constant α ,
(ii) $(\mathbf{AB})^{-1} = \mathbf{B}^{-1} \mathbf{A}^{-1}$,
(iii) $(\mathbf{A}')^{-1} = (\mathbf{A}^{-1})'$

Assignment Project Exam Help

- ▶ For any $n \times n$ matrix A , the trace of A , $tr(A)$, is the sum of diagonal elements

$$tr(A) = \sum_{i=1}^n a_{ii}$$

<https://tutorcs.com>

- ▶ (i) $tr(I_n) = n$, (ii) $tr(A') = tr(A)$, (iii) $tr(A + B) = tr(A) + tr(B)$,
(iv) $tr(\alpha A) = \alpha tr(A)$ for a constant α
(v) $tr(AB) = tr(BA)$ for any dimensionality conformable A and B

WeChat: cstutorcs

- ▶ Let x_1, \dots, x_n be a set of $n \times 1$ vectors. These are **linearly independent** if and only if

$$\alpha_1 x_1 + \alpha_2 x_2 + \dots + \alpha_n x_n = 0$$

implies $\alpha_1 = \alpha_2 = \dots = \alpha_n = 0$.

- ▶ When x_1, \dots, x_n are linearly independent, x_j cannot be represented as a linear combination of others.
- ▶ Let X be an $n \times k$ matrix ($n \geq k$). The rank of X , denoted by $\text{rank}(X)$, is the maximum number of linearly independent column vectors in X .
- ▶ If $\text{rank}(X) = k$, X has full column rank.

Assignment Project Exam Help

- Let A be an $n \times n$ symmetric matrix. The **quadratic form** with A is a function $f(x)$ defined for all $n \times 1$ vectors

$$f(x) = x'Ax$$

Then, A is **positive definite** (p.d.) if $x'Ax > 0$ for all non-zero x .
Moreover, A is **positive semi-definite** (p.s.d.) if $x'Ax \geq 0$ for all x .

- (i) If A is p.d., then $a_{ii} > 0$ for all i .
(ii) If A is p.s.d., then $a_{ii} \geq 0$ for all i .
(iii) If A is p.d., then A is invertible i.e. A^{-1} exists.
(iv) If X is $n \times k$, then $X'X$ and XX' are p.s.d.
(v) If X is $n \times k$ and $\text{rank}(X) = k$, then $X'X$ is p.d. (so invertible)

- For a given $n \times 1$ vector a , consider the linear function

$$f(x) = a'x$$

for all $n \times 1$ vectors x . Then,

$$\frac{\partial f(x)}{\partial x} = a$$

- For an $n \times n$ symmetric matrix A , define a quadratic function

$$g(x) = x'Ax$$

for all $n \times 1$ vectors x . Then,

$$\frac{\partial g(x)}{\partial x} = 2Ax$$

- Note $\frac{\partial f(x)}{\partial x'} = \left(\frac{\partial f(x)}{\partial x} \right)'$ and $\frac{\partial g(x)}{\partial x'} = \left(\frac{\partial g(x)}{\partial x} \right)'$

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

- ▶ Let y be an $n \times 1$ vector and X be $n \times k$. Consider

Assignment Project Exam Help

- ▶ $(y - X\beta)' = y' - (X\beta)' = y' - \beta'X'$. So, the objective function is

<https://tutorcs.com>

$$\begin{aligned} & (y - X\beta)'(y - X\beta) \\ &= (y' - \beta'X')(y - X\beta) \\ &= y'y - y'X\beta - \beta'X'y + \beta'X'X\beta \\ &= y'y - 2y'X\beta + \beta'X'X\beta \end{aligned}$$

- ▶ WeChat: cstutorcs

$$\min_{\beta} (y - X\beta)'(y - X\beta) = \min_{\beta} (y'y - 2y'X\beta + \beta'X'X\beta)$$

- ▶ Then, the first order necessary condition (FOC) is

$$\frac{\partial}{\partial \beta} (y'y - 2y'X\beta + \beta'X'X\beta) = 2\frac{\partial}{\partial \beta} y'X\beta + \frac{\partial}{\partial \beta} \beta'X'X\beta$$

- ▶ Applying the rules above,

$$\frac{\partial}{\partial \beta} y'X\beta = X'y \quad \frac{\partial}{\partial \beta} \beta'X'X\beta = 2X'X\beta$$

- ▶ The FOC is then

$$-2X'y + 2X'X\beta = 0 \iff X'y = X'X\beta$$

at the optimal solution for β , say $\hat{\beta}$. The solution exists if and only if $X'X$ is invertible;

$$\hat{\beta} = (X'X)^{-1}X'y$$

$\text{rank}(X) = k \implies (X'X)$ is p.d. (therefore invertible)

Assignment Project Exam Help

- Suppose we have n random variables, Y_1, \dots, Y_n . Then, define a random vector

$$Y := \begin{pmatrix} Y_1 \\ \vdots \\ Y_n \end{pmatrix}, \quad \text{and then we have } E[Y] := \begin{pmatrix} E[Y_1] \\ \vdots \\ E[Y_n] \end{pmatrix}$$

<https://tutorcs.com>

- Also $V(Y) := E[(Y - E[Y])(Y - E[Y])']$.
- The $(i, j)^{\text{th}}$ element of $V(Y)$ is $C(Y_i, Y_j)$.
- The i^{th} diagonal element of $V(Y)$ is $C(Y_i, Y_i) = V(Y_i)$.
- $V(Y)$ is symmetric because $C(Y_i, Y_j) = C(Y_j, Y_i)$,

[WeChat: cstutorcs](https://tutorcs.com)

Assignment Project Exam Help

- ▶ Let X be another random variable. Then,

$$E[Y|X] := \begin{pmatrix} E[Y_1|X] \\ \vdots \\ E[Y_n|X] \end{pmatrix}$$

and

$$V(Y|X) := E[(Y - E[Y|X])(Y - E[Y|X])'|X]$$

WeChat: cstutorcs