

程序代写代做 CS编程辅导



Lecture - 6
WeChat: estutorcs

Assignment Project Exam Help
Digital Signal Processing

Email: tutorcs@163.com
ENGN 4537 / 6537

QQ: 749389476

Semester 2, 2023
<https://tutorcs.com>

Inverse z-Transform

程序代写代做 CS编程辅导

The z -transform



$$\sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

Question: If we are given $X(z)$, how can we infer $x[n]$?

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Inverse z-Transform

程序代写代做 CS编程辅导

The z -transform



$$\sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

Question: If we are given $X(z)$, how can we infer $x[n]$?

WeChat: cstutorcs

The inverse of z -transform is formally defined as

$$x[n] = \frac{1}{2\pi j} \oint_C X(z)z^{n-1}dz$$

Assignment Project Exam Help
Email: tutorcs@163.com

where C represents a closed contour within ROC of the z -transform.

QQ: 749389476
<https://tutorcs.com>

Inverse z-Transform

程序代写代做 CS编程辅导

The z -transform



$$\sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

Question: If we are given $X(z)$, how can we infer $x[n]$?

WeChat: cstutorcs

The inverse of z -transform is formally defined as

$$x[n] = \frac{1}{2\pi j} \oint_C X(z)z^{n-1}dz$$

Assignment Project Exam Help
Email: tutorcs@163.com

where C represents a closed contour within ROC of the z -transform.

QQ: 749389476
<https://tutorcs.com>

For the purpose of calculating inverse transforms for most practical purposes, we use **simpler techniques** rather than using this integral. (**We are not going to use this formula directly**)

Method 1: Inspection Method

程序代写代做 CS编程辅导

Simply by inspection and familiarity with Table 3.1 and Table 3.2 of the z-transforms.

TABLE 3.1 SOME COM



S

orm

ROC

Sequence		
1. $\delta[n]$	1	All z
2. $u[n]$	$\frac{1}{1 - z^{-1}}$	$ z > 1$
3. $-u[-n - 1]$	$\frac{1}{z^{-1}}$	$ z < 1$
4. $\delta[n - m]$	$\frac{1}{z^{-m}}$	All z except 0 (if $m > 0$) or ∞ (if $m < 0$)
5. $a^n u[n]$	$\frac{1}{1 - az^{-1}}$	$ z > a $
6. $-a^n u[-n - 1]$	$\frac{1}{1 - az^{-1}}$	$ z < a $
7. $na^n u[n]$	$\frac{az^{-1}}{(1 - z^{-1})^2}$	$ z > a $
8. $-na^n u[-n - 1]$	$\frac{az^{-1}}{(1 - az^{-1})^2}$	$ z < a $
9. $\cos(\omega_0 n)u[n]$	$\frac{z - \cos(\omega_0)}{z^2 - 2\cos(\omega_0)z + 1}$	$ z > 1$
10. $\sin(\omega_0 n)u[n]$	$\frac{\sin(\omega_0)z^{-1}}{1 - 2\cos(\omega_0)z^{-1} + z^{-2}}$	$ z > 1$
11. $r^n \cos(\omega_0 n)u[n]$	$\frac{z - \cos(\omega_0)}{z^2 - 2r\cos(\omega_0)z + r^2}$	$ z > r$
12. $r^n \sin(\omega_0 n)u[n]$	$\frac{r \sin(\omega_0)z^{-1}}{1 - 2r\cos(\omega_0)z^{-1} + r^2z^{-2}}$	$ z > r$
13. $\begin{cases} a^n, & 0 \leq n \leq N - 1, \\ 0, & \text{otherwise} \end{cases}$	$\frac{1 - a^N z^{-N}}{1 - az^{-1}}$	$ z > 0$

WeChat: cstutorcs
Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Method 1: Inspection Method

程序代写代做 CS 编程辅导

TABLE 3.2 SOME z -TRANSFORM PAIRS

Property Number	Section Reference	Transform	ROC
		$X(z)$	R_x
		$X_1(z)$	R_{x_1}
		$x_2[n]$	R_{x_2}
1	3.4.1	WeChat: cstutoros	Contains $R_{x_1} \cap R_{x_2}$
2	3.4.2	$x[n - n_0]$	R_x , except for the possible addition or deletion of the origin or ∞
3	3.4.3	$z_0^n x[n]$	$ z_0 R_x$
4	3.4.4	$\frac{d}{dz} X(z)$	R_x
5	3.4.5	$x^*[n]$	R_x
6	3.4.6	$Re\{x[n]\}$	Contains R_x
7	3.4.7	$\frac{1}{2}[X(z) + X^*(z^*)]$	Contains R_x
8	3.4.6	$x^*[-n]$	$1/R_x$
9	3.4.7	$x_1[n] * x_2[n]$	Contains $R_{x_1} \cap R_{x_2}$

Method 1: Inspection Method

程序代写代做 CS编程辅导

Good idea to apply this to our memory:



$$z^{-m} \rightarrow \sigma[n-m], \quad m \text{ is a constant}$$

WeChat: cstutorcs

Assignment Project Exam Help

$$\frac{1}{1 - az^{-1}} \rightarrow \begin{cases} a^n u[n], & \text{ROC: } |z| > a \\ -a^n u[-n-1], & \text{ROC: } |z| < a \end{cases}$$

Email: tutorcs@163.com
 QQ: 749389476
<https://tutorcs.com>

Method 1: Inspection Method

Example

程序代写代做 CS编程辅导



$$a^n u[n] \quad \xleftrightarrow{\mathcal{Z}} \quad \frac{1}{1 - az^{-1}} \quad |z| > |a| \quad (\text{Table 3.1 Transform Pair 5})$$

WeChat: cstutorcs

Find the inverse z-transform of:
Assignment Project Exam Help

$$X(z) = \frac{1}{1 - \frac{1}{2}z^{-1}}$$

QQ: 749389476

<https://tutorcs.com>

Method 1: Inspection Method

Example

程序代写代做 CS编程辅导



$$a^n u[n] \quad \xleftrightarrow{\mathcal{Z}} \quad \frac{1}{1 - az^{-1}} \quad |z| > |a| \quad (\text{Table 3.1 Transform Pair 5})$$

WeChat: cstutorcs

Find the inverse z-transform of:
Assignment Project Exam Help

$$x(z) = \frac{1}{1 - \frac{1}{2}z^{-1}}, \quad \text{Email: tutorcs@163.com}$$

QQ: 749389476

Ans. According to Table 3.1 Transform Pair 5, if we set $a = \frac{1}{2}$,
<https://tutorcs.com>
 then:

$$x[n] = \left(\frac{1}{2}\right)^n u[n]$$

Method 2: Partial Fraction Expansion

程序代写代做 CS编程辅导

- Break down a rational function into a summation of smaller rational functions.
- Generic Rational Functions :

$$X(z) = \frac{\sum_{k=0}^M b_k z^{-k}}{\sum_{k=0}^N a_k z^{-k}} \quad (X(z) \text{ as ratio of polynomials in } z^{-1})$$

WeChat: cstutorcs

An equivalent expression is:

$$X(z) = \frac{\sum_{k=0}^M b_k z^{-k}}{\sum_{k=0}^N a_k z^{-k}} = \frac{b_0 + b_1 z^{-1} + b_2 z^{-2} + \cdots + b_M z^{-M}}{a_0 + a_1 z^{-1} + a_2 z^{-2} + \cdots + a_N z^{-N}}$$

Assignment Project Exam Help
Email: tutores@163.com

Factoring out z^{-M} from numerator and z^{-N} from denominator:

$$\begin{aligned} X(z) &= \frac{z^{-M}(b_0 z^M + b_1 z^{M-1} + b_2 z^{M-2} + \cdots + b_M)}{z^{-N}(a_0 z^N + a_1 z^{N-1} + a_2 z^{N-2} + \cdots + a_N)} \\ &= \frac{z^N \sum_{k=0}^M b_k z^{M-k}}{z^M \sum_{k=0}^N a_k z^{N-k}} \quad (X(z) \text{ as ratio of polynomials in } z) \end{aligned}$$

Method 2: Partial Fraction Expansion

程序代写代做 CS编程辅导



$$\frac{\sum_{k=0}^M b_k z^{M-k}}{z^M \sum_{k=0}^N a_k z^{N-k}}$$

WeChat: cstutorcs

How many poles and zeros are there?

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

程序代写代做 CS编程辅导



$$\frac{\sum_{k=0}^M b_k z^{M-k}}{z^M \sum_{k=0}^N a_k z^{N-k}}$$

WeChat: cstutorcs

How many poles and zeros are there?

Assignment Project Exam Help

- There are M zeros and N poles at non-zero locations,
assuming $a_k > 0$ and $b_k > 0$.

Email: tutorcs@163.com
QQ: 749389476

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

程序代写代做 CS编程辅导



$$\frac{\sum_{k=0}^M b_k z^{M-k}}{z^M \sum_{k=0}^N a_k z^{N-k}}$$

WeChat: cstutorcs

How many poles and zeros are there?

Assignment Project Exam Help

- ▶ There are M zeros and N poles at non-zero locations,
assuming $a_k > 0$ and $b_k > 0$.
Email: tutorcs@163.com
- ▶ If $M > N$, then there are $(M - N)^{\text{th}}$ order poles at $z = 0$.
QQ: [749389476](https://tutorcs.com)
- ▶ If $M < N$, then there are $(N - M)^{\text{th}}$ order zeros at $z = 0$.

Method 2: Partial Fraction Expansion

程序代写代做 CS编程辅导

To obtain the partial expansion, it is most convenient to express $X(z)$ in the form:



WeChat: cstutorcs

$$X(z) = \frac{b_0 \prod_{k=1}^M (1 - c_k z^{-1})}{a_0 \prod_{k=1}^N (1 - d_k z^{-1})}$$

Assignment Project Exam Help
 Email: tutorcs@163.com

QQ: 749389476

where c_k are the zeros of $X(z)$ and d_k are the poles of $X(z)$.

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

Case 1

程序代写代做 CS编程辅导

Given



$$\frac{\prod_{k=1}^M (1 - c_k z^{-1})}{\prod_{k=1}^N (1 - d_k z^{-1})}$$

Case 1: If $M < N$ and the poles are all 1st order, we can decompose $X(z)$ as:

$$\begin{aligned} X(z) &= \sum_{k=1}^N \frac{A_k}{1 - d_k z^{-1}} && \text{Assignment Project Exam Help} \\ &= \frac{A_1}{1 - d_1 z^{-1}} + \frac{A_2}{1 - d_2 z^{-1}} + \dots + \frac{A_m}{1 - d_m z^{-1}} + \dots + \frac{A_N}{1 - d_N z^{-1}} && \text{Email: tutorcs@163.com} \end{aligned}$$

Then $A_m = (1 - d_m z^{-1})X(z)|_{z^{-1}=d_m^{-1}}$

- Note: $\frac{1}{(1-d_k z^{-1})^2}$ will result in second-order poles at $z = d_k$.

Method 2: Partial Fraction Expansion

Case 1

程序代写代做 CS编程辅导

Proof:

Multiply all sides by $(1 - d_m z^{-1})$



$$(1 - d_m z^{-1}) X(z) \Big|_{z=d_m} = \frac{b_0 \prod_{k=1}^M (1 - c_k z^{-1})}{a_0 \prod_{k=1}^m (1 - d_k z^{-1}) \prod_{k=m+1}^N (1 - d_k z^{-1})} \Big|_{z=d_m}$$

Assignment Project Exam Help

$$\text{Email: } \text{futores@163.com} + \frac{A_2(1 - d_m z^{-1})}{1 - d_2 z^{-1}} \Big|_{z=d_m} + \dots$$

QQ: 749389476

$$\dots + \frac{A_m(1 - d_m z^{-1})}{1 - d_m z^{-1}} \Big|_{z=d_m} + \dots + \frac{A_N(1 - d_m z^{-1})}{1 - d_N z^{-1}} \Big|_{z=d_m}$$

$$= 0 + 0 + \dots + A_m + \dots + 0$$

Method 2: Partial Fraction Expansion

Case 1

程序代写代做 CS编程辅导



S.No.	Form of the rational function	Form of the partial fraction
1.	$\frac{px+q}{(x-a)(x-b)}$, $a \neq b$	$\frac{A}{x-a} + \frac{B}{x-b}$
2.	$\frac{px+q}{(x-a)^2}$	WeChat: cstutorcs $\frac{A}{x-a} + \frac{B}{(x-a)^2}$
3.	$\frac{px^2+qx+r}{(x-a)(x-b)(x-c)}$	Assignment Project Exam Help Email: tutorcs@163.com $\frac{A}{x-a} + \frac{B}{x-b} + \frac{C}{x-c}$
4.	$\frac{px^2+qx+r}{(x-a)^2(x-b)}$	QQ: 749389476 $\frac{A}{x-a} + \frac{B}{(x-a)^2} + \frac{C}{x-b}$
5.	$\frac{px^2+qx+r}{(x-a)(x^2+bx+c)}$	https://tutorcs.com $\frac{A}{x-a} + \frac{Bx+C}{x^2+bx+c}$, where $x^2 + bx + c$ cannot be factorised further

Method 2: Partial Fraction Expansion

Case 1: Example 3.9

Consider the z -transform



$$\frac{1}{-\frac{3}{4}z^{-1} + \frac{1}{8}z^{-2}}$$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

Case 1: Example 3.9

Consider the z -transform



$$\frac{1}{-\frac{3}{4}z^{-1} + \frac{1}{8}z^{-2}}$$

$$X(z) = \frac{1}{1 - \frac{3}{4}z^{-1} + \frac{1}{8}z^{-2}} = \underbrace{\frac{1}{(1 - \frac{3}{4}z^{-1})(1 - \frac{1}{2}z^{-1})}}_{\text{Partial fraction expansion}} = \underbrace{\frac{A_1}{1 - \frac{1}{4}z^{-1}}}_{\text{Assignment Project Exam Help}} + \underbrace{\frac{A_2}{1 - \frac{1}{2}z^{-1}}}_{(\text{Enterprise Optimization})}$$

Clearing denominators: $1 + A_1(1 - \frac{1}{2}z^{-1}) = A_2(1 - \frac{1}{4}z^{-1})$

Substituting $z^{-1} = 4$ gives $A_1 = -1$ and substituting $z^{-1} = 2$ gives $A_2 = 2$.

Another method: Solve equations $A_1 + A_2 = 1$ and $\frac{A_1}{2} + \frac{A_2}{4} = 0$

<https://tutorcs.com>

$$X(z) = \frac{-1}{1 - \frac{1}{4}z^{-1}} + \frac{2}{1 - \frac{1}{2}z^{-1}}$$

How do we find A_k in general? Use $A_m = (1 - d_m z^{-1}) X(z)|_{z^{-1}=d_m^{-1}}$

Method 2: Partial Fraction Expansion

Case 1: Example 3.9

$$\begin{aligned} X(z) &= \frac{1}{1 - \frac{1}{4}z^{-1} - \frac{1}{2}z^{-2}} = \frac{1}{(1 - \frac{1}{4}z^{-1})(1 - \frac{1}{2}z^{-1})} \\ &= \frac{A_1}{1 - \frac{1}{4}z^{-1}} + \frac{A_2}{1 - \frac{1}{2}z^{-1}} \end{aligned}$$

WeChat: csfutorcs
Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

Case 1: Example 3.9



$$X(z) = \frac{1}{1 - \frac{1}{4}z^{-1} - \frac{1}{2}z^{-2}} = \frac{1}{(1 - \frac{1}{4}z^{-1})(1 - \frac{1}{2}z^{-1})}$$

$$= \frac{A_1}{1 - \frac{1}{4}z^{-1}} + \frac{A_2}{1 - \frac{1}{2}z^{-1}}$$

WeChat: csfutorcs

To find A_1 corresponding to the pole at $z = \frac{1}{4}$ (or $z^{-1} = 4$),

$$\left(1 - \frac{1}{4}z^{-1}\right) X(z) = \left(\frac{A_1}{1 - \frac{1}{4}z^{-1}} + \frac{A_2}{1 - \frac{1}{2}z^{-1}} \right) \left(1 - \frac{1}{4}z^{-1}\right)$$

Assignment Project Exam Help
Email: tutorcs@163.com
QQ: 749389476
<https://tutorcs.com>

$$\frac{1}{1 - \frac{1}{2}z^{-1}} \Big|_{z^{-1}=4} = A_1 + \frac{A_2(1 - \frac{1}{4}z^{-1})}{1 - \frac{1}{2}z^{-1}} \Big|_{z^{-1}=4}$$

$$\therefore A_1 = -1$$

Method 2: Partial Fraction Expansion

Case 1: Example 3.9



$$\begin{aligned} X(z) &= \frac{1}{1 - \frac{1}{4}z^{-1} - \frac{1}{2}z^{-2}} = \frac{1}{(1 - \frac{1}{4}z^{-1})(1 - \frac{1}{2}z^{-1})} \\ &= \frac{A_1}{1 - \frac{1}{4}z^{-1}} + \frac{A_2}{1 - \frac{1}{2}z^{-1}} \end{aligned}$$

WeChat: csfutorcs
Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

Case 1: Example 3.9



$$X(z) = \frac{1}{1 - \frac{1}{4}z^{-1} - \frac{1}{2}z^{-2}} = \frac{1}{(1 - \frac{1}{4}z^{-1})(1 - \frac{1}{2}z^{-1})}$$

$$= \frac{A_1}{1 - \frac{1}{4}z^{-1}} + \frac{A_2}{1 - \frac{1}{2}z^{-1}}$$

WeChat: csfutorcs

To find A_2 corresponding to the pole at $z = \frac{1}{2}$ (or $z^{-1} = 2$)

$$\left(1 - \frac{1}{2}z^{-1}\right) X(z) = \left(\frac{A_1}{1 - \frac{1}{4}z^{-1}} + \frac{A_2}{1 - \frac{1}{2}z^{-1}} \right) \left(1 - \frac{1}{2}z^{-1}\right)$$

Assignment Project Exam Help
Email: tutorcs@163.com
QQ: 749389476
<https://tutorcs.com>

$$\frac{1}{1 - \frac{1}{4}z^{-1}} \Big|_{z^{-1}=2} \stackrel{\cong}{=} \frac{A_1(1 - \frac{1}{2}z^{-1})}{1 - \frac{1}{4}z^{-1}} \Big|_{z^{-1}=2} + A_2$$

$$\therefore A_2 = 2$$

Method 2: Partial Fraction Expansion

Case 1: Example 3.9



$$\begin{aligned} X(z) &= \frac{1}{1 - \frac{3}{4}z^{-1} + \frac{1}{8}z^{-2}} = \frac{1}{(1 - \frac{1}{4}z^{-1})(1 - \frac{1}{2}z^{-1})} \\ &= \frac{A_1}{1 - \frac{1}{4}z^{-1}} + \frac{A_2}{1 - \frac{1}{2}z^{-1}} \end{aligned}$$

Email: tutorcs@163.com

QQ: 749389476

$$\therefore X(z) = \frac{-1}{1 - \frac{1}{4}z^{-1}} + \frac{2}{1 - \frac{1}{2}z^{-1}}$$

Method 2: Partial Fraction Expansion

Case 1

程序代写代做 CS编程辅导



Note the z-transform

$$\frac{A_k}{1 - d_k z^{-1}} \rightarrow \begin{cases} A_k(d_k)^n u[n] & \text{ROC: } |z| > d_k \\ A_k(d_k)^n u[-n-1] & \text{ROC: } |z| < d_k \end{cases}$$

WeChat: tutorcs
Assignment Project Exam Help
Email: tutorcs@163.com

QQ: 749389476

ROC needs to be used to determine the corresponding sequence.

Method 2: Partial Fraction Expansion

Case 1: Example 3.9 (Completed)



$$X(z) = \frac{1}{1 - \frac{3}{4}z^{-1} + \frac{1}{8}} = \frac{-1}{\frac{1}{4}z^{-1}} + \frac{2}{1 - \frac{1}{2}z^{-1}} = \frac{z^2}{(z - \frac{1}{2})(z - \frac{1}{4})}$$

If further know that ROC is $|z| > \frac{1}{2}$, then we conclude that:


WeChat: cstutorcs

$$x[n] = \left(\frac{1}{4}\right)^n u[n] + 2\left(\frac{1}{2}\right)^n \text{Exam}[n] \text{Help}$$

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



Method 2: Partial Fraction Expansion

Exercise

程序代写代做 CS编程辅导



$$X(z) = \frac{1}{1 - \frac{3}{4}z^{-1} + \frac{1}{8}z^{-2}} = \frac{-1}{1 - \frac{1}{4}z^{-1}} + \frac{2}{1 - \frac{1}{2}z^{-1}}$$

WeChat: cstutorcs

If the ROC is $\frac{1}{4} < |z| < \frac{1}{2}$, what would be the sequence?

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

Exercise

程序代写代做 CS编程辅导



$$X(z) = \frac{1}{1 - \frac{3}{4}z^{-1} + \frac{1}{8}z^{-2}} = \frac{-1}{1 - \frac{1}{4}z^{-1}} + \frac{2}{1 - \frac{1}{2}z^{-1}}$$

WeChat: cstutorcs

If the ROC is $\frac{1}{4} < |z| < \frac{1}{2}$, what would be the sequence?

Assignment Project Exam Help
Email: tutorcs@163.com

Ans.

$$x[n] = \frac{Q(\frac{1}{4})^n}{4} u[n] - \frac{7}{2} \left(\frac{1}{2}\right)^n u[-n-1]$$

<https://tutorcs.com>

As an exercise if ROC is $|z| < \frac{1}{4}$, determine the corresponding sequence.

Method 2: Partial Fraction Expansion

Case 2 - First-order poles



$$X(z) = \frac{\prod_{k=1}^M (1 - c_k z^{-1})}{a_0 \prod_{k=1}^N (1 - d_k z^{-1})}$$

WeChat: cstutorcs

Assignment Project Exam Help

Case 2: If $M \geq N$, a polynomial of the order $M - N$ has to be added. If all poles are first order, then we will have:

QQ: 749389476

$$X(z) = \sum_{r=0}^{M-N} \frac{B_r}{1 - d_r z^{-1}} + \sum_{k=1}^N \frac{A_k}{1 - d_k z^{-1}}.$$

<https://Btutorcs.com>

Method 2: Partial Fraction Expansion

Case 2 - First-order poles



$$X(z) = \sum_{r=0}^M B_r z^{-r} + \sum_{k=1}^N \frac{A_k}{1 - d_k z^{-1}}.$$

WeChat: cstutorcs

Assignment Project Exam Help

- ▶ Coefficients B_r can be obtained by long division of the numerator of $X(z)$ by its denominator.
Email: tutors@163.com
QQ: 749389476
- ▶ Coefficients A_k are obtained as in the case of $M < N$ (by multiplying $X(z)$ by $1 - d_k z^{-1}$ and evaluating at $z = d_k$ or $z^{-1} = d_k^{-1}$).

Method 2: Partial Fraction Expansion

Case 2 - First-order poles



Note the Z transform

$$B_r z^{-r} \rightarrow B_r \delta[n - r]$$

WeChat: cstutorcs

and

Assignment Project Exam Help

$$\frac{A_k}{1 - d_k z^{-1}} \rightarrow \begin{cases} A_k (d_k)^n u[n] & \text{ROC: } |z| > d_k \\ \text{Email: } \text{tutorcs}@163.com \\ \text{QQ: } 749389476 & \text{ROC: } |z| < d_k \end{cases}$$

ROC needs to be used to determine the corresponding sequence.

Method 2: Partial Fraction Expansion

Case 2: Example 程序代写代做 CS编程辅导



$$\frac{+ 2z^{-1} + z^{-2}}{1 - \frac{3}{2}z^{-1} + \frac{1}{2}z^{-2}}$$

WeChat: cstutorcs

- From $X(z)$, $M = N$.
- We need to rearrange $X(z)$ to have the following form:

Email: tutorcs@163.com

$$X(z) \underset{\text{QQ: 749389476}}{\underset{M-N}{=}} \sum_{r=0}^N B_r z^{-r} + \sum_{k=1}^N \frac{A_k}{1 - d_k z^{-1}}.$$

<https://tutorcs.com>

- To find B_r we need to calculate the long division of $X(z)$, which gives quotient 2 with a remainder $(-1 + 5z^{-1})$.

Empty page

程序代写代做 CS编程辅导



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

Case 2: Example

程序代写代做 CS编程辅导



$$X(z) = 2 + \frac{1}{(1 - \frac{1}{2}z^{-1})(1 - z^{-1})} = 2 + \frac{A_1}{1 - \frac{1}{2}z^{-1}} + \frac{A_2}{1 - z^{-1}}$$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

Case 2: Example

程序代写代做 CS 编程辅导



$$X(z) = 2 + \frac{1}{\left(1 - \frac{1}{2}z^{-1}\right)} = 2 + \frac{A_1}{1 - \frac{1}{2}z^{-1}} + \frac{A_2}{1 - z^{-1}}$$

To find A_1 corresponding to the pole at $z = \frac{1}{2}$ (or $z^{-1} = 2$)

$$\left(1 - \frac{1}{2}z^{-1}\right) X(z) = \left(2 + \frac{A_1}{1 - \frac{1}{2}z^{-1}} + \frac{A_2}{1 - z^{-1}}\right) \left(1 - \frac{1}{2}z^{-1}\right)$$

$$\left[2\left(1 - \frac{1}{2}z^{-1}\right) + \frac{\text{QQ: } 549389476}{1 - z^{-1}}\right]_{z^{-1}=2} = 2\left(1 - \frac{1}{2}z^{-1}\right) \Big|_{z^{-1}=2} + A_1 + \frac{A_2(1 - \frac{1}{2}z^{-1})}{1 - z^{-1}} \Big|_{z^{-1}=2}$$

$$\therefore A_1 = -9$$

Method 2: Partial Fraction Expansion

Case 2: Example

程序代写代做 CS 编程辅导

$$X(z) = 2 + \frac{1}{(1 - \frac{1}{2}z^{-1})(1 - z^{-1})} = 2 + \frac{A_1}{1 - \frac{1}{2}z^{-1}} + \frac{A_2}{1 - z^{-1}}$$



To find A_2 corresponding to the pole at $z = 1$ (or $z^{-1} = 1$)

WeChat: cstutorcs

$$(1 - z^{-1}) X(z) = \left(2 + \frac{A_1}{1 - \frac{1}{2}z^{-1}} + \frac{A_2}{1 - z^{-1}} \right) (1 - z^{-1})$$

Email: tutorcs@163.com

$$\left[2(1 - z^{-1}) + \frac{-1 + 5z^{-1}}{1 - \frac{1}{2}z^{-1}} \right]_{z^{-1}=1} = 2(1 - z^{-1}) \Big|_{z^{-1}=1} +$$

$$\frac{A_1(1 - z^{-1})}{1 - \frac{1}{2}z^{-1}} \Big|_{z^{-1}=1} + A_2$$

$$\therefore A_2 = 8$$

Method 2: Partial Fraction Expansion

Case 2: Example

程序代写代做 CS编程辅导



$$X(z) = 2 + \frac{-9}{1 - \frac{1}{2}z^{-1}} + \frac{8}{1 - z^{-1}}$$

WeChat: cstutorcs

- If ROC $|z| > 1$: Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

Case 2: Example

程序代写代做 CS编程辅导



$$X(z) = 2 + \frac{-9}{1 - \frac{1}{2}z^{-1}} + \frac{8}{1 - z^{-1}}$$

WeChat: cstutorcs

- If ROC $|z| > 1$: $x[n] = 2\delta[n] - 9(\frac{1}{2})^n u[n] + 8u[n]$

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

Case 2: Example

程序代写代做 CS编程辅导



$$X(z) = 2 + \frac{-9}{1 - \frac{1}{2}z^{-1}} + \frac{8}{1 - z^{-1}}$$

WeChat: cstutorcs

- If ROC $|z| > 1$: $x[n] = 2\delta[n] - 9(\frac{1}{2})^n u[n] + 8u[n]$

Email: tutorcs@163.com

- If ROC $|z| < \frac{1}{2}$: QQ: 749389476

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

Case 2: Example

程序代写代做 CS编程辅导



$$X(z) = 2 + \frac{-9}{1 - \frac{1}{2}z^{-1}} + \frac{8}{1 - z^{-1}}$$

WeChat: cstutorcs

- If ROC $|z| > 1$: $x[n] = 2\delta[n] - 9(\frac{1}{2})^n u[n] + 8u[n]$

Email: tutorcs@163.com

- If ROC $|z| < \frac{1}{2}$: $x[n] = 2\delta[n] - 9(\frac{1}{2})^n u[-n-1] - 8u[-n-1]$

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

Case 2: Example

程序代写代做 CS编程辅导



$$X(z) = 2 + \frac{-9}{1 - \frac{1}{2}z^{-1}} + \frac{8}{1 - z^{-1}}$$

WeChat: cstutorcs

- If ROC $|z| > 1$: $x[n] = 2\delta[n] - 9(\frac{1}{2})^n u[n] + 8u[n]$

Email: tutorcs@163.com

- If ROC $|z| < \frac{1}{2}$: $x[n] = 2\delta[n] - 9(\frac{1}{2})^n u[-n-1] - 8u[-n-1]$

<https://tutorcs.com>

- If ROC $\frac{1}{2} < |z| < 1$:

Method 2: Partial Fraction Expansion

Case 2: Example

程序代写代做 CS编程辅导



$$X(z) = 2 + \frac{-9}{1 - \frac{1}{2}z^{-1}} + \frac{8}{1 - z^{-1}}$$

WeChat: cstutorcs

- If ROC $|z| > 1$: $x[n] = 2\delta[n] - 9(\frac{1}{2})^n u[n] + 8u[n]$

Email: tutorcs@163.com

- If ROC $|z| < \frac{1}{2}$: $x[n] = 2\delta[n] - 9(\frac{1}{2})^n u[-n-1] - 8u[-n-1]$

<https://tutorcs.com>

- If ROC $\frac{1}{2} < |z| < 1$: $x[n] = 2\delta[n] - 9(\frac{1}{2})^n u[n] - 8u[-n-1]$

Method 2: Partial Fraction Expansion

Generalisation

程序代写代做 CS编程辅导



How to determine whether partial fraction expansion is right-sided or left-sided for the ROC?

WeChat: cstutorcs

- To decide whether each term within $X(z)$, i.e.,

Assignment Project Exam Help

Email: tutores@163.com

QQ: 749389476

corresponds to $(d_k)^n u[n]$ or $-(d_k)^n u[-n - 1]$, we need to determine the relationship of d_k position with respect to a given ROC.

Method 2: Partial Fraction Expansion

Generalisation

程序代写代做 CS编程辅导



To decide whether each term $\frac{A_k}{1-d_k z^{-1}}$ within $X(z)$, i.e., $\frac{A_k}{1-d_k z^{-1}}$, is a right-sided or left-sided sequence in the time domain:

- ▶ Lets assume the ROC is of the form $r_R < |z| < r_L$.

WeChat: cstutorcs
Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

Generalisation

程序代写代做 CS编程辅导



To decide whether each term $\frac{A_k}{1-d_k z^{-1}}$ within $X(z)$, i.e., $\frac{A_k}{1-d_k z^{-1}}$, is a right-sided or left-sided sequence in the time domain:

- Lets assume the ROC is of the form $r_R < |z| < r_L$.

WeChat: cstutorcs
Assignment Project Exam Help

- If a pole at d_k is $d_k > r_R$ then this will correspond to a right-sided sequence $(d_k)^n u[n]$.

Email: tutorcs@163.com
QQ: 749389476

<https://tutorcs.com>

Method 2: Partial Fraction Expansion

Generalisation

程序代写代做 CS 编程辅导



To decide whether each term $\frac{A_k}{1-d_k z^{-1}}$ within $X(z)$, i.e., $\frac{A_k}{1-d_k z^{-1}}$, is a right-sided or left-sided sequence in the time domain:

- Lets assume the ROC is of the form $r_R < |z| < r_L$.

**WeChat: cstutorcs
Assignment Project Exam Help**

- If a pole at d_k is $d_k < r_R$ then this will correspond to a right-sided sequence $(d_k)^n u[n]$.

**Email: tutorcs@163.com
QQ: 749389476**

<https://tutorcs.com>

- If a pole at d_k is $d_k > r_L$ then this will correspond to a left-sided sequence $-(d_k)^n u[-n - 1]$.

Method 3: Power Series Expansion

程序代写代做 CS编程辅导

We can write a given sequence in the form of the z -transform definition:



$$X(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

WeChat: cstutorcs
 Assignment Project Exam Help
 Email: tutorcs@163.com

$$= \cdots + x[-2]z^2 + x[-1]z + x[0] + x[1]z^{-1} + x[2]z^{-2} + \cdots$$

QQ: 749389476

We can determine any particular value of the sequence by finding the coefficient of the appropriate power of z^{-1} .

Method 3: Power Series Expansion

Example

程序代写代做 CS编程辅导

$$X(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$


$$x[-2]z^2 + x[-1]z + x[0] + x[1]z^{-1} + \dots$$

Example: $X(z) = z^2 - 0.5z - 1 + 0.5z^{-1}$. Then, by inspection we can evaluate $X(z)$ to

WeChat: cstutorcs

Assignment Project Exam Help

$$\text{Email: } \text{tutorcs@163.com}$$

$$x[n] = \begin{cases} 1, & n = -2 \\ -0.5, & n = -1 \\ -1, & n = 0 \\ 0.5, & n = 1 \\ 0, & \text{otherwise} \end{cases}$$

QQ: 749389476

<https://tutorcs.com>

Therefore $x[n]$ can be written as a summation of delta functions:

$$x[n] = \delta[n+2] - 0.5\delta[n+1] - \delta[n] + 0.5\delta[n-1]$$

Method 3: Power Series Expansion

程序代写代做 CS编程辅导



- Sometimes, a given function can be converted to a power series using *Taylor Expansion*. See Example 3.12 in the textbook.
Taylor series expansion is NOT assessable in the exams.

WeChat: cstutorcs
Assignment Project Exam Help

- When $X(z)$ is a ratio of polynomials, a long division can be used to convert $X(z)$ to a power series.
Email: tutorcs@163.com
QQ: 749389476

<https://tutorcs.com>

Method 3: Power Series Expansion

Example

程序代写代做 CS编程辅导



If the ROC is $|z| > a$, expand $X(z)$ as:

$$X(z) = \frac{1}{1 - az^{-1}} = 1 + az^{-1} + a^2z^{-2} + a^3z^{-3} + \dots$$

**WeChat: cstutorcs
Assignment Project Exam Help**

Email: tutorcs@163.com

Then,

QQ: 749389476

$x[n] = a^n u[n]$
<https://tutorcs.com>

Method 3: Power Series Expansion

Example

程序代写代做 CS编程辅导



ROC is $|z| < a$ or $|a|^{-1} < |z| < \infty$, we expand $X(z)$ as:

$$X(z) = \frac{1}{1 - az^{-1}} = -a^{-1}z - a^{-2}z^2 - a^{-3}z^3 - \dots$$

**WeChat: cstutorcs
Assignment Project Exam Help**

Email: tutorcs@163.com

Then,

QQ: 749389476

$x[n] = -a^n u[-n - 1]$
<https://tutorcs.com>

A Final Example

程序代写代做 CS编程辅导



- ▶ The z-transform of $x[n]$ is

WeChat: cstutorcs

$$X(z) = \frac{1 - 2z^{-1}}{1 - \frac{1}{2}z^{-1}}$$

Assignment Project Exam Help

for $|z| > \frac{1}{2}$. Find $x[n]$ using any valid method.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

A Final Example (Continued)

程序代写代做 CS编程辅导



This is one approach and be used to solve this problem:

$$X(z) = \frac{1 - 2z^{-1}}{1 - \frac{1}{2}z^{-1}} = 4 + \frac{-3}{1 - \frac{1}{2}z^{-1}}$$

WeChat: cstutorcs

Assignment Project Exam Help

Since ROC is $|z| > \frac{1}{2}$, we have Email: tutorcs@163.com

QQ: 749389476
 $x[n] = 4\delta[n] - 3(\frac{1}{2})^n u[n]$
<https://tutorcs.com>

A Final Example (Continued)

程序代写代做 CS 编程辅导

Here is another approach that could be used to solve this problem:

$$X(z) = \frac{1}{1 - \frac{1}{2}z^{-1}} + \frac{-2z^{-1}}{1 - \frac{1}{2}z^{-1}}$$


For the first term we have WeChat: cstutorcs

Assignment Project Exam Help

$$x_1[n] = \left(\frac{1}{2}\right)^n u[n]$$

Email: tutorcs@163.com

Using the time shift property for the second term we have

QQ: 749389476

$$x_2[n] = -2\left(\frac{1}{2}\right)^{n-1} u[n-1] = -4\left(\frac{1}{2}\right)^n u[n-1]$$

Verify that:

$$x_1[n] + x_2[n] = 4\delta[n] - 3\left(\frac{1}{2}\right)^n u[n]$$

A Final Example (Continued)

程序代写代做 CS编程辅导

Proof of the second approach to solve this problem:



$$= \left(\frac{1}{2}\right)^n u[n]$$

Using the time shift property of the second term we have

$$x_2[n] = \frac{1}{2}^{n-1} u[n-1] - 4\left(\frac{1}{2}\right)^n u[n-1]$$

Note that for $n = 0$, we have

Assignment Project Exam Help

$$x_1[n] + x_2[n] = \left(\frac{1}{2}\right)^0 u[0] - 4\left(\frac{1}{2}\right)^0 u[0-1] = \left(\frac{1}{2}\right)^0 u[0] - 1 = 4 - 3 = 4\delta[n] - 3\left(\frac{1}{2}\right)^0 u[0]$$

Note that for $n \geq 1$ we have

QQ: 749389476

$$x_1[n] + x_2[n] = \left(\frac{1}{2}\right)^n u[n] - 4\left(\frac{1}{2}\right)^n u[n-1]$$

$$= \left(\frac{1}{2}\right)^n - 4\left(\frac{1}{2}\right)^n = -3\left(\frac{1}{2}\right)^n = 0 - 3\left(\frac{1}{2}\right)^n = 4\delta[n] - 3\left(\frac{1}{2}\right)^n u[n]$$

for $n < 0$, $x_1[n] + x_2[n] = 0$. So overall for all n , we have the result.

Homework

程序代写代做 CS编程辅导



WeChat: cstutorcs

1. Read and understand Chapter 3 of the textbook
2. Related Problems: Assignment, Project, Exam, Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>