

程序代写代做 CS编程辅导



Lecture - 4
WeChat: cstutors

Assignment Project Exam Help
Digital Signal Processing

Email: tutors@163.com

ENGN 4537/ 6537

QQ: 749389476

Semester 2, 2023
<https://tutors.com>

z-Transform: Definition (Revisit)

程序代写代做 CS编程辅导

The z-transform of



$$\sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

WeChat: cstutorcs

- ▶ z is considered to be a complex variable
 - ▶ $z = re^{j\omega}$, where r is the radius, and ω is the phase

Email: tutorcs@163.com

- ▶ Notation $\mathcal{Z}\{x[n]\} = X(z)$

<https://tutorcs.com>

- ▶ How can you evaluate the FT from the z-transform?
 - ▶ Evaluate the z-transform at $z = e^{j\omega}$ (*unit circle on complex z-plane*), to get the FT.

Region of Convergence (ROC)

程序代写代做 CS编程辅导

- For a given sequence, the set of z values for which the z-transform converges is called the *Region of Convergence (ROC)* of the z-transform.

- Condition on convergence is

WeChat: cstutorcs

$$\sum_{n=-\infty}^{\infty} |x[n]| |z|^{-n} < \infty$$

Email: tutorcs@163.com

- ROC of the z-transform consists of all values of z such that the above inequality holds.
- If for some value of z , say $z = z_1$, is in the ROC ($z_1 \in \text{ROC}$), then all values of z on the circle defined by $|z| = |z_1|$ will also be in the ROC.

QQ: 749389476

<https://tutorcs.com>

Poles and Zeros of z-Transform

程序代写代做 CS编程辅导



- Let the z-transform be presented inside ROC as

$$X(z) = \frac{P(z)}{Q(z)}$$

WeChat: cstutorcs

where $P(z)$ and $Q(z)$ are polynomials in z

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Poles and Zeros of z-Transform

程序代写代做 CS编程辅导



- Let the z-transform be presented inside ROC as

$$X(z) = \frac{P(z)}{Q(z)}$$

WeChat: cstutorcs

where $P(z)$ and $Q(z)$ are polynomials in z

Email: tutorcs@163.com

- The **zeros** are the roots of the numerator polynomial (i.e., $P(z) = 0$) and the **poles** are the roots of the denominator polynomial (i.e., $Q(z) \neq 0$)

QQ: 749389476

<https://tutorcs.com>

Reminder: Sum of a Finite Power Series: $S_N = \sum_{n=0}^N b^n$

程序代写代做 CS编程辅导



$$S_N = 1 + b + b^2 + b^3 + \dots + b^N \quad (1)$$

$$bS_N = b + b^2 + b^3 + \dots + b^N + b^{N+1} \quad (2)$$

By subtracting (2) from (1), we have,

$$(1 - b)S_N = 1 - b^{N+1}$$

$$S_N = \frac{1 - b^{N+1}}{1 - b} \quad \text{provided } b \neq 1$$

Reminder: Sum of a Infinite Power Series: $S_N = \sum_{n=0}^{\infty} b^n$

程序代写代做 CS编程辅导



Similarly, $S_{\infty} = \sum_{n=0}^{\infty} b^n$

$$S_{\infty} = 1 + b + b^2 + b^3 + \dots$$

WeChat: cstutorcs

$$bS_{\infty} = b + b^2 + b^3 + \dots$$

Assignment Project Exam Help

Email: tutorcs@163.com

$$S_{\infty} = \frac{1}{1-b} \quad \text{provided } |b| < 1$$

QQ: 749389476

<https://tutorcs.com>

Reminder: Sum of Power Series: $S = \sum_{n=N_1}^{N_2} b^n$

程序代写代做 CS编程辅导



$$S = b^{N_1} + b^{N_1+1} + \dots + b^{N_2} \quad (3)$$

$$bS = b^{N_1+1} + b^{N_1+2} + \dots + b^{N_2} + b^{N_2+1} \quad (4)$$

By subtracting (4) from (3), we have,

$$(1 - b)S = b^{N_1} - b^{N_2+1}$$

$$S = \frac{b^{N_1} - b^{N_2+1}}{1 - b} \quad \text{provided } b \neq 1$$

Example 3.1 Right-sided Exponential sequences

程序代写代做 CS编程辅导



$$= a^n u[n]$$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3.1 Right-sided Exponential sequences

程序代写代做 CS编程辅导



$$= a^n u[n]$$

Then:

$$X(z) = \sum_{n=-\infty}^{\infty} a^n u[n] z^{-n} = \sum_{n=0}^{\infty} a^n z^{-n}.$$

WeChat: cstutorcs

For convergence, we need the series absolutely summable, i.e.,

Assignment Project Exam Help

Email: tutors@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3.1 Right-sided Exponential sequences

程序代写代做 CS编程辅导



$$= a^n u[n]$$

Then:

$$X(z) = \sum_{n=-\infty}^{\infty} a^n u[n] z^{-n} = \sum_{n=0}^{\infty} a^n z^{-n}.$$

WeChat: cstutorcs

For convergence, we need the series absolutely summable, i.e.,

Assignment Project Exam Help

Email: tutorcs@163.com

$$\sum_{n=0}^{\infty} |az^{-1}|^n < \infty$$

QQ: 749389476

<https://tutorcs.com>

- ▶ Above condition holds if $|az^{-1}| < 1$. Thus, the ROC is the range of values for which $|az^{-1}| < 1$ or $|z| > |a|$. i.e., outside of the circle with radius $|a|$ in the z -plane.

Example 3.1: Right-sided Exponential sequences

程序代写代做 CS编程辅导

Inside the ROC, $|z| > 1$ series converges to



$$X(z) = \sum_{n=0}^{\infty} a^n z^{-n}$$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

$= \frac{1}{1 - az^{-1}}$ (Geometric series)

QQ: 749389476

$= \frac{z}{z - a}$
<https://tutorcs.com>

Example 3.1: Right-sided Exponential sequences

程序代写代做 CS编程辅导

- ▶ Thus, $x[n] = a^n u[n]$ has a closed form z-Transform:



$$\frac{z}{z - a} \quad \text{for } |z| > |a|$$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3.1: Right-sided Exponential sequences

程序代写代做 CS编程辅导

- ▶ Thus, $x[n] = a^n u[n]$ has a closed form z-Transform:



$$\frac{z}{z - a} \quad \text{for } |z| > |a|$$

- ▶ Recall: The **zeros** are the roots of the numerator, and the **poles** are the roots of the denominator.

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3.1: Right-sided Exponential sequences

程序代写代做 CS编程辅导

- Thus, $x[n] = a^n u[n]$ has a closed form z-Transform:



$$\frac{z}{z - a} \quad \text{for } |z| > |a|$$

- Recall: The **zeros** are the roots of the numerator, and the **poles** are the roots of the denominator.
- Note that there is a zero at $z = 0$ and a pole at $z = a$.
- If $|a| < 1$ then ROC contains the unit circle, and hence, the FT of the sequence converges to:

$$X(e^{j\omega}) = \frac{1}{1 - ae^{-j\omega}} \quad (\text{Table 2.3 Prop. 4})$$

Assignment Project Exam Help
Email: tutuorcs@163.com
QQ: 749389476
https://tutors.com

Example 3.1: Right-sided Exponential sequences

程序代写代做 CS编程辅导

- Thus, $x[n] = a^n u[n]$ has a closed form z-Transform:



$$\frac{z}{z - a} \quad \text{for } |z| > |a|$$

- Recall: The **zeros** are the roots of the numerator, and the **poles** are the roots of the denominator.
- Note that there is a zero at $z = 0$ and a pole at $z = a$.
- If $|a| < 1$ then ROC contains the unit circle, and hence, the FT of the sequence converges to:


$$X(e^{j\omega}) = \frac{1}{1 - ae^{-j\omega}} \quad (\text{Table 2.3 Prop. 4})$$

- However, if $|a| \geq 1$ then ROC does not contain the unit circle, and hence, the FT of the sequence does not converge.

Example 3.1: Right-sided Exponential sequences

程序代写代做 CS编程辅导

ROC and pole-zero plot for $|a| < 1$



The symbol 'o' denotes the zero and 'x' denotes the pole.

Example 3.2: Left-sided Exponential sequences

程序代写代做 CS编程辅导



$$x[n] = -a \cdot u[-n-1] = \begin{cases} -a^n & n \leq -1 \\ 0 & n > -1. \end{cases}$$

WeChat: cstutorcs

Since the sequence is nonzero for $n \leq -1$, this is a **left-sided sequence**.

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3.2: Left-sided Exponential sequences

程序代写代做 CS编程辅导



$$x[n] = -a^{-n} u[-n-1] = \begin{cases} -a^n & n \leq -1 \\ 0 & n > -1. \end{cases}$$

WeChat: cstutors

Since the sequence is nonzero for $n \leq -1$, this is a **left-sided sequence**. The z- transform:

Assignment Project Exam Help

Email: tutorms@163.com


$$X(z) = - \sum_{n=-\infty}^{\infty} a^n u[-n-1] z^{-n} = - \sum_{n=-\infty}^{-1} a^n z^{-n}$$

QQ: 749389476

<https://tutorms.com>

Example 3.2: Left-sided Exponential sequences

程序代写代做 CS编程辅导



$$X(z) = - \sum_{n=-\infty}^{-1} [a^{n-1}] z^{-n} = - \sum_{n=-\infty}^{-1} a^n z^{-n}$$

We need to change the sequence to start from zero and increase to infinity.

WeChat: tutorcs

Assignment Project Exam Help


Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3.2: Left-sided Exponential sequences

程序代写代做 CS编程辅导



$$X(z) = - \sum_{n=-\infty}^{-1} a^{n+1} z^{-n} = - \sum_{n=-\infty}^{-1} a^{n+1} z^{-n}$$

We need to change the sequence to start from zero and increase to infinity.

WeChat: tutorscs

Assignment Project Exam Help

Change of index $n = -m$, we have

Email: tutorscs@163.com

$$X(z) = - \sum_{m=\infty}^1 a^{-m} z^m = - \sum_{m=1}^{\infty} (a^{-1} z)^m = 1 - \sum_{m=0}^{\infty} (a^{-1} z)^m$$

QQ: 749389476

https://tutorscs.com

Example 3.2: Left-sided Exponential sequences

程序代写代做 CS编程辅导



The z-Transform:

WeChat: cstutorcs

$$X(z) = 1 - \sum_{m=0}^{\infty} (a^{-1}z)^m$$

Assignment Project Exam Help

The series converges if $|(a^{-1}z)| < 1$, i.e., $|z| < |a|$ which is the ROC.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3.2: Left-sided Exponential sequences

程序代写代做 CS编程辅导



Using the formula for sum of the geometric series

$$X(z) = 1 - \sum_{m=0}^{\infty} (a^{-1}z)^m = 1 - \frac{1}{1 - a^{-1}z} = 1 - \frac{a}{z - a} = \frac{z}{z - a}$$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

We can see that there is a zero at $z = 0$ and a pole at $z = a$.

QQ: 749389476

<https://tutorcs.com>

Example 3.2: Left-sided Exponential sequences

程序代写代做 CS编程辅导

The z-Transform: The sequence $x[n] = a^n u[-n-1]$ converges if $|(a^{-1}z)| < 1$, i.e., $|z| < |a|$. The z-transform is



$$X(z) = \frac{z}{z - a}$$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3.2: Left-sided Exponential sequences

程序代写代做 CS编程辅导

The z-Transform: The sequence converges if $|(a^{-1}z)| < 1$, i.e., $|z| < |a|$. The z-transform



$$X(z) = \frac{z}{z - a}$$

WeChat: cstutorcs

- Note that if $|a| \leq 1$ then the unit circle is not in ROC and the FT of the sequence does not exist.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3.2: Left-sided Exponential sequences

程序代写代做 CS编程辅导

The z-Transform: The sequence converges if $|a^{-1}z| < 1$, i.e., $|z| < |a|$. The z-transform



$$X(z) = \frac{z}{z - a}$$

WeChat: cstutorcs

- ▶ Note that if $|a| \leq 1$ then the unit circle is not in the ROC and the FT of the sequence does not exist.

Email: tutorcs@163.com

- ▶ But if $|a| > 1$ then the unit circle is in the ROC and the FT is

QQ: 749389476


<https://tutorcs.com>

$$X(e^{j\omega}) = \frac{1}{1 - ae^{-j\omega}}$$

Example 3.2: Left-sided Exponential sequences

程序代写代做 CS编程辅导

ROC and pole-zero $p < 1$



Example 3.2: Left-sided Exponential sequences

程序代写代做 CS编程辅导



- ▶ **Important:** Note that the algebraic expressions for Example 3.1 and 3.2 (right-sided and left-sided sequences) are the same. Also, their pole-zero locations are the same. However, the regions of convergence are different.
- ▶ Thus, the ROC needs to be found, in order to understand the convergence of each sequence.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3.3: Sum of Two exponential Series

$$x[n] = \left(\frac{1}{2}\right)^n u[n] + \left(-\frac{1}{3}\right)^n u[n]$$

The z-transform is



$$X(z) = \sum_{n=-\infty}^{\infty} \left(\left(\frac{1}{2}\right)^n u[n] + \left(-\frac{1}{3}\right)^n u[n] \right) z^{-n}$$

Assignment Project Exam Help

$$= \sum_{n=-\infty}^{\infty} \left(\frac{1}{2}\right)^n u[n] z^{-n} + \sum_{n=-\infty}^{\infty} \left(-\frac{1}{3}\right)^n u[n] z^{-n}$$

QQ: 749389476

$$= \sum_{n=0}^{\infty} \left(\frac{1}{2} z^{-1}\right)^n + \sum_{n=0}^{\infty} \left(-\frac{1}{3} z^{-1}\right)^n$$

Example 3.3: Sum of Two exponential Series

$$x[n] = \left(\frac{1}{2}\right)^n u[n] + \left(-\frac{1}{3}\right)^n u[n]$$

The z-transform is



$$X(z) = \sum_{n=0}^{\infty} \left(\frac{1}{2}z^{-1}\right)^n + \sum_{n=0}^{\infty} \left(-\frac{1}{3}z^{-1}\right)^n$$

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3.3: Sum of Two exponential Series

$$x[n] = \left(\frac{1}{2}\right)^n u[n] + \left(-\frac{1}{3}\right)^n u[n]$$

The z-transform is



$$X(z) = \underbrace{\sum_{n=0}^{\infty} \left(\frac{1}{2}z^{-1}\right)^n}_{\substack{|z^{-1}| < 1 \\ |z| > \frac{1}{2}}} + \underbrace{\sum_{n=0}^{\infty} \left(-\frac{1}{3}z^{-1}\right)^n}_{\substack{|z^{-1}| < 1 \\ |z| > \frac{1}{3}}}$$

Assignment Project Exam Help

Email: tutorcs@163.com

Convergence requires both summations to converge, i.e., $|z| > \frac{1}{2}$ and $|z| > \frac{1}{3}$.

<https://tutorcs.com>

Thus the ROC is the region of overlap, which is $|z| > \frac{1}{2}$.

Example 3.3: Sum of Two exponential Series

$$x[n] = \left(\frac{1}{2}\right)^n u[n] + \left(-\frac{1}{3}\right)^n u[n]$$



$$X(z) = \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^{-n} + \sum_{n=0}^{\infty} \left(-\frac{1}{3}\right)^n z^{-n}$$

WeChat: cstutorcs

$$= \frac{1}{1 - \frac{1}{2}z^{-1}} + \frac{1}{1 + \frac{1}{3}z^{-1}} \quad (\text{Geometric Series})$$

Email: tutorcs@163.com

$$= \frac{z}{z - \frac{1}{2}} + \frac{z}{z + \frac{1}{3}}$$

QQ: 749389476
<https://tutorcs.com>

$$= \frac{2z(z - \frac{1}{12})}{(z - \frac{1}{2})(z + \frac{1}{3})}$$

Example 3.3: Sum of Two exponential Series

$$x[n] = \left(\frac{1}{2}\right)^n u[n] + \left(-\frac{1}{3}\right)^n u[n]$$



What are the poles and zeros of $X(z) = \frac{2z(z - \frac{1}{12})}{(z - \frac{1}{2})(z + \frac{1}{3})}$?

WeChat: cstutorcs

Assignment Project Exam Help


Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3.3: Sum of Two exponential Series

$$x[n] = \left(\frac{1}{2}\right)^n u[n] + \left(-\frac{1}{3}\right)^n u[n]$$



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example 3.5: Two-sided Exponential Sequence

$$x[n] = \left(-\frac{1}{3}\right)^n u[n] + \left(\frac{1}{2}\right)^{-n} u[-n-1]$$



Verify this as an exercise

Figure 3.6 Pole-zero plot



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Homework

程序代写代做 CS编程辅导



1. Read and understand Chapter 3 (Sections 1 to 2) of the textbook
2. Related Problems: 3.1, 3.2, 3.3, 3.4, 3.5, 3.10, 3.19, 3.20, 3.54

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>