

程序代写代做 CS编程辅导



Lecture - 8  
WeChat: cstutors

Assignment Project Exam Help  
Digital Signal Processing

Email: [tutors@163.com](mailto:tutors@163.com)

ENGN 4537/ 6537

QQ: 749389476

Semester 2, 2023  
<https://tutors.com>

# Transform Analysis of Linear Time-Invariant (LTI) Systems

程序代写代做 CS编程辅导



We will study how to analyze a system given the Z-Transform and/or Fourier Transform

WeChat: cstutorcs

- ▶ Analysis of LTI systems described by Difference Equations
- ▶ Analysis of LTI systems described by Fourier Transform
- ▶ All Pass and Minimum Phase Systems

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Frequency Response of LTI Systems:

## Magnitude and Phase response

程序代写代做 CS编程辅导

- Frequency response of an LTI system is a complex gain that the system has for a complex exponential input  $e^{j\omega n}$ :



$$H(e^{j\omega}) = \sum_{n=-\infty}^{\infty} h[n]e^{-j\omega n}$$

Assignment Project Exam Help

- System input/output Fourier transforms are related by:

$$Y(e^{j\omega}) = H(e^{j\omega})X(e^{j\omega})$$

$$|Y(e^{j\omega})| = |H(e^{j\omega})||X(e^{j\omega})|$$

$$\angle Y(e^{j\omega}) = \angle H(e^{j\omega}) + \angle X(e^{j\omega})$$

$|H(e^{j\omega})|$  - Magnitude response

$\angle H(e^{j\omega})$  - Phase response

# Frequency Response: Magnitude

Magnitude response  $|H(e^{j\omega})|$  defines how different frequencies of the input signal gets by the system



程序代写代做 CS编程辅导



WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)



QQ: 749389476

<https://tutorcs.com>



# Frequency Response Magnitude:

## Influence of poles and zeros



An LTI system described by a difference equation has a system function

$$H(z) = \frac{\sum_{k=0}^M b_k z^{-k}}{\sum_{k=0}^N a_k z^{-k}} = \frac{b_0 \prod_{k=1}^M (1 - c_k z^{-1})}{a_0 \prod_{k=1}^N (1 - d_k z^{-1})}$$

If the system is stable, ROC includes the unit circle and will have a Fourier response

$$H(e^{j\omega}) = \frac{b_0 \prod_{k=1}^M (1 - c_k e^{-j\omega})}{a_0 \prod_{k=1}^N (1 - d_k e^{-j\omega})}$$

## Frequency Response Magnitude:

## Influence of poles and zeros



WeChat: cstutorcs

$$H(e^{j\omega}) = \frac{b_0 \prod_{k=1}^M (1 - c_k e^{-j\omega})}{a_0 \prod_{k=1}^N (1 - d_k e^{-j\omega})}$$

Assignment, Project, Exam Help

$$|H(e^{j\omega})| = \frac{|b_0| \prod_{k=1}^M |1 - c_k e^{-j\omega}|}{|a_0| \prod_{k=1}^N |1 - d_k e^{-j\omega}|} = \frac{|b_0| \prod_{k=1}^M |e^{j\omega} - c_k|}{|a_0| \prod_{k=1}^N |e^{j\omega} - d_k|}$$

Email: tutores@163.com

https://tutorcs.com

# Frequency Response Magnitude:

## Influence of poles and zeros

程序代写代做 CS编程辅导

$|H(e^{j\omega})|$  depends on the form  $|e^{j\omega} - a|$ . Therefore if we understand how that leaves at poles and zeros of the system, we can characterize and intuitively understand the magnitude response of the system.

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>



# Frequency Response Magnitude:

## Influence of poles and zeros



$$|H(e^{j\omega})| = \frac{|b_0| \prod_{k=1}^M |e^{j\omega} - c_k|}{|a_0| \prod_{k=1}^N |e^{j\omega} - d_k|}$$

WeChat: cstutors

$$|H(e^{j\omega})| = \frac{|b_0| \prod_{k=1}^M \text{"distance from } e^{j\omega} \text{ to zeros"} }{|a_0| \prod_{k=1}^N \text{"distance from } e^{j\omega} \text{ to poles"} }$$

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

- ▶ When  $e^{j\omega}$  is close to a zero,  $|H(e^{j\omega})|$  is small- zeros pull magnitude response down.
- ▶ When  $e^{j\omega}$  is close to a pole,  $|H(e^{j\omega})|$  is large - poles push magnitude response up.

QQ: 749389476

<https://tutorcs.com>




## Frequency Response Magnitude:

Influence of poles and zeros



Example 1:




WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)



QQ: 749389476

<https://tutorcs.com>

# Frequency Response Magnitude: Influence of poles and zeros

程序代写代做 CS编程辅导

Example 2:



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Summary: Given the poles and zeros of a system, we can infer the magnitude characteristics Fourier response.

# Frequency Response: Phase

程序代写代做 CS编程辅导

Given the Frequency Response  $H(e^{j\omega})$  of a system,  $\angle H(e^{j\omega})$  is its "Phase response".



WeChat: estutores

- ▶ Due to the periodicity of  $\omega$  the phase is not unique.
- ▶ We define phase in two ways, "principle value (wrapped phase)" and "continuous phase (unwrapped phase)".
- ▶ The principal value or wrapped phase is:

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

- ▶ The continuous phase can be represented in terms of the principal value as:

QQ: 749389476

<https://tutorcs.com>

$$\arg[H(e^{j\omega})] = \text{ARG}[H(e^{j\omega})] + 2\pi r(\omega)$$

where  $r(\omega)$  is an integer that is somewhat arbitrary.

# Frequency Response: Phase

程序代写代做 CS编程辅导


$$\arg[H(e^{j\omega})] = \text{ARG}[H(e^{j\omega})] + 2\pi r(\omega)$$



(a)

(a) The continuous phase exceeds the range  $-\pi$  to  $\pi$ .


WeChat: cstutorcs



(b)

(b) The principal value is restricted within the range.

Email: [tutorcs@163.com](mailto:tutorcs@163.com)



(c)

(c) Integer multiples of  $2\pi$  to be added to  $\text{ARG}[H(e^{j\omega})]$  to obtain  $\arg[H(e^{j\omega})]$ .

<https://tutorcs.com>

# Frequency Response Phase: Group Delay

程序代写代做 CS编程辅导

- ▶ Group delay is a useful representation of phase. It measures the [link between phase and predicts the system's delay to different frequency components](#):

WeChat: cstutorcs

Assignment Project Exam Help

$$\tau(\omega) = \text{grd}[H(e^{j\omega})] = -\frac{d}{d\omega} \{\arg[H(e^{j\omega})]\}$$

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

- ▶ Group delay is [represented by https://tutorcs.com](https://tutorcs.com)
- ▶ The units of group delay is samples.

# Frequency Response Phase: Group Delay

程序代写代做 CS编程辅导

Let us look at an ideal system:



$$h_{id}[n] = \delta[n - n_d]$$

WeChat: cstutorcs

- ▶ Frequency response

Assignment Project Exam Help

$$H_{id}(e^{j\omega}) = e^{-j\omega n_d}$$

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

$$\angle H_{id}(e^{j\omega}) = -\omega n_d$$

QQ: 749389476

- ▶ Time delay is associated with phase that is linear with frequency.

<https://tutorcs.com>

# Frequency Response Phase: Group Delay

程序代写代做 CS编程辅导

The ideal delay system



$$H_{id}(e^{j\omega}) = e^{-j\omega n_d},$$

where  $|H_{id}(e^{j\omega})| = 1$  and  $\angle H_{id}(e^{j\omega}) = -\omega n_d$ .

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

# Frequency Response Phase: Group Delay

程序代写代做 CS编程辅导

The ideal delay system



$$H_{id}(e^{j\omega}) = e^{-j\omega n_d},$$

where  $|H_{id}(e^{j\omega})| = 1$  and  $\angle H_{id}(e^{j\omega}) = -\omega n_d$ .

WeChat: cstutorcs

Assignment {Project} Exam Help

Email: tutorcs@163.com

- ▶ The group delay:  $\tau(\omega) = n_d$  samples
- ▶ This is a constant group delay. Phase is linear (perfect scenario).
- ▶ Group delay is independent of the frequency in an ideal delay system.

<https://tutorcs.com>



# Frequency Response Phase: Group Delay

程序代写代做 CS编程辅导

We use the group delay to measure the linearity of the phase.



WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

# Frequency Response Phase: Group Delay

程序代写代做 CS编程辅导

We use the group delay to measure the linearity of the phase.



- a broadband signal can be thought of as a superposition of narrowband signals

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

# Frequency Response Phase: Group Delay

程序代写代做 CS编程辅导

We use the group delay to measure the linearity of the phase.



- a broadband signal can be thought of as a superposition of narrowband signals

WeChat: cstutorcs

Assignment Project Exam Help

- If the group delay is constant with frequency, then each narrowband component undergoes the same time delay.

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

# Frequency Response Phase: Group Delay

程序代写代做 CS编程辅导

We use the group delay to measure the linearity of the phase.



- a broadband signal can be thought of as a superposition of narrowband signals

WeChat: cstutorcs

Assignment Project Exam Help

- If the group delay is constant with frequency, then each narrowband component undergoes the same time delay.

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476


- If group delay varies with frequency (is not constant), then each narrowband component undergoes different time delays - time dispersion of the output energy.

<https://tutorcs.com>

# Effects of Group Delay and Attenuation


程序代写代做 CS编程辅导

Consider an example system with the phase response shown below:



(a) Principle Value of Phase Response

- (a) The principal value phase response exhibits multiple discontinuities, due to the modulo  $2\pi$  computation of the wrapped phase.



(b) Unwrapped Phase Response

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476


<https://tutorcs.com>

- (b) The unwrapped (continuous) phase is smoother by removing the jumps of size  $2\pi$ .

- ▶ Is the phase linear?
- ▶ What does this mean in terms of group delay?

# Effects of Group Delay and Attenuation


程序代写代做 CS编程辅导



(a) Group delay of  $H(z)$

- (a) The group delay is not a constant, or the phase is not linear phase, negative at

**WeChat: cstutorcs**  $\omega = \pm 0.8\pi$  and a large positive peak at  $0.17\pi < |\omega| < 0.23\pi$ .



(b) Magnitude of Frequency Response

**Email: tutors@163.com**

**QQ: 749389476**


**https://tutorcs.com**

- (b) The magnitude response  $(|H(e^{j\omega})|)$ .

- ▶ What do the peaks mean in the group delay?
- ▶ What happens when the magnitude is zero?


# Effects of Group Delay and Attenuation

程序代写代做 CS编程辅导



(a) Waveform of signal  $x[n]$

(a) An input signal  $x[n]$  consisting of three narrowband pulses separated in time.



(b) Magnitude of DTFT of  $x[n]$

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>


(b) The corresponding DTFT magnitude  $|X(e^{j\omega})|$  shows the frequency components at  $0.2\pi$ ,  $0.4\pi$  and  $0.8\pi$ .

# Effects of Group Delay and Attenuation


System:

程序代写代做 CS编程辅导


Input:



(a) Group delay of  $H(z)$



(a) Waveform of signal  $x[n]$




(b) Magnitude of Frequency Response

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)




(b) Magnitude of DTFT of  $x[n]$

Output:

QQ: 749389476

<https://tutorcs.com>







The output signal  $y[n]$

- ▶  $\omega = 0.8\pi$  eliminated.
- ▶  $\omega = 0.2\pi$  pulse experience the most delay.



**Problem 5.34:**



程序代写代做 CS编程辅导

WeChat: cstutorcs





Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Determine which one of the following signals is the output of filter A when the input is  $x[n]$ :



# Homework

程序代写代做 CS编程辅导



WeChat: cstutorcs

- ▶ Read and understand Section 5.1 of the textbook.
- ▶ Related Problems Assignment, Project, Exam Help 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>