

# Assignment Project Exam Help

Unit Roots, Stochastic Trends, ARIMA Forecasting Models

~~<https://tutorcs.com>~~

Ka-fu WONG  
University of Hong Kong  
January 17, 2020

## WeChat: estutorcs

## CONTENTS

1. A quick review of autoregressive models	5
2. Random walk as a unit root	13
3. Drastic difference between stationary $AR(1)$ and unit root	17
4. Non-stationary of unit root process	20
5. Random walk with drift	24
6. ARMA with Unit Root	28
7. Similarity of $ARIMA(p, 1, q)$ to random walk	34
8. Application: Forecast of US GDP per capita trend	35
9. Detecting Unit Roots	40
9.1. Detecting unit root without drift	41
9.2. Detecting unit root <i>with drift</i>	44
9.3. Detecting unit root with drift (stochastic versus deterministic trend)	47
9.4. Detecting unit root with higher order autoregressive dynamics versus $AR(2)$	50

9.5. Detecting unit root with higher order autoregressive dynamics versus $AR(3)$	52
9.6. Detecting unit root with higher order autoregressive dynamics versus $AR(p)$	53
9.7. Detecting unit root with higher order autoregressive dynamics <i>with trend</i>	54
9.8. Detecting unit root with general ARMA models	56
10. Difference or not?	57
11. Forecasting the Yen/Dollar Exchange Rate	58
12. Level data	59
12.1. Preliminary data analysis	59
12.2. Box-Pierce Test of White Noise	63
12.3. Model selection	64
12.4. Residual check	67
12.5. Forecast	68
13. Augmented Dickey-Fuller Unit Root Test	71
14. Differenced Data	74

14.1.	Preliminary data analysis	74
14.2.	Box-Pierce Test of White Noise	78
14.3.	Model selection	79
14.4.	Residual check	82
14.5.	Forecast	83
15.	Concluding remarks:	86
16.	R packages used	87
16.1.	Package "forecast"	87
16.2.	Package "UnitRoots"	87

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

## 1. A QUICK REVIEW OF AUTOREGRESSIVE MODELS

Recall that the simple AR(1) model with zero mean:

$$(1.1) \quad y_t = \rho y_{t-1} + \epsilon_t \quad (\epsilon_t \sim WN(0, \sigma^2))$$

$$y_t - \rho y_{t-1} = \epsilon_t$$

$$(1 - \rho L)y_t = \epsilon_t$$

Stationarity requires the root of  $(1 - \rho z) = 0$  has absolute value larger than 1, i.e.,  $|\rho| < 1$ . With this condition, the Wold representation of  $y_t$  exists.

$$(1.2) \quad y_t = \sum_{i=0}^{\infty} \rho^i \epsilon_{t-i} = \epsilon_t + \rho \epsilon_{t-1} + \rho^2 \epsilon_{t-2} + \dots$$

Stationarity, as illustrated in this AR(1) process, has several important implications.

- (1) The impact of an innovation in time  $t$ ,  $\epsilon_t$ , on future  $y_t$ , i.e.,  $y_{t+h}$ , diminishes *towards zero* as the horizon  $h$  increases. Or, if  $y_t$  has a non-zero mean,  $y_{t+h}$  diminishes *towards the unconditional mean* of  $y_t$  as the horizon  $h$  increases.
- (2) The unconditional variance of  $y_t$  is finite.

$$\begin{aligned}
 V(y_t) &= V\left(\sum_{i=0}^{\infty} \rho^i \epsilon_{t-i}\right) = \sum_{i=0}^{\infty} V(\rho^i \epsilon_{t-i}) \\
 &= \sum_{i=0}^{\infty} \rho^{2i} V(\epsilon_t) = \sum_{i=0}^{\infty} \rho^{2i} \sigma^2 \\
 &= \frac{\sigma^2}{1 - \rho^2} < \infty
 \end{aligned}$$

A higher order AR process have similar properties. An AR(p) model with zero mean:

$$y_t = \rho_1 y_{t-1} + \rho_2 y_{t-2} + \dots + \rho_p y_{t-p} + \epsilon_t$$

$$(1 - \rho_1 L - \rho_2 L^2 - \dots - \rho_p L^p) y_t = \epsilon_t$$

Stationarity requires the roots of

$$(1 - \rho_1 x - \rho_2 x^2 - \dots - \rho_p x^p) = 0$$

to have absolute value (or modulus for complex roots) larger than 1.

With this condition, the Wold representation of  $y_t$  exists.

$$(1.3) \quad y_t = \sum_{i=0}^{\infty} b_i \epsilon_{t-i} = b_0 \epsilon_t + b_1 \epsilon_{t-1} + b_2 \epsilon_{t-2} + \dots$$

where  $b_0$  equals to 1 and  $b_i$  will be a function of the  $\rho_j$ 's.

Again, stationarity has several important implications.

- (1) The impact of an innovation in time  $t$ ,  $\epsilon_t$ , on future  $y_t$ , i.e.,  $y_{t+h}$ , diminishes *towards zero* as the horizon  $h$  increases. Or, if  $y_t$  has a non-zero mean,  $y_{t+h}$  diminishes *towards the unconditional mean* of  $y_t$  as the horizon  $h$  increases.

- (2) The unconditional variance of  $y_t$  is finite.


Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs



White Noise:  $y_t = \epsilon_t$ ,  $\epsilon_t \sim WN(0,1)$




Grey dotted line:  $y_t = \epsilon_t$ ,  $\epsilon_t \sim WN(0, 1)$

Blue solid line:  $y_t = 0.5y_{t-1} + \epsilon_t$ ,  $\epsilon_t \sim WN(0, 1)$

# Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs




Blue solid line:  $y_t = 0.5y_{t-1} + \epsilon_t$ ,  $\epsilon_t \sim WN(0, 1)$

Red solid line:  $y_t = 0.9y_{t-1} + \epsilon_t$ ,  $\epsilon_t \sim WN(0, 1)$

# Assignment Project Exam Help

<https://tutores.com>

WeChat: cstutores




Blue solid line:  $y_t = 0.9y_{t-1} + \epsilon_t$ ,  $\epsilon_t \sim WN(0, 1)$

Red solid line:  $y_t = 0.95y_{t-1} + \epsilon_t$ ,  $\epsilon_t \sim WN(0, 1)$

# Assignment Project Exam Help

<https://tutores.com>

WeChat: cstutores



## 2. RANDOM WALK AS A UNIT ROOT

Random walk has the following form:

$$(2.1) \quad y_t = y_{t-1} + \epsilon_t$$

The optimal forecast of  $y_{t+1}$  given time  $t$  information is

$$\begin{aligned} E\{y_{t+1} \mid y_t, y_{t-1}, \dots\} &= E\{y_t + \epsilon_{t+1} \mid y_t, y_{t-1}, \dots\} \\ &= E(y_t \mid y_t, y_{t-1}, \dots) + E(\epsilon_{t+1} \mid y_t, y_{t-1}, \dots) \\ &= y_t \end{aligned}$$

The optimal forecast of  $y_{t+2}$  given time  $t$  information is

$$\begin{aligned} E\{y_{t+2} \mid y_t, y_{t-1}, \dots\} &= E\{y_{t+1} + \epsilon_{t+2} \mid y_t, y_{t-1}, \dots\} \\ &= E(y_{t+1} \mid y_t, y_{t-1}, \dots) + E(\epsilon_{t+2} \mid y_t, y_{t-1}, \dots) \\ &= y_t \end{aligned}$$

Similarly, the optimal forecast of  $y_{t+h}$  given time  $t$  information is

$$E\{y_{t+h} \mid y_t, y_{t-1}, \dots\} = y_t.$$

Whatever happen in time  $t$  (a shock that changes  $y_t$ ) has *permanent effect* on future  $y_{t+h}$  and hence its forecast.

<https://tutorcs.com>

WeChat: cstutorcs

RW is considered as a naive and lazy forecasting model. Using RW as a forecasting model, there is no need to estimate any regression model before we make a forecast. As examples of short horizon forecast, we would use

- (1) today's closing as a forecast of tomorrow's Hang Seng Index closing.
- (2) today's yen-dollar rate as a forecast of tomorrow's yen-dollar exchange rate.
- (3) today's oil price as a forecast of tomorrow's oil price.
- (4) today's gold price as a forecast of tomorrow's gold price.
- (5) the last target federal funds rate as a forecast of the target federal funds rate after the next FOMC meeting.
- (6) past one year's GDP growth rate as a forecast of next year's GDP growth rate.
- (7) past one year's inflation rate as a forecast of next year's inflation rate.

Even if we may not consider RW naive in the short horizon forecast, it would definitely sound *naive* when we use the model to perform *long-horizon forecast*, say 10-year ahead.

Comparing equations (1.1) and (2.1), we can easily see that RW is like  $AR(1)$  with  $\rho = 1$ .

# Assignment Project Exam Help

Interpreting RW in the  $AR(1)$  framework, for RW the root of  $(1 - \rho x) = 0$  takes a value of 1, a *unit root*. That is why RW is also known as unit root.

<https://tutorcs.com>

The root ( $x$ ) in  $(1 - \rho x) = 0$  equals to "1"  $\iff \rho = 1$   
 (The root ( $x$ ) in  $(1 - \rho x) = 0$  equals to "-1"  $\iff \rho = -1$ )

WeChat: cstutorcs

So, one may think that RW is a limiting case of  $AR(1)$  when  $\rho$  approaches 1 and thus they should have *similar properties*.

That turns out to be *WRONG*!



### 3. DRASTIC DIFFERENCE BETWEEN STATIONARY AR(1) AND UNIT ROOT

Table 3.1 shows that the dependence of the forecast  $E(y_{t+h} | y_t, y_t - 1, \dots)$  on  $y_t$  at different horizons for AR(1) and unit root processes.

## Assignment Project Exam Help

TABLE 3.1. The forecast  $E(y_{t+h} | y_t, y_t - 1, \dots)$  at different horizons

	$\rho = 0.5$	$\rho = 0.9$	$\rho = 0.99$	$\rho = 0.999$	$\rho = 1$
$h = 1$	$0.5y_t$	$0.9y_t$	$0.99y_t$	$0.999y_t$	$y_t$
$h = 5$	$0.031y_t$	$0.59y_t$	$0.951y_t$	$0.995y_t$	$y_t$
$h = 10$	$0.001y_t$	$0.349y_t$	$0.904y_t$	$0.99y_t$	$y_t$
$h = 100$	$0.000$	$0.000$	$0.368y_t$	$0.905y_t$	$y_t$
$h = 1000$	$0.000$	$0.000$	$0.000$	$0.368y_t$	$y_t$
$h = 10000$	$0.000$	$0.000$	$0.000$	$0.000$	$y_t$

The difference becomes visible in the long horizons. As long as  $|\rho| < 1$ ,  $y_t$  *would not help forecast*  $y_{t+h}$  eventually, at *long horizons*.


Drastic difference between stationary AR(1) and unit root

// 17 ....

Blue solid line:  $y_t = 0.5y_{t-1} + \epsilon_t$ ,  $\epsilon_t \sim WN(0, 1)$

Red solid line:  $y_t = 0.9y_{t-1} + \epsilon_t$ ,  $\epsilon_t \sim WN(0, 1)$

Assignment Project Exam Help



Drastic difference between stationary AR(1) and unit root

// 18 ....

Blue solid line:  $y_t = 0.95y_{t-1} + \epsilon_t$ ,  $\epsilon_t \sim WN(0, 1)$

Red solid line:  $y_t = y_{t-1} + \epsilon_t$ ,  $\epsilon_t \sim WN(0, 1)$

Assignment Project Exam Help



WeChat: cstutorcs

## 4. NON-STATIONARY OF UNIT ROOT PROCESS

Given a random walk, we can use repeat substitutions to obtain

$$y_t = y_{t-1} + \epsilon_t$$

$$= y_{t-2} + \epsilon_{t-1} + \epsilon_t$$

$$= y_{t-3} + \epsilon_{t-2} + \epsilon_{t-1} + \epsilon_t$$

$$\vdots$$

$$= y_0 + \epsilon_1 + \epsilon_2 + \dots + \epsilon_{t-1} + \epsilon_t$$

$$= y_0 + \sum_{i=1}^t \epsilon_i$$

where  $y_0$  is the initial value of the process.

It is easy to see that

- (1)  $E(y_t) = y_0$ , i.e., the unconditional mean depends on the initial value of the process – even for large  $t$ .

Assignment Project Exam Help

$$E(y_t) = E\left(y_0 + \sum_{i=1}^t \epsilon_i\right)$$

<https://tutorcs.com>

$$= y_0 + \sum_{i=1}^t E(\epsilon_i)$$

WeChat: cstutorcs

(2)  $Var(y_t) = t\sigma^2$ , i.e., variance increases with time.

$$\begin{aligned} Var(y_t) &= Var\left(y_0 + \sum_{i=1}^t \epsilon_i\right) \\ &= Var\left(\sum_{i=1}^t \epsilon_i\right) \end{aligned}$$

$$= \sum_{i=1}^t Var(\epsilon_i) + \text{a bunch of covariances between } \epsilon_i \text{ and } \epsilon_j, i \neq j$$

$$= t\sigma^2 + 0$$

(3)  $\lim_{t \rightarrow \infty} Var(y_t) = \infty$ , i.e. the variance approaches infinity as  $t$  increases.

Clearly, a unit root process is *not covariance stationary*.

A unit root process is *not covariance stationary* because

- Unconditional mean depends on the initial condition/value.
- Unconditional variance depends on time, and approaches infinity as  $t$  increases

# Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

## 5. RANDOM WALK WITH DRIFT

By adding a constant term to the random walk process, we will have a random walk with **drift**.

Assignment Project Exam Help

$$y_t = \delta + y_{t-1} + \epsilon_t$$

↑  
drift

<https://tutorcs.com>

$$y_t = \delta + y_{t-1} + \epsilon_t$$

WeChat: cstutorcs

$$y_t - y_{t-1} = \delta + \epsilon_t$$

$$(1-L)y_t = \delta + \epsilon_t$$

$$\Delta y_t = \delta + \epsilon_t$$



Again we can use repeat substitution to obtain

$$y_1 = \delta + y_0 + \epsilon_1$$

$$y_2 = \delta + y_1 + \epsilon_2 = 2\delta + y_0 + \epsilon_2 + \epsilon_1$$

$$y_3 = \delta + y_2 + \epsilon_3 = 3\delta + y_0 + \epsilon_3 + \epsilon_2 + \epsilon_1$$

$$y_t = \delta + y_{t-1} + \epsilon_t = t\delta + y_0 + \epsilon_t + \epsilon_{t-1} + \dots + \epsilon_1$$

or

$$y_t = t\delta + y_0 + \sum_{i=1}^t \epsilon_i$$

Thus, the random walk with drift model implies a **linear time trend**,  $t\delta$ .

- This time trend ~~would not result if~~ we have  $y_t = \delta + \rho y_{t-1} + \epsilon_t$  and  $|\rho| < 1$ .
- This time trend is different from the conventional deterministic trend model such as  $y_t = \delta t + \epsilon_t$ .


To distinguish it from the deterministic trend model, random walk with drift is often called a model of **stochastic trend**.

Blue solid line:  $y_t = 0.95y_{t-1} + \epsilon_t$ ,  $\epsilon_t \sim WN(0, 1)$

Red solid line:  $y_t = y_{t-1} + \epsilon_t$ ,  $\epsilon_t \sim WN(0, 1)$

Purple solid line:  $y_t = 0.3 + y_{t-1} + \epsilon_t$ ,  $\epsilon_t \sim WN(0, 1)$

# Assignment Project Exam Help



It is easy to see that

(1)  $E(y_t) = t\delta + y_0$ , i.e., the unconditional mean depends on the initial value of the process.

(2)  $Var(y_t) = t\sigma^2$ , i.e., variance increases with time.

(3)  $\lim_{t \rightarrow \infty} Var(y_t) = \infty$ , i.e., the variance approaches infinity as  $t$  increases.

Thus, a *random walk with drift* process is *not covariance stationary*.

<https://tutorcs.com>

WeChat: cstutorcs

## 6. ARMA WITH UNIT ROOT

While a unit root in  $y_t$  is not stationary,  $\Delta y_t$ , the difference of  $y_t$  is *stationary*.

# Assignment Project Exam Help

$\Delta y_t$  is in fact ARMA(0,0).

Sometimes, we want to talk about the process of  $y_t$ , and we say  $y_t$  is ARIMA(0,1,0).

<https://tutorcs.com>

AR | MA  
0 1 0

The middle “integration” order “1” is used to tell readers that  $y_t$  has a unit root, and the “first” difference of  $y_t$  ( $\Delta y_t$ ) will be stationary.

Consider an AR(2) process

$$y_t = \rho_1 y_{t-1} + \rho_2 y_{t-2} + \epsilon_t$$

We can rewrite it as

$$(1 - \rho_1 L - \rho_2 L^2)y_t = \epsilon_t$$

# Assignment Project Exam Help

or

$$(1 - c_1 L)(1 - c_2 L)y_t = \epsilon_t$$

## <https://tutorcs.com>

Stationarity requires  $|c_1| < 1$  and  $|c_2| < 1$ , or the roots of  $(1 - c_1 x)(1 - c_2 x) = 0$  is larger than 1 in absolute values.

## WeChat: cstutorcs

If one of the roots is equal to 1, we have one unit root, say,  $c_1 = 1$ , we can rewrite

$$(1 - c_2 L)(1 - L)y_t = \epsilon_t$$

Assignment Project Exam Help

Thus,  $\Delta y_t$ , the difference of  $y_t$  is *stationary*. In fact,  $\Delta y_t$  is  $ARMA(1, 0)$ , or  $y_t$  is  $ARIMA(1, 1, 0)$ .

If  $y_t$  has two unit roots, i.e.,  $c_1 = 1$  and  $c_2 = 1$ , we have

$$(1 - L)(1 - L)y_t = \epsilon_t$$

$$\Delta \Delta y_t = \epsilon_t$$

or

WeChat: cstutorcs

$$\Delta^2 y_t = \epsilon_t$$

Thus,  $\Delta^2 y_t$ , the “second” difference of  $y_t$  is *stationary*. In fact,  $\Delta^2 y_t$  is  $ARMA(0, 0)$ , or  $y_t$  is  $ARIMA(0, 2, 0)$ . Again, the middle “integration” order “2” is used to tell readers that  $y_t$  has two unit roots.

Generally, consider an  $AR(p)$  process

$$y_t = \rho_1 y_{t-1} + \rho_2 y_{t-2} + \dots + \rho_p y_{t-p} + \epsilon_t$$

We can check stationarity by solving for the roots of

$$1 - \rho_1 x - \rho_2 x^2 - \dots - \rho_p x^p = 0$$

If there is exactly one unit root,  $\Delta y_t$ , the first difference of  $y_t$  is *stationary*. And,  $\Delta y_t$  is  $ARMA(p-1, 0)$  or  $y_t$  is  $ARIMA(p-1, 1, 0)$ .

If there is exactly two unit roots,  $\Delta^2 y_t$ , the second difference of  $y_t$  is *stationary*. And,  $\Delta^2 y_t$  is  $ARMA(p-2, 0)$ , or  $y_t$  is  $ARIMA(p-2, 2, 0)$ .

If there is exactly  $d$  unit roots,  $\Delta^d y_t$ , the  $d$ -th difference of  $y_t$  is *stationary*. And,  $\Delta^d y_t$  is  $ARMA(p-d, 0)$ , or  $y_t$  is  $ARIMA(p-d, d, 0)$ .

Remark:

Often, when the data is found non-stationary (say, by estimating a  $AR(p)$  and computing the roots), we will take the first difference of the data. If the first differenced data is found non-stationary, we will take an additional difference. If the second differenced data is found non-stationary, we will continue to take an additional difference.

That is, take  $n$ -th difference, until the  $n$ -th differenced data appears stationary.

<https://tutorcs.com>

WeChat: cstutorcs



It is easy to generalize the discussion to ARMA(p,q) models.

$$y_t = \rho_1 y_{t-1} + \rho_2 y_{t-2} + \dots + \rho_p y_{t-p} + \epsilon_t + \theta_1 \epsilon_{t-1} + \theta_2 \epsilon_{t-2} + \dots + \theta_q \epsilon_{t-q}$$

We can check stationarity by solving for the roots of

$$1 - \rho_1 x - \rho_2 x^2 - \dots - \rho_p x^p = 0$$

If there is exactly one unit root,  $\Delta y_t$ , the first difference of  $y_t$  is *stationary*. And,  $\Delta y_t$  is ARMA(p-1, q), or  $y_t$  is ARIMA(p-1, 1, q).

If there is exactly two unit roots,  $\Delta^2 y_t$ , the second difference of  $y_t$  is *stationary*. And,  $\Delta^2 y_t$  is ARMA(p-2, q), or  $y_t$  is ARIMA(p-2, 2, q).

If there is exactly  $d$  unit roots,  $\Delta^d y_t$ , the  $d$ -th difference of  $y_t$  is *stationary*. And,  $\Delta^d y_t$  is ARMA(p-d, q), or  $y_t$  is ARIMA(p-d, d, q).

## 7. SIMILARITY OF $ARIMA(p, 1, q)$ TO RANDOM WALK

$ARIMA(p, 1, q)$  processes are appropriately made stationary by differencing.

Shocks ( $\epsilon_t$ ) to  $ARIMA(p, 1, q)$  processes have *permanent* effects.

- Hence, shock persistence means that optimal forecasts even at very long horizons *do not completely revert to a mean or a trend*.

(Shocks ( $\epsilon_t$ ) to  $ARIMA(p, 0, q)$  processes have *transitory* effects.)

The variance of an  $ARIMA(p, 1, q)$  process grows without bound as time progresses.

- Uncertainty associated with our forecasts grows with horizon of our forecast.
- Width of our interval forecast grows *without bound* with the horizon of our forecast.


## 8. APPLICATION: FORECAST OF US GDP PER CAPITA TREND

- Estimation: 1859-1933 US GDP per capita  
(data from <http://www.measuringworth.com/usgdp/?q=hmit/gdp>)
- Forecast: 1934-1993 US GDP per capita
- Two types of models
  - $ARMA(p, q)$ : model selection criteria suggest  $p = 2$ ,  $q = 0$ , i.e.,  $ARMA(2, 0)$
  - $ARIMA(p, d, q)$ : model selection criteria suggest  $AR(1)$  in difference (i.e.,  $y_t - y_{t-1}$ ) with drift, i.e.,  $ARIMA(1, 1, 0)$

Note: The example is drawn from Diebold and Senhadji (1996).<sup>1</sup>

---


<sup>1</sup>Diebold, Francis X. and Abdelhak S. Senhadji (1996): "The Uncertain Unit Root in Real GDP: Comment," *American Economic Review*, 86: 1291-1298.




Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs







## 9. DETECTING UNIT ROOTS

Random walk may be viewed as a limiting case of  $AR(1)$ .

The Correlogram (ACF and PACF) looks like that of  $AR(1)$ , and thus cannot be used as a test of unit root.

Several statistical tests are available.

- Augmented Dickey–Fuller test
- Dickey–Fuller test
- Phillips–Perron test
- KPSS test
- Zivot–Andrews test

The most common ones are Dickey–Fuller (DF) test or augmented Dickey–Fuller (ADF) test.



### 9.1. Detecting unit root without drift.

A unit root process is a limiting case of  $AR(1)$ , which  $AR$  coefficient  $\rho = 1$ .

$y_t = \rho y_{t-1} + \epsilon_t$  versus  $y_t = y_{t-1} + \epsilon_t$   
 Assignment Project Exam Help

To test

$H_0$ :  $AR$  coefficient  $\rho = 1$  versus  $H_1$ :  $AR$  coefficient  $\rho < 1$

<https://tutorcs.com> (stationary alternative)

regress  $y_t$  on  $y_{t-1}$  and compute the test statistic in the usual way.

$\hat{\tau} = \frac{\hat{\rho} - 1}{s \sqrt{\frac{1}{T-2} \sum_{t=2}^T \hat{\epsilon}_t^2}}$   
 WeChat: [tutorcs](https://tutorcs.com)

where

$$s = \sqrt{\frac{\sum_{t=2}^T \hat{\epsilon}_t^2}{T-2}}$$

Sometimes, we would like to rewrite the model as

$$y_t - y_{t-1} = (\rho - 1)y_{t-1} + \epsilon_t$$

$$\Delta y_t = \beta y_{t-1} + \epsilon_t$$

To test **Assignment Project Exam Help**

$H_0$ : AR coefficient  $\rho = 1$       versus       $H_1$ : AR coefficient  $\rho < 1$

is almost equivalent to test **<https://tutorcs.com>**

$H_0: \beta = (\rho - 1) = 0$       versus       $H_1: \beta = (\rho - 1) < 0$

To conduct the test, regress  $\Delta y_t$  on  $y_{t-1}$  and compute the test statistic in the usual way. **WeChat: cstutors**

Almost all statistical softwares will compute *automatically* the corresponding  $t$ -statistic for us.

While the test looks like the usual  $t$ -test, we *cannot use the usual  $t$ -test critical values* for our test, because it turns out that the  $t$ -statistic

$$\hat{\tau} = \frac{\hat{\rho} - 1}{s \sqrt{\sum_{i=2}^n \frac{1}{q_i^2}}}$$

# Assignment Project Exam Help

does not have the usual Student- $t$  distribution. That is, the  $p$ -value spitted out automatically by the OLS regression procedure of usual statistical software is not to be trusted.

<https://tutorcs.com>

The statistics has shown to have a special distribution, called the *Dickey-Fuller* distribution. That is, we have to use critical values based on Dickey-Fuller distribution, not Student- $t$  distribution.

WeChat: cstutorcs

## 9.2. Detecting unit root *with drift*.

When the null is unit root with drift  $y_t = \alpha + y_{t-1} + \epsilon_t$ , it seems logical to consider the alternative of  $y_t = \alpha + \rho y_{t-1} + \epsilon_t$ . That is

**Assignment Project Exam Help**

We can rewrite the model as

$$y_t = \alpha + \rho y_{t-1} + \epsilon_t$$

$$y_t = \mu(1 - \rho) + \rho y_{t-1} + \epsilon_t$$

$$(y_t - \mu) = \rho(y_{t-1} - \mu) + \epsilon_t$$

Under null of unit root,  $\rho = 1$ , and *in addition*,  $\rho = 1$  implies  $\alpha = 0$ .  
(Note the inconsistency of unit root with drift?)

If  $\mu$  is known, we can simply subtract  $\mu$  from  $y_t$  and it reduces to the simple case without drift. We can then estimate

$$(y_t - \mu) = \rho(y_{t-1} - \mu) + \epsilon_t \text{ and test } H_0 : \rho = 1$$

Since  $\mu$  is not known, we have to estimate  $y_t = \alpha + \rho y_{t-1} + \epsilon_t$  and test  $H_0 : (\alpha, \rho) = (0, 1)$ .

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

$$y_t = \alpha + \rho y_{t-1} + \epsilon_t$$

# Assignment Project Exam Help

If we rewrite the model as

$$\begin{aligned} y_t - y_{t-1} &= \alpha + (\rho - 1)y_{t-1} + \epsilon_t \\ &= \alpha + \beta y_{t-1} + \epsilon_t \end{aligned}$$

we would be testing  $\beta = 0$  and  $\alpha = 0$ . Thus, some researchers will consider two possible null hypotheses in conducting unit root test.

- $H_0 : \beta = 0$
- $H_0 : \beta \neq 0, \alpha = 0$

Again, the test statistic will have a non-standard distribution. So, we have to refer to the Dickey-Fuller distribution.

## 9.3. Detecting unit root with drift (stochastic versus deterministic trend).

$y_t = \alpha + \beta TIME_t + \rho y_{t-1} + \epsilon_t$  versus  $y_t = \alpha + y_{t-1} + \epsilon_t$

# Assignment Project Exam Help

We can rewrite the model as

$$y_t = \alpha + \beta TIME_t + \rho y_{t-1} + \epsilon_t$$

$$y_t - (a + bTIME_t) = \rho [y_{t-1} - (a + bTIME_t)] + \epsilon_t$$

$$y_t - (a + bTIME_t) = \rho [y_{t-1} - (a + bTIME_{t-1})] - \rho b + \epsilon_t$$

Under null of unit root,  $\rho = 1$ .

If  $a$  and  $b$  are known, we can simply subtract  $(a + bTIME_t)$  from  $y_t$  and it reduces to the simple case without drift and trend.

Since  $a$  and  $b$  are not known, we have to estimate  $y_t = \alpha + \beta TIME_t + \rho y_{t-1} + \epsilon_t$  and test  $\rho = 1$ .

If we are rewrite the model as

$$y_t - y_{t-1} = \alpha + \beta TIME_t + (\rho - 1)y_{t-1} + \epsilon_t$$

we would be testing  $\gamma = 0$ .

Testing  $\gamma = 0$  is equivalent to testing

$$\begin{array}{cc} \text{Null} & \text{vs} & \text{Alternative} \\ y_t = \alpha + y_{t-1} + \epsilon_t & & y_t = \alpha + \beta TIME_t + \rho y_{t-1} + \epsilon_t, |\rho| < 1 \end{array}$$

Note that when the nonzero  $\alpha$  (drift) in the null specification is consistent with the time trend in data ( $y_t$ ).



Some researchers will consider the additional two hypotheses when a *deterministic trend* is included:

- $H_0 : \rho = 1, \alpha = 0$  (or  $H_0 : \gamma = (\rho - 1) = 0, \alpha = 0$ ) ,

Assignment Project Exam Help

Null vs Alternative

$$y_t = y_{t-1} + \epsilon_t \quad y_t = \alpha + \beta TIME_t + \rho y_{t-1} + \epsilon_t, |\rho| < 1$$

- and sometimes  $H_0 : \rho = 1, \alpha = 0, \beta = 0$  (or  $H_0 : \gamma = (\rho - 1) = 0, \alpha = 0, \beta = 0$ )

<https://tutorcs.com>

Null vs Alternative

$$y_t = y_{t-1} + \epsilon_t \quad y_t = \alpha + \beta TIME_t + \rho y_{t-1} + \epsilon_t, |\rho| < 1$$

(Note that if the null is  $y_t = y_{t-1} + \epsilon_t$ , we should not see a time trend in  $y_t$ .)

WeChat: cstutores

Again, the test statistic will have a non-standard distribution. So, we have to refer to the Dickey-Fuller distribution.

## 9.4. Detecting unit root with higher order autoregressive dynamics versus $AR(2)$ .

We can rewrite the  $AR(2)$

Assignment Project Exam Help  
as

$$y_t = -\rho_1 y_{t-1} - \rho_2 y_{t-2} + \epsilon_t$$

$$y_t = -(\rho_1 + \rho_2) y_{t-1} + \rho_2 (y_{t-1} - y_{t-2}) + \epsilon_t$$

$$y_t = \beta_1 y_{t-1} + \beta_2 (y_{t-1} - y_{t-2}) + \epsilon_t$$

where

$$\beta_1 = -(\rho_1 + \rho_2), \quad \beta_2 = \rho_2$$

To test for unit root, regress  $y_t$  on  $y_{t-1}$  and  $(y_{t-1} - y_{t-2})$ , and test the null of  $\beta_1 = 1$ , using the Dickey-Fuller distribution.

If we rewrite the model as

$$y_t - y_{t-1} = (\beta_1 - 1)y_{t-1} + \beta_2 (y_{t-1} - y_{t-2}) + \epsilon_t$$

$$= \gamma y_{t-1} + \beta_2 (y_{t-1} - y_{t-2}) + \epsilon_t$$

Assignment Project Exam Help

we would be testing  $\gamma = 0$ .

<https://tutorcs.com>

WeChat: cstutorcs

## 9.5. Detecting unit root with higher order autoregressive dynamics versus $AR(3)$ .

We can rewrite the  $AR(3)$

Assignment Project Exam Help

$$y_t = \beta_1 y_{t-1} + \beta_2 (y_{t-1} - y_{t-2}) + \beta_3 (y_{t-2} - y_{t-3}) + \epsilon_t$$

To test for unit root, regress  $y_t$  on  $y_{t-1}$ ,  $(y_{t-1} - y_{t-2})$  and  $(y_{t-2} - y_{t-3})$ , and test the null of  $\beta_1 = 1$ , using the Dickey-Fuller distribution.

If we rewrite the model as

$$\begin{aligned} y_t - y_{t-1} &= (\beta_1 - 1)y_{t-1} + \beta_2 (y_{t-1} - y_{t-2}) + \beta_3 (y_{t-2} - y_{t-3}) + \epsilon_t \\ &= \gamma y_{t-1} + \beta_2 (y_{t-1} - y_{t-2}) + \beta_3 (y_{t-2} - y_{t-3}) + \epsilon_t \end{aligned}$$

$$\Delta y_t = \gamma y_{t-1} + \beta_2 \Delta y_{t-1} + \beta_3 \Delta y_{t-2} + \epsilon_t$$

we would be testing  $\gamma = 0$ .

## 9.6. Detecting unit root with higher order autoregressive dynamics versus $AR(p)$ .

We can rewrite the  $AR(p)$

Assignment Project Exam Help  
as

$$y_t = \beta_1 y_{t-1} + \beta_2 (y_{t-1} - y_{t-2}) + \dots + \beta_p (y_{t-p+1} - y_{t-p}) + \epsilon_t$$

To test for unit root, regress  $y_t$  on  $y_{t-1}$ ,  $(y_{t-1} - y_{t-2})$  to  $(y_{t-p+1} - y_{t-p})$ , and test the null of  $\beta_1 = 1$ , using the Dickey-Fuller distribution.

If we rewrite the model as

$$\begin{aligned} y_t - y_{t-1} &= (\beta_1 - 1)y_{t-1} + \beta_2 (y_{t-1} - y_{t-2}) + \dots + \beta_p (y_{t-p+1} - y_{t-p}) + \epsilon_t \\ &= \gamma y_{t-1} + \beta_2 (y_{t-1} - y_{t-2}) + \dots + \beta_p (y_{t-p+1} - y_{t-p}) + \epsilon_t \\ \Delta y_t &= \gamma y_{t-1} + \beta_2 \Delta y_{t-1} + \dots + \beta_p \Delta y_{t-p+1} + \epsilon_t \end{aligned}$$

we would be testing  $\gamma = 0$ .

## 9.7. Detecting unit root with higher order autoregressive dynamics *with trend*.

We can rewrite

$$(y_t - a - bT) - \sum_{j=1}^p \rho_j (y_{t-j} - a - bT) = \epsilon_t$$

as

$$y_t = k_1 + k_2T + \beta y_{t-1} + \sum_{j=2}^p \beta_j (y_{t-j+1} - y_{t-j}) + \epsilon_t$$

where

$$k_1 = a \left( 1 + \sum_{i=1}^p \rho_i \right) - b \sum_{i=1}^p i \rho_i$$

and

$$k_2 = b \left( 1 + \sum_{i=1}^p \rho_i \right)$$

Under the null of unit root with drift, we would test  $\beta_1 = 1$ .

Under the null of unit root without drift, we would test

•  $H_0 : \beta_1 = 1, k_1 = 0$ , and  
 • sometimes  $H_0 : \beta_1 = 1, k_1 = 0, k_2 = 0$

Again, to test the null of  $\beta_1 = 1$ , and those  $k_1$  and  $k_2$ , using the Dickey-Fuller distribution.

If we rewrite the model as

$$y_t - y_{t-1} = k_1 + k_2 TIME_t + (\beta_1 - 1)y_{t-1} + \sum_{j=2}^p \beta_j (y_{t-j+1} - y_{t-j}) + \epsilon_t$$

$$\Delta y_t = k_1 + k_2 TIME_t + \gamma y_{t-1} + \sum_{j=2}^p \beta_j \Delta y_{t-j+1} + \epsilon_t$$

we would be testing  $\gamma = 0$ , or  $\gamma = 0, k_1 = 0, k_2 = 0$

### 9.8. Detecting unit root with general ARMA models.

- Complicated as any MA terms will map into infinite AR terms. But in practice we can include small number of AR terms as an approximation.
- As an *approximation*, *add many AR terms* and still use the standard Dickey-Fuller distribution.

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs



## 10. DIFFERENCE OR NOT?

In certain respects, the most important part of unit root theory for forecasting concerns estimation, not testing. It's important for forecasters to understand the effects of unit roots on consistency and small-sample bias. Such understanding on the one hand leads to the insight that at least asymptotically *we're probably better estimating forecasting models in levels with trends included* because then we'll get an accurate approximation to the dynamics in the data regardless of the true state of the world, unit root or no unit root. If there's no unit root, then of course it's desirable to work in levels; if there is a unit root, then the estimated long-run root will converge appropriately to unity, and at a fast rate. On the other hand, *differencing is appropriate only in the unit root case, and inappropriate differencing can be harmful, even asymptotically.*

(Francis Diebold, *Elements of Forecasting*)

## 11. FORECASTING THE YEN/DOLLAR EXCHANGE RATE

- $y$  = yen/dollar series
- Work with Levels:
  - $\ln(y)$  or
  - Differences:  $\Delta \ln(y)$
- Data: 1973.01 to 2014.12, *end-of-month* data (not monthly average), from <http://research.stlouisfed.org/fred2/series/DEXJPUS#>
- Estimation: 1973.01 – 2014.12
- Out-of-sample Forecast: 2013.01 – 2014.12

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

## 12. LEVEL DATA

### 12.1. Preliminary data analysis.

- First, we will do some preliminary data analysis.

- plot time series data
- ACF and PACF

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

FIGURE 12.1. Log JPY/USD Rate (1971-2012)




FIGURE 12.2. ACF of Log JPY/USD Rate



FIGURE 12.3. PACF of Log JPY/USD Rate



Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

12.2. **Box-Pierce Test of White Noise.** Obviously, if there series is white noise, we have nothing else to do but to report the non-forecastability.

Here, we adopt the simple Box-Pierce Test:

$$Q_{BP} = \sum_{k=1}^m \hat{\rho}_k^2 \sim \chi^2(m)$$

Setting  $m = 21 \approx \sqrt{480}$ , we obtain

- $\hat{Q}_{BP} = 82.12, 427, \text{ df} = 21, p\text{-value} < 2.2\text{e-}16$

The white noise null is obviously rejected at all conventional level of significance.

WeChat: cstutorcs

12.3. **Model selection.** We consider ARMA model with linear trend, with maximum AR order ( $p$ ) and MA order ( $q$ ) of 5.

# Assignment Project Exam Help

TABLE 12.1.  $p$ -value of Box Test

	$q = 0$	$q = 1$	$q = 2$	$q = 3$	$q = 4$	$q = 5$
$p = 0$	0	0	0	0	0	0
$p = 1$	0.0900	0.1608	0.2045	0.3023	0.3633	0.4252
$p = 2$	0.1707	0.2192	0.2449	0.3229	0.3732	0.5848
$p = 3$	0.2312	0.2393	0.6898	0.6879	0.3208	0.3601
$p = 4$	0.2663	0.2665	0.2589	0.4195	0.4102	0.7076
$p = 5$	0.2663	0.2685	0.7888	0.6905	0.2709	0.5736



TABLE 12.2. AIC and SIC of Log JPY/USD Rate

AIC	q = 0	q = 1	q = 2	q = 3	q = 4	q = 5
p = 0	-359.2032	-900.1638	-1229.3996	-1440.2734	-1548.9953	-1643.7729
p = 1	-1912.3581	-1911.4145	-1911.0634	-1910.9277	-1909.4812	-1908.0462
p = 2	-1911.5457	-1911.2376	-1910.1545	-1909.146	-1907.5734	-1907.5175
p = 3	-1911.3974	-1909.9885	-1914.6008	-1913.0301	-1905.2847	-1903.5307
p = 4	-1910.4738	-1908.4732	-1906.2927	-1915.9402	-1914.1297	-1912.9009
p = 5	-1908.4775	-1906.4953	-1911.3525	-1909.1295	-1903.9339	-1909.5526

SIC	q = 0	q = 1	q = 2	q = 3	q = 4	q = 5
p = 0	-340.6319	-883.4687	-1208.5306	-1415.2307	-1519.7788	-1610.3826
p = 1	-1895.6729	-1890.5451	-1886.0256	-1881.7112	-1876.0909	-1870.4821
p = 2	-1890.6767	-1886.1949	-1880.938	-1875.7557	-1870.0093	-1865.7796
p = 3	-1886.3547	-1880.772	-1881.2106	-1875.466	-1863.5469	-1857.6191
p = 4	-1881.2572	-1875.083	-1868.7286	-1874.2024	-1868.218	-1862.8155
p = 5	-1875.0872	-1868.9322	-1869.6147	-1863.2179	-1853.8485	-1855.2934

Box test suggests we cannot reject white noise residuals as long as the model include an AR term.

Conclusion from Box, AIC and SIC: AR(1) with trend.

# Assignment Project Exam Help


$$y_t = \beta_0 + \gamma_1 y_{t-1} + \beta_1 t + \epsilon_t$$

TABLE 12.3. Log JPY/USD Rate, best-fitting AR(1) with deterministic trend model

Variable	Coefficient	Standard Errors
Intercept	5.6357	0.1152
AR1	0.9791	0.0086
Trend	-0.0026	0.0004

12.4. **Residual check.** The residuals show no strong persistence.

FIGURE 12.4. Log JPY/USD Rate, best-fitting AR(1) with deterministic trend model, residual plot



12.5. **Forecast.** Note that the forecast exchange rate is decreasing with forecast horizon, essentially converging to the linear trend.

# Assignment Project Exam Help

FIGURE 12.3. Log (PY/USD) Rate, History and Forecast, AR(1) in levels with linear trend



FIGURE 12.6. Log JPY/USD Rate, History and Long-horizon Forecast, AR(1) in levels with linear trend





FIGURE 12.7. Log JPY/USD Rate, History, Forecast, and Realization, AR(1) in levels with linear trend



## 13. AUGMENTED DICKEY-FULLER UNIT ROOT TEST

Assignment Project Exam Help

$$\Delta y_t = \alpha + \gamma y_{t-1} + \sum_{j=1}^k \Delta y_{t-j} + \epsilon_t$$

<https://tutorcs.com>

$$\Delta y_t = \alpha + \gamma y_{t-1} + \beta t + \sum_{j=1}^k \Delta y_{t-j} + \epsilon_t$$

WeChat: cstutorcs

TABLE 13.1. Unit Root Test against the alternative of AR(1)

Variable	Coef.	Std.E.	Coef.	Std.E.	Coef.	Std.E.
Intercept			2.18E-02	1.87E-02	1.06E-01	5.09E-02
$y_{t-1}$	-5.49E-04	3.00E-04	-4.89E-03	3.74E-03	-1.95E-02	9.04E-03
Trend					-4.66E-05	2.62E-05
DF test						
t-stat	-1.8338		-1.3059		-2.159	
p-value	0.06761		0.5724		0.5108	

<https://tutorcs.com>  
WeChat: cstutorcs



TABLE 13.2. Unit Root Test against alternative of AR(2)

Variable	Coef.	Std.E.	Coef.	Std.E.	Coef.	Std.E.
Intercept			1.70E-02	1.86E-02	1.03E-01	5.69E-02
$y_{t-1}$	-4.70E-04	2.97E-04	-3.85E-03	3.71E-03	-1.97E-02	8.96E-03
$\Delta y_{t-1}$	3.92E-02	4.52E-02	4.06E-02	4.52E-02	4.90E-02	4.53E-02
Trend					-5.03E-05	2.60E-05
DF test						
t-stat	-1.5836		-1.0377		-2.1949	
p-value	0.1116		0.6726		0.4956	

## 14. DIFFERENCED DATA

### 14.1. Preliminary data analysis.

- First, we will do some preliminary data analysis.

- plot time series data
- ACF and PACF

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

FIGURE 14.1. Plot of Differenced Log JPY/USD Rate (1971-2012)




FIGURE 14.2. ACF of Differenced Log JPY/USD Rate




Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

FIGURE 14.3. PACF of Differenced Log JPY/USD Rate



14.2. **Box-Pierce Test of White Noise.** Here, we adopt the simple Box-Pierce Test on the first difference of log exchange rate:

$$Q_{BP} = \sum_{k=1}^m \hat{\eta}_k^2 \sim \chi^2(m)$$

Setting  $m = 21 \approx \sqrt{480}$ , we obtain

$$\hat{Q}_{BP} = 29.8071, \text{ df} = 21, p\text{-value} = 0.09593$$

The white noise null is rejected at 10% level of significance but not at 5% and 1%. This result suggests that we may choose ARIMA(0,1,0) model of the level data, or the ARMA(0,0) of the differenced data.

## 14.3. Model selection.

TABLE 14.1. p-value of Box Test (Differenced Log JPY/USD Rate)

	q = 0	q = 1	q = 2	q = 3	q = 4	q = 5
p = 0	0.0954	0.1511	0.1795	0.2326	0.2441	0.3609
p = 1	0.1574	0.1842	0.1945	0.1772	0.2865	0.5793
p = 2	0.1873	0.1761	0.0924	0.1488	0.1998	0.8029
p = 3	0.2049	0.2221	0.1552	0.1351	0.2187	0.1932
p = 4	0.2087	0.2986	0.2429	0.1652	0.1773	0.2379
p = 5	0.3536	0.508	0.8363	0.2547	0.2428	0.0497

TABLE 14.2. AIC and SIC of Differenced Log JPY/USD Rate

AIC	q = 0	q = 1	q = 2	q = 3	q = 4	q = 5
p = 0	-1910.1332	-1908.8196	-1908.0154	-1907.205	-1905.3261	-1904.5182
p = 1	-1908.8105	-1907.7687	-1906.5527	-1904.0224	-1901.7878	-1904.6594
p = 2	-1908.1568	-1908.0108	-1902.1975	-1900.9416	-1900.5459	-1907.0197
p = 3	-1906.7613	-1907.76	-1901.052	-1900.4727	-1904.151	-1899.698
p = 4	-1904.8048	-1903.9363	-1904.3559	-1900.1902	-1903.762	-1899.0594
p = 5	-1904.3389	-1904.0252	-1907.1	-1904.3722	-1899.1125	-1892.3261

SIC	q = 0	q = 1	q = 2	q = 3	q = 4	q = 5
p = 0	-1901.7857	-1896.2983	-1891.3202	-1886.3361	-1880.2834	-1875.3017
p = 1	-1896.3692	-1891.0731	-1885.6837	-1878.9797	-1874.5713	-1871.2791
p = 2	-1891.4617	-1887.1419	-1877.1548	-1871.7251	-1867.1556	-1869.4556
p = 3	-1885.8924	-1882.7173	-1871.8355	-1867.0825	-1866.587	-1857.9601
p = 4	-1879.7621	-1874.7198	-1870.9656	-1862.6262	-1862.0241	-1853.1477
p = 5	-1875.1724	-1870.6359	-1869.9359	-1862.6343	-1853.2008	-1842.2407



Box test suggests we cannot reject white noise residuals for all ARMA combination, including ARMA(0,0) on the differenced data.

Conclusion from Box, AIC and SIC: ARIMA(0,1,0).

# Assignment Project Exam Help

$$\Delta y_t = \alpha + \epsilon_t$$


TABLE 14.3 Regression results of Log JPY/USD Rate, ARIMA(0,1,0).

Variable	Coefficient	Standard Errors
Intercept	-0.0026	0.0015

14.4. **Residual check.** The residuals look like white noise and show no strong persistence.

# Assignment Project Exam Help

FIGURE 14.4 Log JPY/USD Rate, ARIMA(0,1,0) residual plot



14.5. **Forecast.** Note the wider forecast interval when compared with the one based on ARMA model of log exchange rate.

## Assignment Project Exam Help

FIGURE 14.1. Log JPY/USD Rate, History and Forecast,  $ARIMA(0,1,0)$



FIGURE 14.6. Log JPY/USD Rate, History and Long-horizon Forecast, ARIMA(0,1,0)

# Assignment Project Exam Help



FIGURE 14.7. Log JPY/USD Rate, History, Forecast, and Realization, ARIMA(0,1,0)

# Assignment Project Exam Help



Differenced Data

## 15. CONCLUDING REMARKS:

The two sets of forecasts based on the two different assumption (without and with unit roots) do not differ much. This is not suprising because if there is indeed unit root the AR(1) model should be able to approximate it well. And, indeed, the AR(1) coefficient is very close to one.

<https://tutorcs.com>

WeChat: cstutorcs

## 16. R PACKAGES USED

### 16.1. Package “forecast”.

```
Arima(x, order=c(0,0,0), seasonal=c(0,0,0), xreg=NULL,  
include.mean=TRUE, include.drift=FALSE, include.constant,  
lambda=model$lambda, transform.pars=TRUE, fixed=NULL,  
init=NULL, method=c("CSS-ML","ML","CSS"), n.cond,  
optim.control=list(), kappa=1e6, model=NULL)
```

```
forecast(object, h=felse(object$arima[5]>1, 2*object$arima[5], 10),  
level=c(80,95), fan=FALSE, xreg=NULL, lambda=object$lambda,  
bootstrap=FALSE, npaths=5000, ...)
```

### 16.2. Package “UnitRoots”

```
adfTest(x, lags = 1, type = c("nc", "c", "ct"), title = NULL,  
description = NULL)
```