



MONASH University

Information Technology

程序代写代做 CS编程辅导

FIT1006



Business Information Analysis

WeChat: cstutorcs

Assignment Project Exam Help

Lecture 18

Email: tutorcs@163.com

Hypothesis Testing (Part 2)

QQ: 749389476

<https://tutorcs.com>

Topics covered:

程序代写代做 CS编程辅导

- p-value

- Type I and II



- The significance and power of a test.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

The Steps of Hypothesis Testing

1. Decide on a null hypothesis H_0 .
2. Decide on an alternative hypothesis H_1 .
3. Decide on a significance level.
4. Calculate the appropriate test statistic.
5. Find from tables the corresponding tabulated test statistic.
6. Compare calculated and tabulated test statistics and decide whether to accept or reject the null hypothesis.
7. State the conclusion and assumptions of the test.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

(Source: Rees, D.G. Essential Statistics, Chapman and Hall 1995.)

程序代写代做 CS编程辅导

p-value

- The p-value of a test is the smallest value of the critical region that leads to the rejection of H_0 . For example, if a test has the following test statistic:


WeChat: cstutorcs

Assignment Project Exam Help (p = 2.685) = 0.004
i.e., $p = 0.004$

Email: tutores@163.com

QQ: 749389476

<https://tutorcs.com>



程序代写代做 CS编程辅导

Calculating the p-value

- Modified Question for next lecture:
- A hypothesis test for population mean when the population variance is known. (Note: unrealistic situation)
- The Axle manufacturing company has been making axles for a long time and kept records for every axle produced.
- Population parameters are $\mu = 90\text{mm}$ and $\sigma = 2.5\text{mm}$.
- A sample of 100 axles from new machine has **mean = 90.5**.
- Is the new machine making parts with same average length required (90mm) – **or are they longer than this?**
- **What is the p-value of the test?**



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

<https://flux.qa> 程序代写代做CS编程辅导 (Feed code: SJ6KGV)


Question 1

From the tables



2) =

$$P(Z > 2) = 1 - 0.9772 = 0.0228$$



This area = 0.9772; $z = 2.0$ (one tail)

A. 0.0114



B. 0.0228

C. 0.4886

WeChat: cstutorcs

D. 0.9772

Cumulative Probabilities for the Standard Normal Distribution

Assignment Project Exam Help

Table gives $P(Z < z)$ for $Z = N(0,1)$

Email: tutorcs@163.com

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952

QQ: 749389476

<https://tutorcs.com>

Solution

1. $H_0, \mu = 90\text{mm}$

2. $H_1, \mu > 90\text{mm}$

3. Significance = 0.01.

4. The test statistic, $x = 90.5$. We calculate Z_x .

(standardising) $Z = (90.5 - 90) / (2.5 / \sqrt{100}) = 2$.

5. From tables the calculated critical value is 2.33.

6. We see that $2 < 2.33$ and thus do not reject H_0 .

7. Conclude ...

8. p-value of the test is 0.0228 – this is the highest level of significance at which H_0 will be rejected.

程序代写代做 CS编程辅导 = 0.01,



Area = $1 - 0.01$
 $= 0.99$

One tail critical value
for $\alpha = 0.01$, from
table: $z = 2.33$


WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



$P(Z > 2) = 1 - 0.9772$
 $= 0.0228$

... visually...

程序代写代做 CS编程辅导

Interpreting the p-value:

- A small p-value indicates there is ample evidence to support the alternative hypothesis.
- A big p-value indicates that there is little evidence to support the alternative hypothesis.



Since, $2 < 2.33$

→ do not reject H_0

Test statistic = 90.5

Standardised test statistic = 2

WeChat: cstutorcs

Assignment Project Exam Help
 $P(Z > 2) = 1 - 0.9772 = 0.0228$

Email: tutorcs@163.com → p-value = 0.0228

Acceptance Region
QQ: 749389476

<https://tutorcs.com>

$\mu = 90$

For $\alpha = 0.01$, one tail;
critical value = 2.33

程序代写代做 CS编程辅导

Errors in Hypothesis Testing



- We can make two errors when testing hypotheses
 - A Type I error occurs when the null hypothesis is correct, but is rejected.
 - A Type II error occurs when the null hypothesis is incorrect but is not rejected.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

	H_0 is Not Rejected	H_0 is Rejected
H_0 is Correct	Correct Decision ($1 - \alpha$)	Type I Error α
H_1 is Correct	Type II Error β	Correct Decision ($1 - \beta$)

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

...

- The distinction between type I and II errors is easier to understand using a legal law analogy.



	Person Acquitted	Person Convicted
Person is Innocent	Correct Decision	Type I Error
Person is Guilty	Type II Error	Correct Decision

QQ: 749389476

- When we attempt to minimize the type I error, the consequence is an increase in the type II error.

The Power of the Test

程序代写代做 CS编程辅导

Rejecting the Null Hypothesis
when you should



- The probability of committing a Type II error is β . If the null hypothesis is incorrect, the probability of avoiding a Type II error is $(1-\beta)$, this is the power of the test.
- α and β are interrelated: increasing α reduces β and *vice versa*. To reduce both, sample size needs to be increased.
- Only α or β (usually α) can be chosen. The true value of β also depends on the actual value of the population parameter – which is often unknown.

WeChat: cstutorcs

Assignment Project Exam Help


Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

The Power of the Test ctd..



程序代写代做 CS编程辅导

Calculating β

■ Using the same ϵ



- The Axle manufacturing company has been making axles for a long time and kept records for every axle produced.
- Population parameters are $\mu = 90\text{mm}$ and $\sigma = 2.5\text{mm}$.
- A sample of 100 axles from new machine has mean = 92.7.
- Is the new machine making parts with same average length required (90mm)?
- Assume a 1% significance.
- What is the power of the test if it eventuates that the new machines produces axles with a mean length of 91mm?

WeChat: estutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Solution



- From our previous questions
- $H_0, \mu = 90\text{mm}$
- $H_1, \mu \neq 90\text{mm}$ (a two tailed experiment)
- Significance = 0.01.
- From tables the critical values are ± 2.58
- Use upper critical value since alternative is $> H_0$
- Calculate upper critical value = $90 + 2.58(0.25) = 90.65$
- $P(x < 90.65)$ when $X \sim N(91, 0.25^2)$ is $P(z < -1.42) = 0.08$
- This is the probability that we fail to reject the null hypothesis, when the population mean for the new axle is 91mm. The power of the test = 0.92

WeChat: cstutorcs

Assignment Project Exam Help


Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

The Power of the Test ctd..

程序代写代做 CS编程辅导



Power Curves

程序代写代做 CS编程辅导

- To see how the power of a test changes as the value of the alternative hypothesis changes, we can calculate the power for a range of values and view the resulting curve.


WeChat: cstutores

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutores.com>



Example

- From last lecture
- It is claimed that Melbourne families are spending more than \$150 per week on food and grocery items on average. A sample of 15 families was conducted and the amount spent each week was recorded. Do these results support this thesis? (assume a 1% significance)
- Weekly food and grocery expenditure (\$)
- 156, 234, 199, 78, 256, 189, 221, 49, 220, 178, 120, 290, 97, 177, 231.
- What is the power of the test if the average cost of groceries is \$160?



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

Summary Statistics from Excel



Mean	179.7
Standard Deviation	17.7
Median	189.0
Mode	#N/A
Standard Deviation	68.5
Sample Variance	4697.0
Kurtosis	10.5
Skewness	-0.5
Range	241.0
Minimum	49.0
Maximum	290.0
Sum	2695.0
Count	15.0

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

<https://flux.qa> (Feed code: SU0KGV)

Question 2

The power of the test is:



A. 0.99

B. 0.01

C. 0.98

D. 0.02

WeChat: cstutorcs

Assignment Project Exam Help

Hypothetical Distribution


Critical Value

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

'True' Distribution



<https://flux.qa> ^{程序代写代做CS编程辅导} (Feed code: SJ6KGV)

Question 3



For a one-sided t Statistic for a 1% significance is:

A. 2.6025

WeChat: [tutorcs](https://tutorcs.com)

B. 2.9467

✓ C. 2.6245

Assignment Project Exam Help

D. 2.9768

Email: tutorcs@163.com

Critical Values of the t Distribution

Table gives upper critical values only

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

n	α							
	0.300	0.200	0.100	0.050	0.025	0.010	0.005	
12	0.5386	0.8726	1.0832	1.3562	1.7823	2.1788	2.6810	
13	0.5375	0.8702	1.0795	1.3502	1.7709	2.1604	2.6503	
14	0.5366	0.8681	1.0763	1.3450	1.7613	2.1448	2.6245	
15	0.5357	0.8662	1.0735	1.3406	1.7531	2.1314	2.6025	
16	0.5350	0.8647	1.0711	1.3368	1.7459	2.1199	2.5835	
17	0.5344	0.8633	1.0690	1.3334	1.7396	2.1098	2.5669	
18	0.5338	0.8620	1.0672	1.3304	1.7341	2.1009	2.5524	

程序代写代做 CS编程辅导

...

- $H_0, \mu = \$150$

- $H_1, \mu > \$150$ (a c experiment)

- Significance = 0.01



- From tables the critical value $T_{(0.01)(v=14)}$ is $2.624\sigma_x$

WeChat: cstutorcs

- Calculate upper critical value: $150 + 2.624(17.7) = 196.5$

Assignment Project Exam Help

- Assume alternative distribution Normal, new μ , original SE.

Email: tutors@163.com

- $P(x < 196.5)$ when $X \sim N(160, 17.7) = P(z < 2.061) = 0.98$

QQ: 749389476

- This is β , the probability that we fail to reject the null hypothesis, when the population mean for expenditure is \$160.

<https://tutorcs.com>

- The power of the test is 0.02.

程序代写代做 CS编程辅导

Other Hypothesis Tests

- We have looked at hypothesis tests for the population mean and a population proportion.
- There are a range of other hypothesis tests which are regularly used, these include tests for:



- The difference of means
- Variance
- Distribution Shape
- Independence

WeChat: estutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

→ We're not going to cover this.

程序代写代做 CS编程辅导

Reading/Questions (Selvanathan)



- Reading: Hypothesis Testing
 - 7th Ed. Sections 12.3, 12.5, 12.6.

WeChat: cstutorcs

- Questions: Hypothesis Testing

Assignment Project Exam Help

- 7th Ed. Questions 12.1, 12.19, 12.25, 12.26, 12.56, 12.59, 12.65, 12.66, 12.67, 12.70, 12.72, 12.74.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>