



MONASH University

Information Technology

程序代写代做 CS编程辅导

FIT1006



Business Information Analysis

WeChat: cstutorcs

Assignment Project Exam Help

Lecture 11

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

Binomial and Poisson Distribution

QQ: 749389476

<https://tutorcs.com>

# Topics covered:

程序代写代做 CS编程辅导



- The Binomial Distribution
- The Poisson Distribution
- Poisson approximation to Binomial
- Probability calculations (manual & calculators)

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

# Motivating problem



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcscs@163.com

QQ: 749389476

<https://tutorcs.com>

- A cosmetics retailer sells 60 of a particular lipstick during the year. The shop is open 300 days per year. The shop keeps one lipstick on the counter and no others in stock. If the lipstick sells it can be replaced prior to opening the next trading day. However if a second or third... customer tries to buy the lipstick on any given day they lose the sale.
- What is the probability that the retailer loses a sale because they are out of stock?

<https://flux.qa> <sup>程序代写代做CS编程辅导</sup> (Feed code: SJ6KGV)

## Question 1 – ating Problem

What is the probability that the retailer loses a sale on a particular day because they are out of stock?

WeChat: cstutorcs

- A. 0.002      Assignment Project Exam Help
- ✓ B. 0.02      Email: [tutorcs@163.com](mailto:tutorcs@163.com)
- C. 0.2      QQ: 749389476
- D. 0.5      <https://tutorcs.com>
- E. None of these.

<https://flux.qa> <sup>程序代写 代做 CS 编程辅导</sup> (Feed code: SJ6KGV)

## Question 2



Which of the following situations could not be modelled using a Binomial distribution?

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

- A. A student guesses correct answers to a multiple choice test where each question has 4 choices.
- B. Students in class today indicate whether they are left or right-handed.
- ✓ C. I count the number of customers who visited Chadstone shopping centre today.
- D. I count the number of bad eggs in a carton of 12.

程序代写代做 CS编程辅导

# The Binomial Distribution



- We use the Binomial Distribution to determine the 'number of successes', each with a probability  $p$  of occurring in  $n$  independent trials.
- Typical questions:
  - A coin is tossed 50 times, what is the probability that 22 heads will be tossed? What is the probability of more than 40 heads being tossed?
  - A machine produces bolts and from past experience it is known that there is a 0.01 probability that a bolt produced will be defective. If a box contains 100 bolts, what is the probability that there are less than 5 defectives in the box.

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

# Formal Statement



- The use of the Binomial Distribution is valid under the following conditions:

1. Trials are independent
2. There are only two outcomes for each trial
3. The probability of success in each trial is constant

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

*For  $n$  independent trials, each with a probability*

*$p$ , of success, we define the the number of successes  $X$ ,*

*as a Binomial Distribution with the following formula :*

<https://tutorcs.com>

$$P(X = x) = {}^nC_x p^x (1 - p)^{(n-x)} \text{ for } x = 0, 1, 2, 3 \dots n$$

<https://flux.qa> <sup>程序代写代做CS编程辅导</sup> (Feed code: SJ6KGV)

## Question 3



Which of the following situations could not be modelled using a Binomial distribution?

WeChat: cstutores

A. A player tosses a coin 10 times and records the number of tails.

Assignment Project Exam Help  
Email: [tutorcs@163.com](mailto:tutorcs@163.com)

✓ B. The number of student Moodle access complaints each day is counted.

QQ: 749389476

C. Throwing 6 sixes (on a die) in a row.

<https://tutorcs.com>

D. Choosing 100 people at random and recording their gender.



# Example

程序代写代做 CS编程辅导

- A fair coin is tossed 10 times and the number of heads appearing uppermost is counted. What is the probability distribution for the number of heads thrown?



WeChat: cstutorcs


- Using table constructed in Excel:
- Probabilities for each x below.

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>



n =	10
p =	0.50
0	0.0010
1	0.0098
2	0.0439
3	0.1172
4	0.2051
5	0.2461
6	0.2051
7	0.1172
8	0.0439
9	0.0098
10	0.0010
sum =	1.0000

程序代写代做 CS编程辅导

# Calculating a Binomial Probability



- Example: A fair coin is tossed 12 times and the number of heads appearing uppermost is counted. What is the probability that 8 heads are thrown?

WeChat: cstutorcs

We have  $n = 12$ ,  $p = 0.5$  and  $x = 8$

thus Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

$$P(x) = {}^nC_x p^x (1-p)^{(n-x)}$$

QQ: 749389476

$$P(8) = {}^{12}C_8 0.5^8 (1-0.5)^{(12-8)}$$

<https://tutorcs.com>

$$= \frac{12 \times 11 \times 10 \times 9}{4 \times 3 \times 2 \times 1} \times 0.5^8 \times 0.5^4$$

$$= 0.1208$$

# Binomial Tables ( $n = 12$ )

程序代写代做 CS编程辅导



Recall:  $n = 12$ ,  $p = 0.5$  and  $x = 8$

Table gives  $P(X=x)$  for  $X = \text{Bi}(n,p)$

$n = 12$											
$p =$	0.05	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	0.95
0	0.5404	0.2824	0.0687	0.0138	0.0022	0.0002	0.0000	0.0000	0.0000	0.0000	0.0000
1	0.3413	0.3766	0.2062	0.0711	0.0199	0.0009	0.0003	0.0000	0.0000	0.0000	0.0000
2	0.0988	0.2301	0.2835	0.1678	0.0639	0.0161	0.0025	0.0002	0.0000	0.0000	0.0000
3	0.0173	0.0852	0.2362	0.2397	0.1419	0.0537	0.0125	0.0015	0.0001	0.0000	0.0000
4	0.0021	0.0213	0.1329	0.2311	0.2128	0.1208	0.0420	0.0078	0.0005	0.0000	0.0000
5	0.0002	0.0038	0.0532	0.1585	0.2270	0.1934	0.1009	0.0291	0.0033	0.0000	0.0000
6	0.0000	0.0005	0.0155	0.0492	0.1056	0.2256	0.1765	0.0792	0.0155	0.0005	0.0000
7	0.0000	0.0000	0.0033	0.0291	0.1009	0.1934	0.2270	0.1585	0.0532	0.0038	0.0002
8	0.0000	0.0000	0.0005	0.0078	0.0420	0.1208	0.2128	0.2311	0.1329	0.0213	0.0021
9	0.0000	0.0000	0.0001	0.0015	0.0125	0.0537	0.1419	0.2397	0.2362	0.0852	0.0173
10	0.0000	0.0000	0.0000	0.0002	0.0025	0.0161	0.0639	0.1678	0.2835	0.2301	0.0988
11	0.0000	0.0000	0.0000	0.0000	0.0003	0.0009	0.0174	0.0712	0.2062	0.3766	0.3413
12	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0022	0.0138	0.0687	0.2824	0.5404

WeChat: cstutors

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389406

<https://tutorcs.com>

程序代写代做CS编程辅导

# Cumulative Probabilities



- Example: A fair coin is tossed 12 times and the number of heads appearing uppermost is counted. What is the probability that up to and including 8 heads are thrown?

WeChat: cstutorcs

Assignment Project Exam Help

- Use the table of cumulative probabilities!

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

# Cumulative Binomial Table

Table gives  $P(X \leq x)$  for  $X = \text{Bi}(n, p)$



n =	12											
p =	0.05	0.10	0.20	0.40	0.50	0.60	0.70	0.80	0.90	0.95		
0	0.5404	0.2824	0.0687	0.0022	0.0002	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
1	0.8816	0.6590	0.2749	0.0850	0.0196	0.0032	0.0003	0.0000	0.0000	0.0000	0.0000	0.0000
2	0.9804	0.8891	0.5583	0.2538	0.0834	0.0193	0.0028	0.0002	0.0000	0.0000	0.0000	0.0000
3	0.9978	0.9744	0.7946	0.4925	0.2253	0.0730	0.0153	0.0017	0.0001	0.0000	0.0000	0.0000
4	0.9998	0.9957	0.9274	0.7237	0.4382	0.1938	0.0573	0.0095	0.0006	0.0000	0.0000	0.0000
5	1.0000	0.9995	0.9806	0.8812	0.6652	0.3812	0.1582	0.0386	0.0039	0.0001	0.0000	0.0000
6	1.0000	0.9999	0.9961	0.9614	0.8418	0.6128	0.3348	0.1178	0.0194	0.0005	0.0000	0.0000
7	1.0000	1.0000	0.9994	0.9905	0.9427	0.8663	0.5618	0.2763	0.0726	0.0043	0.0002	0.0000
8	1.0000	1.0000	0.9999	0.9983	0.9847	0.9270	0.7747	0.5075	0.2054	0.0256	0.0022	0.0000
9	1.0000	1.0000	1.0000	0.9998	0.9972	0.9807	0.9166	0.7472	0.4417	0.1109	0.0196	0.0000
10	1.0000	1.0000	1.0000	1.0000	0.9987	0.9968	0.9804	0.9150	0.7251	0.3410	0.1184	0.0000
11	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9978	0.9862	0.9313	0.7176	0.4596	0.0000
12	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

WeChat: cstutores

Assignment Project Exam Help

Email: tutores@163.com

QQ: 749389476

https://tutores.com

程序代写代做 CS编程辅导

# Binomial Probabilities in EXCEL



- We can use the functions in EXCEL to calculate binomial probabilities

- For  $X = Bi(n, p)$

$$P(X = x) = \text{BINOMDIST}(x, n, p, \text{false})$$

WeChat: cstutorcs  
Assignment Project Exam Help  
Probability of success

- The cumulative probability is given by

$$P(X \leq x) = \text{BINOMDIST}(x, n, p, \text{true})$$

Email: tutorcs@163.com  
QQ: 749389476

<https://tutorcs.com>

- ... or learn to use your calculator for these.

# Problem



- A farmer produces a range of eggs which are collected, graded and packaged into cartons of 12. The probability of a defective egg getting through the packaging process is 0.05. What is the probability that a carton will contain more than one defective egg.

WeChat: cstutorcs

- Now,  $n = 12$ ,  $p = 0.05$ ,

Assignment Project Exam Help

We want  $P(x \geq 1) = P(x=1) + P(x=2) + P(x=3) + \dots + P(x=12)$

Email: tutorcs@163.com

- A quicker way to use our knowledge of complementary events. \*

QQ: 749389476

<https://tutorcs.com>

- $P(x > 1) = 1 - [P(x = 0) + P(x = 1)]$

$$= 1 - 0.5404 - 0.3413 = 0.1183$$

<https://flux.qa> <sup>程序代写代做CS编程辅导</sup> (Feed code: SJ6KGV)

## Question 4



Which of the following situations could not be modelled using a Poisson distribution?

- 
- <sup>WeChat: cstutorcs</sup>
- <sup>Assignment Project Exam Help</sup>
- <sup>Email: tutorcs@163.com</sup>
- <sup>QQ: 749389476</sup>
- <sup><https://tutorcs.com></sup>
- A. The number of mutations in a strand of DNA.
  - ✓ B. The number of damaged books in a box of 50.
  - C. The number of earthquakes occurring in the Pacific Rim during 2013.
  - D. The number of hail stones landing on a 1m<sup>2</sup> area during a storm.



程序代写代做 CS编程辅导

# The Poisson Distribution



- We use the Poisson distribution to determine the number of occurrences of a random event, distributed over time or space.
- Typical Poisson Distribution questions:
  - On average 100 customers per day visit a particular shop. What is the probability that 10 customers enter the shop over one hour?  
WeChat: cstutorcs  
Assignment Project Exam Help  
Email: [tutorcs@163.com](mailto:tutorcs@163.com)
  - A fabric is known to have, on average, one defect per 10 meters. What is the probability that 5 meters of fabric will have two defects?  
QQ: 749389476  
<https://tutorcs.com>
  - We may also use the Poisson distribution as an approximation to the Binomial distribution.

# Formal Statement



- The use of the Poisson distribution is valid under the following conditions

1. Trials record the number of occurrences of a random event distributed over time or space.

2. The number of occurrences is theoretically unlimited

*For a random event occurring with an average rate  $\lambda$ , the mean number of occurrences over a period or area  $t$*

*is given by  $\mu = \lambda t$ . The number of occurrences over the time or period  $x$ , has a Poisson distribution with the formula :*

$$P(X = x) = \frac{e^{-\mu} \mu^x}{x!} \text{ for } x = 0, 1, 2, 3, \dots, \infty$$

# More Poisson examples

程序代写代做 CS编程辅导

From Wikipedia: [http://en.wikipedia.org/wiki/Poisson\\_distribution](http://en.wikipedia.org/wiki/Poisson_distribution)



- Electrical system example: calls arriving in a system.
- Astronomy example: photons arriving at a telescope.
- Biology example: the number of mutations on a strand of DNA per unit length.
- Management example: customers arriving at a counter or call centre.
- Civil engineering example: cars arriving at a traffic light.
- Finance and insurance example: number of losses or claims occurring in a given time.
- Radioactivity Example: Decay of a radioactive nucleus.
- The number of soldiers killed by horse-kicks each year in each corps in the Prussian cavalry.
- The number of yeast cells used when brewing Guinness beer. This example was made famous by William Sealy Gosset (1876–1937).
- The number of goals in sports involving two competing teams.
- The number of deaths per year in a given age group.
- The number of jumps in a stock price in a given time interval.

WeChat: estutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Example



- A certain fabric has a probability of 0.05 that a given metre will have a defect. Find the probability that a 10m length will have 3 defects?

We have  $t = 10$ ,  $\lambda = 0.05$  thus  $\mu = 0.5$

Assignment Project Exam Help

$$P(x) = \frac{e^{-\mu} \mu^x}{x!}$$

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

$$P(3) = \frac{e^{-0.5} \times 0.5^3}{3!}$$

<https://tutorcs.com>

$$= \frac{0.6065 \times 0.125}{6}$$

$$= 0.0126$$

# Poisson Tables

程序代写代做 CS编程辅导

Table gives  $P(X=x)$  for  $X = \text{Poi}(\mu)$

$\mu$	0.5	1.0	2.0	3.0	4.0	5.0	6.0	7.0	8.0	9.0	10.0
0	0.6065	0.3679	0.1353	0.0540	0.0183	0.0067	0.0025	0.0009	0.0003	0.0001	0.0000
1	0.3033	0.3679	0.2707	0.1490	0.0733	0.0337	0.0149	0.0064	0.0027	0.0011	0.0005
2	0.0758	0.1839	0.2707	0.2240	0.1465	0.0842	0.0446	0.0223	0.0107	0.0050	0.0023
3	0.0126	0.0613	0.1804	0.2240	0.1954	0.1404	0.0892	0.0521	0.0286	0.0150	0.0076
4	0.0016	0.0153	0.0902	0.1880	0.1954	0.1755	0.1339	0.0912	0.0573	0.0337	0.0189
5	0.0002	0.0031	0.0361	0.1008	0.1563	0.1755	0.1606	0.1277	0.0916	0.0607	0.0378
6	0.0000	0.0005	0.0120	0.0504	0.1042	0.1462	0.1606	0.1490	0.1221	0.0911	0.0631
7	0.0000	0.0001	0.0034	0.0216	0.0595	0.1044	0.1377	0.1490	0.1396	0.1171	0.0901
8	0.0000	0.0000	0.0009	0.0081	0.0298	0.0653	0.1033	0.1304	0.1396	0.1318	0.1126
9	0.0000	0.0000	0.0002	0.0027	0.0132	0.0363	0.0688	0.1014	0.1241	0.1318	0.1251
10	0.0000	0.0000	0.0000	0.0008	0.0063	0.0181	0.0413	0.0710	0.0993	0.1186	0.1251
11	0.0000	0.0000	0.0000	0.0002	0.0019	0.0082	0.0225	0.0452	0.0722	0.0970	0.1137
12	0.0000	0.0000	0.0000	0.0001	0.0006	0.0034	0.0113	0.0263	0.0481	0.0728	0.0948
13	0.0000	0.0000	0.0000	0.0000	0.0002	0.0013	0.0052	0.0142	0.0296	0.0504	0.0729
14	0.0000	0.0000	0.0000	0.0000	0.0001	0.0005	0.0022	0.0071	0.0169	0.0324	0.0521
15	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0009	0.0033	0.0090	0.0194	0.0347
16	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0014	0.0045	0.0109	0.0217
17	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0006	0.0021	0.0058	0.0128
18	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0009	0.0029	0.0071
19	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0004	0.0014	0.0037
20	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0006	0.0019

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcsl63.com

QQ: 749389476

https://tutorcs.com

程序代写代做 CS编程辅导

# Cumulative Probabilities



- Customers enter at an average rate of one per minute. What is the probability that in a 10 minute period more than 7 customers will enter?

- Now,  $\lambda = 1$ ,  $t = 10$ , and  $\mu = 10$

- We want  $P(x > 7)$

- $P(x > 7) = 1 - P(x = 7) - P(x = 6) - P(x = 0)$

QQ: 749389476

- (we can use the table of cumulative probabilities here)

- $P(x > 7) = 1 - 0.2202 = 0.7798$

<https://tutorcs.com>

程序代写代做 CS编程辅导

# Cumulative Poisson Tables

Table gives  $P(X \leq x)$  for  $X = \text{Poi}(\mu)$

mu	0.5	1.0	2.0	3.0	4.0	5.0	6.0	7.0	8.0	9.0	10.0
0	0.6065	0.3679	0.1353	0.0540	0.0183	0.0067	0.0025	0.0009	0.0003	0.0001	0.0000
1	0.9098	0.7358	0.4060	0.2466	0.1491	0.0916	0.0404	0.0174	0.0073	0.0030	0.0012
2	0.9856	0.9197	0.6767	0.4232	0.2381	0.1247	0.0620	0.0296	0.0138	0.0062	0.0028
3	0.9982	0.9810	0.8571	0.6472	0.4335	0.2650	0.1512	0.0818	0.0424	0.0212	0.0103
4	0.9998	0.9963	0.9473	0.8153	0.6288	0.4405	0.2851	0.1730	0.0996	0.0550	0.0293
5	1.0000	0.9994	0.9834	0.9161	0.7851	0.6160	0.4457	0.3007	0.1912	0.1157	0.0671
6	1.0000	0.9999	0.9955	0.9665	0.8893	0.7622	0.6063	0.4497	0.3134	0.2068	0.1301
7	1.0000	1.0000	0.9984	0.9881	0.9489	0.8666	0.7440	0.5987	0.4320	0.3239	0.2202
8	1.0000	1.0000	0.9998	0.9962	0.9786	0.9319	0.8472	0.7291	0.5925	0.4557	0.3328
9	1.0000	1.0000	1.0000	0.9989	0.9919	0.9682	0.9161	0.8305	0.7166	0.5874	0.4579
10	1.0000	1.0000	1.0000	0.9997	0.9970	0.9863	0.9574	0.9015	0.8159	0.7060	0.5830
11	1.0000	1.0000	1.0000	0.9999	0.9991	0.9945	0.9799	0.9467	0.8881	0.8030	0.6968
12	1.0000	1.0000	1.0000	1.0000	0.9997	0.9980	0.9912	0.9730	0.9362	0.8758	0.7916
13	1.0000	1.0000	1.0000	1.0000	0.9999	0.9993	0.9964	0.9872	0.9658	0.9261	0.8645
14	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9986	0.9943	0.9827	0.9585	0.9165
15	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9995	0.9976	0.9918	0.9780	0.9513
16	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9998	0.9990	0.9963	0.9889	0.9730
17	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9996	0.9984	0.9947	0.9857
18	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9993	0.9976	0.9928
19	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9997	0.9989	0.9965
20	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	0.9999	0.9996	0.9984

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

https://tutorcs.com

程序代写代做 CS编程辅导

# Poisson Probabilities in EXCEL

- We can use the functions in EXCEL to calculate Poisson probabilities



- For  $X = Poi(\mu)$  WeChat: cstutorcs

$$P(X = x) = POISSON(\mu, x, false)$$

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

- The cumulative probability is given by

QQ: 749389476

$$P(X \leq x) = POISSON(\mu, x, true)$$

<https://tutorcs.com>

- ... or learn to use your calculator for these.




# Poisson Approximation to the Binomial

程序代写代做 CS编程辅导



- When  $n$  is large and  $p$  is small let  $\mu = np$ .
- The graph below shows this for  $n = 20, p = 0.1$ .



# Poisson Approximation to the Binomial



程序代写代做CS编程辅导

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutormcs@163.com

QQ: 749389476

https://tutorcs.com

- A farmer produces 12 eggs which are collected, graded and packaged into 12. The probability that a defective egg gets through the packaging process is 0.05. What is the probability that a carton will contain more than one defective egg.
- Now,  $n = 12$ ,  $p = 0.05$
- When  $n$  is large and  $p$  is small, we let  $\mu = np$
- $P(X = x) = Poi(0.6)$   $\mu = 12 \times 0.05 = 0.6$
- $P(x > 1) = 1 - P(x = 0) - P(x = 1)$  \*  
 $= 1 - 0.5488 - 0.3293 = 0.1219$
- Using a Poisson approximation to the Binomial Distribution

程序代写代做 CS编程辅导

# Sample Calculations



- For the previous question,  $\mu = 0.05 \times 12 = 0.6$

$$P(x) = \frac{e^{-\mu} \mu^x}{x!}$$

WeChat: cstutorcs

$$\begin{aligned} P(0) &= \frac{e^{-0.6} \times 0.6^0}{0!} \\ &= \frac{0.5488 \times 1}{1} \\ &= 0.5488 \end{aligned}$$

Assignment Project Exam Help

$$\begin{aligned} P(1) &= \frac{e^{-0.6} \times 0.6^1}{1!} \\ &= \frac{0.5488 \times 0.6}{1} \\ &= 0.3293 \end{aligned}$$

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

$$\begin{aligned} P(2) &= \frac{e^{-0.6} \times 0.6^2}{2!} \\ &= \frac{0.5488 \times 0.36}{2} \\ &= 0.0988 \end{aligned}$$

程序代写代做 CS编程辅导

# Mean and Variance



- The table below is a summary of the formula for each distribution as well as the mean and variance of each.

WeChat: cstutorcs

Distribution	Formula	Mean	Variance
Binomial	$P(X = x) = {}^nC_x p^x (1-p)^{(n-x)}$	$np$	$np(1-p)$
Poisson	$P(X = x) = \frac{e^{-\mu} \mu^x}{x!}$	$\mu$	$\mu$

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

Know how to calculate Binomial and Poisson probabilities with your calculator.

# Binomial vs Poisson

程序代写代做 CS编程辅导

- Binomial:

2 para



&  $p$

Number of trials

Probability of success

Series

– Continuous

Finite – Maximum known

Random variable – count number of successes in fixed number of trials

WeChat: cstutorcs

Assignment Project Exam Help

- Poisson:

1 parameter,  $\mu$  (= mean)

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

Discrete (0, 1, 2, ...)

QQ: 749389476

Maximum theoretically infinite

<https://tutorcs.com>

Random variable – count number of independent arrivals or independent events

程序代写代做 CS编程辅导

# Motivating problem



- A cosmetics retailer has 60 of a particular lipstick during the year. The shop is open 300 days per year. The shop keeps one lipstick on the counter and no others in stock. If the lipstick sells it can be replaced prior to opening the next trading day. However if a second or third... customer tries to buy the lipstick on any given day they lose the sale.
- What is the probability that the retailer loses a sale because they are out of stock?

WeChat: estutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

<https://flux.qa> <sup>程序代写 代做CS编程辅导</sup> (Feed code: SJ6KGV)

## Question 5 – ating Problem

The probability that a retailer loses a sale because they are out of stock:

<sup>WeChat: cstutorcs</sup>

- 
- <sup>Assignment Project Exam Help</sup>
- A. has a Binomial distribution.
  - ✓ B. has a Poisson distribution.

<sup>Email: tutorcs@163.com</sup>

<sup>QQ: 749389476</sup>

<sup><https://tutorcs.com></sup>

程序代写代做 CS编程辅导

# Solution to the motivating problem:

$$\mu = 60/300 = 0.2$$



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutors@163.com

QQ: 749389476

<https://tutorcs.com>

$P(x) = \frac{e^{-\mu} \mu^x}{x!}$	mand =	60
$P(0) = \frac{e^{-0.2} \times 0.5^0}{0!}$	year =	300
$= 0.8187$	aily demand =	0.2
	Distribution =	Poisson
Daily Demand	Probability	Lost Sale
0	0.8187	-
1	0.1637	-
2	0.0164	0.0164
3	0.0011	0.0011
4	0.0001	0.0001
5	0.0000	0.0000
Total		0.0175



程序代写代做 CS编程辅导

# Reading/Questions



- Reading:

- 7<sup>th</sup> Ed. Section 7.2, 7.6, 7.7.

- Questions:

WeChat: cstutorcs

- 7<sup>th</sup> Ed. Questions 7.50, 7.53, 7.54, 7.56, 7.58, 7.60, 7.62, 7.64, 7.63, 7.68, 7.69.

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

- Also file: Probability Distributions.xls

QQ: 749389476

<https://tutorcs.com>