



MONASH University

Information Technology

程序代写代做 CS编程辅导

FIT1006



Business Information Analysis

WeChat: cstutorcs

Assignment Project Exam Help

Lecture 17

Email: tutorcs@163.com

Hypothesis Testing

QQ: 749389476

<https://tutorcs.com>

Topics covered:

程序代写代做 CS编程辅导

- The Null and Alternative Hypothesis
- The test procedure



- The hypothesis test for a population mean and proportion.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Student Evaluation of Teaching

程序代写代做 CS编程辅导



- SETU, Please take 5 minutes to evaluate this unit.
 - Your feedback is confidential and used to improve teaching and learning across all units.
 - There is a \$5,000 prize draw on offer to students who complete their evaluations this semester (100 * \$50 gift vouchers)
 - Two options to access SETU:
 1. Use the SETU link on the right hand panel in the Moodle homepage.
- Or:
2. At the bottom of 'Week 10' content page


WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



程序代写代做 CS编程辅导

Mid-semester Test Results

- Well done! Test marks were released this week. (Sorry for the delay as we need to cater for the consideration cases who sat for the test on Monday)



Test Scores (100%)

Mean	50.8
Standard Error	1.2
Median	50.0
Mode	50.2
Standard Deviation	15.8
Sample Variance	250.2
Kurtosis	-0.4
Skewness	0.1
Range	73.3
Minimum	17.5
Maximum	90.8
Sum	8816.5
Count	176

FIT1006 2021 S1 Mid Sem Test


WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



程序代写代做 CS编程辅导

Motivating Problem

- Would Labor have an election today?
- The Australian poll had the two-party preferred vote at: ALP 51% vs Coalition (Liberal) 49% from a sample of 1,160 people chosen at random (taken on 25 April 2021).

WeChat: estutorcs

Assignment Project Exam Help

- Hint: Find a 95% CI for the expected Liberal-NP vote.

QQ: 749389476

- Ref: <http://www.theaustralian.com.au/national-affairs/newspoll>
<https://tutorcs.com>

程序代写代做 CS编程辅导

Statistical Inference



- Statistical inference is concerned with the way we draw conclusions about the population using a sample.

WeChat: cstutorcs

- There are two approaches to statistical inference:

Assignment Project Exam Help

- confidence intervals for population parameters,

Email: tutorcs@163.com

- hypothesis testing, in which we test an assumption, or point of view about a population parameter.

QQ: 749389476

<https://tutorcs.com>

- The two approaches are closely related.

程序代写代做 CS编程辅导

Chance Difference?



- When we are hypothesis testing, we are attempting to determine whether a sample statistic could plausibly have come from a population having a certain parameter value, or whether it just occurred by chance.
- For example, if we take a sample of 100 student test results and find the sample mean is 37. We would be more accepting of the hypothesis that the population mean is 40 rather than 60, as it is unlikely that a population with mean 60 would yield a sample of 100 having a mean of 37.
- But at what value would your acceptance change?

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

What is an Hypothesis?



- An hypothesis is an assumption, or statement about a population parameter. Some possible hypotheses are:

- The population mean = 50.
- The proportion of left handed people in the class is 0.1.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

- We then test the hypothesis in a systematic way and state a conclusion based on the test.

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

The Null and Alternate Hypothesis



- When we test the null hypothesis, we test against an alternative. For example, the hypothesis:
 - H_0 : The population mean = 50
- Could be tested against the alternative hypothesis:
 - H_1 : The population mean > 50.
- We use H_0 to denote the null hypothesis – this is the hypothesis that our population parameter has no difference from a particular value.
- The alternative hypothesis is denoted H_1 .

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

One and Two Sided Tests



- H_0 : The population mean = 50
- Could be tested against **one sided** hypothesis:
 - H_1 : The population mean > 50 .
 - *Reject the hypothesis for high values of the sample statistic*
- Alternatively H_0 could be tested against the **two sided** hypothesis:
 - H_1 : The population mean $\neq 50$.
 - *Reject the hypothesis for high **or** low values of the sample statistic.*

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

One Sided Test: rejection region



- For a one sided test there is an upper or lower rejection region. Reject H_0 if the statistic lies in the rejection region as defined by our test.

Reject if values less than a certain value


WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



Two Sided Test: rejection region

程序代写代做 CS编程辅导



- For a two sided test there is an upper and lower rejection region. Reject H_0 if the statistic lies in either region.

Reject if smaller

Area = $\alpha/2$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Rejection Regions

Reject if larger

Acceptance Region

Critical Values

Estimation

程序代写代做 CS编程辅导

Population



WeChat: cstutorcs

Assignment Project Exam Help

Sample

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>
We are $(1-\alpha)\%$ sure the
population parameter
falls within this range

Hypothesis testing

程序代写代做 CS编程辅导



Population

Sample

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Set critical value(s) to be $(1-\alpha)\%$ certain about μ .

\bar{X}

μ

Reject

Reject

Assume population

Accept
assumption

H_0

程序代写代做 CS编程辅导

The Steps of Hypothesis Testing

1. Decide on a null hypothesis H_0 .
2. Decide on an alternative hypothesis H_1 .
3. Decide on a significance level.
4. Calculate the appropriate test statistic.
5. Find from tables the corresponding tabulated test statistic.
6. Compare calculated and tabulated test statistics and decide whether to accept or reject the null hypothesis.
7. State the conclusion and assumptions of the test.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

(Source: Rees, D.G. Essential Statistics, Chapman and Hall 1995.)

Example 1

程序代写代做 CS编程辅导



- A hypothesis test for population mean when the population variance is known.

(Note: unrealistic situation that lets us use the Normal distribution)

- The Axle manufacturing company has been making axles for a long time and kept records for every axle produced.
 - Population parameters are $\mu = 90\text{mm}$ and $\sigma = 2.5\text{mm}$.
 - A sample of 100 axles from new machine has mean = 92.7.
 - Is the new machine making parts with same average length required (90mm)?
 - Assume a 1% significance.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

<https://flux.qa> (Feed code: SJ6KGV)

Question 1



From previous samples, the company is using a new machine... a sample of 100 yielded a mean length of 92.7mm. Which hypothesis test do we perform to test whether the new machine is producing parts with the same average length (of 90mm) as required?

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

- A. $H_0: \mu = 90, H_1: \mu > 90.$
- ✓ B. $H_0: \mu = 90, H_1: \mu \neq 90.$
- C. $H_0: \mu = 92.7, H_1: \mu > 92.7.$
- D. $H_0: \mu = 92.7, H_1: \mu \neq 92.7.$

<https://flux.qa> 程序代写代做CS编程辅导 (Feed code: SJ6KGV)

Question 2

For a 1% significance, two-sided test, the z statistic is:

A. 1.645

B. 1.96 $\alpha = \text{significance} = 0.01$

C. 2.33

D. 2.58

WeChat: cstutorcs




Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



Cumulative Probabilities for the Standard Normal Distribution

Table gives $P(Z < z)$ for $Z = N(0,1)$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952

Given in Slide 16:
 $\mu = 90\text{mm}$ and $\sigma = 2.5\text{mm}$

Solution

程序代写代做 CS编程辅导

$$\frac{x - \mu}{\sigma_x} = \frac{92.7 - 90}{2.5/\sqrt{100}} = 10.8$$

1. $H_0, \mu = 90\text{mm}$

2. $H_1, \mu \neq 90\text{mm}$

3. Significance = 0.01.

4. The test statistic, $x = 92.7$. We calculate Z_x .
(standardising) $Z = (92.7 - 90)/(2.5/\sqrt{100}) = 10.8$

5. From tables the calculated critical values are ± 2.58

6. We see that $10.8 > 2.58$ and thus we reject H_0 .

7. Thus we conclude that the axles produced by the new machine have a mean significantly different from 90mm (1% level), assuming that axle length is normally distributed.



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

From the previous slide: Two tail critical value is 2.58

程序代写代做 CS编程辅导
...visually, this is what it means...



and thus we reject H_0

WeChat: cstutorcs


Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Standardised test
statistic = 10.8



Example 2

程序代写代做 CS编程辅导



- A hypothesis test of the population mean when the population variance is not known.
- It is claimed that Melbourne families are spending more than \$150 per week on food and grocery items on average. A sample of 15 families was surveyed and the amount spent each week was recorded. Do these results support this thesis?
- Weekly food and grocery expenditure (\$):
- 156, 234, 199, 78, 256, 189, 221, 49, 220, 178, 120, 290, 97, 177, 231.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com


QQ: 749389476

<https://tutorcs.com>

Summary Statistics from Excel

程序代写代做 CS编程辅导

$$\sigma_x = \frac{s}{\sqrt{n}} = \frac{68.5}{\sqrt{15}} = 17.7$$



✓ Mean	179.7
✓ Standard Deviation	17.7
Median	189.0
Mode	#N/A
Standard Deviation	68.5
Sample Variance	4697.0
Kurtosis	0.5
Skewness	-0.5
Range	241.0
Minimum	49.0
Maximum	290.0
Sum	2695.0
Count	15.0

Website: tutorcs.com

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

<https://flux.qa> 程序代写代做CS编程辅导 (Feed code: SJ6KGV)

Question 3



For a one sided 15 observations the t statistic for a 1% significance is:

A. 2.6025

WeChat: cstutorcs B. 2.9467 $\alpha = \text{significance} = 0.01$

✓ C. 2.6245

Assignment Project Exam Help D. 2.9768 (since one tail)
 $V = 15 - 1 = 14$

Email: tutors@163.com

Critical Values of the t Distribution								
Table gives upper critical values only								
	α							
n	0.300	0.200	0.150	0.100	0.050	0.025	0.010	0.005
12	0.5386	0.8726	1.0832	1.3562	1.7823	2.1788	2.6810	3.0545
13	0.5375	0.8702	1.0795	1.3502	1.7709	2.1604	2.6503	3.0123
14	0.5366	0.8681	1.0763	1.3450	1.7613	2.1448	2.6245	2.9768
15	0.5357	0.8662	1.0735	1.3406	1.7531	2.1314	2.6025	2.9467
16	0.5350	0.8647	1.0711	1.3368	1.7459	2.1199	2.5835	2.9208
17	0.5344	0.8633	1.0690	1.3334	1.7396	2.1098	2.5669	2.8982
18	0.5338	0.8620	1.0672	1.3304	1.7341	2.1009	2.5524	2.8784

Given in Slide 22:
 $\mu = 179.7$ and $\sigma = 68.5$

Solution

程序代写代做 CS编程辅导

$$\frac{\bar{x} - \mu}{\sigma_x} = \frac{179.7 - 150}{17.7} = 1.67$$

1. $H_0, \mu = \$150$
2. $H_1, \mu > \$150$ (one-tailed experiment)
3. Significance = 0.01.



From the previous slide: One tail at $\alpha = 0.01$ and $v = 14$, critical value = 2.625

4. The test statistic, $\bar{x} = 179.7$. We calculate T_x .
(standardising) $T = (179.7 - 150) / (68.5 / \sqrt{15}) = 1.67$
5. From tables our calculated critical value $T_{(0.01)(v=14)}$ is 2.625
6. We see that $1.67 < 2.625$ and thus we do not reject H_0 .
7. Thus we conclude that the mean expenditure is not significantly greater than \$150, assuming that expenditure is normally distributed.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

SYSTAT Solution

程序代写代做 CS编程辅导



- One-sample t-test: GROCERIES with 15 cases
- Ho: Mean = 150.0; Test Alternative = 'greater than'

- Mean = 179.667
- SD = 68.534
- t = 1.677
- df = 14
- p-value = 0.058


WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



程序代写代做 CS编程辅导
...visually, this is what it means...



1. 5 and thus we do not reject H_0

Standardised test

statistic = 1.67

WeChat: cstutorcs


Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Critical Value = 2.625



Example 3

程序代写代做 CS编程辅导

- A hypothesis test for a population proportion.



- In a contest of two political parties, a party will win if it gets more than 50% of the vote. In a contest between the Liberal Party and the Australian Labor Party for a particular electorate as survey of 237 voters, 180 indicated that they would vote for the ALP in the next election. Test the hypothesis that the ALP will win the next election.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com


QQ: 749389476

<https://tutorcs.com>

$$p = \frac{180}{237}$$

程序代写代做 CS编程辅导

Example 3 continued

- We assume that s of binomial probabilities hold (that is, n independent trials with probability π of success, $np > 5$ and $n(1-p) > 5$) and test the hypothesis that the population parameter $\pi > 0.5$.

WeChat: cstutorcs

- The standard error of the sample is determined using the assumed proportion in the hypothetical distribution, $H_0 (\pi)$.

Email: tutorcs@163.com

$$p = \frac{180}{237} = 0.7594$$

QQ: 749389476

$$\sigma_p = \sqrt{\frac{\pi(1-\pi)}{n}} = \sqrt{\frac{0.5(1-0.5)}{237}} = 0.0325$$

<https://tutorcs.com>

<https://flux.qa>

程序代写代做CS编程辅导
(Feed code: SJ6r)

Remember,
99% confident
for one-sided
test, $Z = 2.33$

Question 4

For a one sided and 1% significance, $z =$



A. 1.645

B. 1.96

C. 2.33

D. 2.58

$\alpha = \text{significance} = 0.01$
(since one tail)

WeChat: cstutorcs

Cumulative Probabilities for the Standard Normal Distribution

Table gives $P(Z < z)$ for $Z = N(0,1)$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Given in Slide 28:

$$p = 0.7594 \text{ and } \sigma_p = 0.0325$$

Solution

$$\frac{0.7594 - 0.5}{0.0325} = 7.98$$

1. $H_0, \pi = 0.5$
2. $H_1, \pi > 0.5$ (a one-tailed experiment)
3. Significance = 0.01.



From the previous slide: One tail at $\alpha = 0.01$ critical value

4. The test statistic, $p = 0.7594$. We calculate the Z_x .
(standardising) $Z = \frac{(0.7594 - 0.5)}{0.0325} = 7.98$

5. From tables our calculated critical value $Z_{(0.01)}$ is 2.33

6. We see that $7.98 > 2.33$ and thus we reject H_0 .

7. Thus we conclude that proportion of voters intending to vote ALP is greater than 50% at the 1% level.

Assuming the rules for a binomial probability hold.

程序代写代做 CS编程辅导

Motivating Problem

- Would Labor have an election today?
- The Australian poll had the two-party preferred vote at: ALP 51% vs Coalition (Liberal) 49% from a sample of 1,160 people chosen at random (taken on 25 April 2021).

WeChat: estutorcs

Assignment Project Exam Help

- Ref: <http://www.theaustralian.com.au/national-affairs/newspoll>
Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



<https://flux.qa> (Feed code: 8J0KGV)

Question 5



Would Labor have won a Federal Election if one is held today? Newspoll had the two-party preferred vote at: Labor 51% Liberal-NP 49% (taken on 25th April 2021). For this question, the null and alternate hypothesis is:

WeChat: estutorcs

Assignment Project Exam Help

Email: tutores@163.com

✓ A. $H_0: \pi = 0.50, H_1: \pi > 0.50$.

QQ: 749389476

B. $H_0: \pi = 0.50, H_1: \pi \neq 0.50$.

<https://tutorcs.com>

C. $H_0: \pi = 0.51, H_1: \pi > 0.51$.

D. $H_0: \pi = 0.51, H_1: \pi \neq 0.51$.

程序代写代做 CS编程辅导

Motivating Problem in Groups



(a) Is Labor going to win the next election based on these data? (Use 1% Significance)

$$p = 0.51$$

WeChat: cstutorcs

$$\sigma_p = \sqrt{\frac{\pi(1-\pi)}{n}} = \sqrt{\frac{0.5(1-0.5)}{1160}} = 0.0147$$

Email: tutorcs@163.com

$$Z_{Calc} = \frac{0.51 - 0.50}{0.0147} = 0.68$$

0.0147
<https://tutorcs.com>

$$Z_{Tables} = 2.33$$

Since $0.68 < 2.33$ we do not reject H_0

<https://flux.qa> (Feed code: 8J0KGV)

Question 6



The Newspoll (2021) gave the two-party preferences at 51% Labor and 49% Coalition, for a sample size of 1160 ($Calc\ z = 0.68$). Based on this information, if an election had been held at the time:

Assignment Project Exam Help

- A. Labor would have won (99% confidence)
- B. Coalition would have won (99% confidence)
- ✓ C. Too close to call (1% sig)
- D. Just don't know...

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

Reading/Questions (Selvanathan)



- Reading: Hypothesis Testing
 - 7th Ed. Sections 12.3, 12.5, 12.6.

WeChat: cstutorcs

- Questions: Hypothesis Testing

Assignment Project Exam Help

- 7th Ed. Questions 12.1, 12.19, 12.25, 12.26, 12.56, 12.59, 12.65, 12.66, 12.67, 12.70, 12.72, 12.74.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>