

程序代写代做 CS编程辅导

FIT2014 Theory of Computation



ecture 11

(A) Closure properties;

WeChat: cstutorcs

(B) Pumping Lemma for Regular Languages

Assignment Project Exam Help

Email: tutorcs@163.com

slides by Graham Farr

based in part on previous slides by David Albrecht

QQ: 749389476

<https://tutorcs.com>

COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

Warning

This material has been reproduced and communicated to you by or on behalf of Monash University in accordance with s113P of the Copyright Act 1968 (the Act).

The material in this communication may be subject to copyright under the Act.

Any further reproduction or communication of this material by you may be the subject of copyright protection under the Act.

Do not remove this notice.

Overview

程序代写代做 CS编程辅导



- ▶ Closure properties of regular languages
- ▶ Circuits in FAs
- ▶ Pumping Lemma
- ▶ Non-regular Languages

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Closure properties of regular languages

Definition

If doing some operation on regular languages always produces another regular language, then we say that the class of regular languages is **closed** under that operation.

程序代写代做 CS编程辅导



We will see that regular languages are closed under:

- ▶ complement
- ▶ union
- ▶ intersection
- ▶ concatenation

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

Theorem.

The complement of a regular language is regular.

<https://tutorcs.com>

We prove this using Kleene's Theorem.

Closure properties of regular languages

程序代写代做 CS编程辅导

Theorem.

The complement of a regular language is regular.



Proof. (outline)

Suppose we have a Regular Language.

There must be a regular expression that defines it.

So, by Kleene's Theorem, there is a Finite Automaton (FA) that defines this language.

We can convert this FA into one that defines the complement of the language. (See Lecture 7.)

So, by Kleene's Theorem, there is a regular expression that defines the complement. \square

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutors@163.com

QQ: 749389476

<https://tutorcs.com>

Closure properties of regular languages

Theorem.

程序代写代做 CS编程辅导

The union of two regular languages is regular.



Proof.

Suppose L_1 and L_2 are regular.

WeChat: cstutorcs

By definition of “regular language”,

there exist regular expressions R_1 and R_2

that describe L_1 and L_2 , respectively.

Email: tutorcs@163.com

Then $R_1 \cup R_2$ is a regular expression that describes $L_1 \cup L_2$.

QQ: 749889476

- This uses part 3(iii) of the inductive definition of regular expressions in Lecture 6.

https://tutorcs.com

So $L_1 \cup L_2$ is regular. \square

Closure properties of regular languages

程序代写代做 CS编程辅导



Theorem.

The intersection of two regular languages is regular.

WeChat: cstutorcs

Assignment Project Exam Help

We can't just mimic the proof that regular languages are closed under union, since there is no \cap operation on regular expressions.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Closure properties of regular languages

Theorem.

程序代写代做 CS编程辅导

The intersection of two regular languages is regular.



Proof.

Suppose L_1 and L_2 are regular.

WeChat: cstutorcs

We know that their complements $\overline{L_1}$ and $\overline{L_2}$ are regular.

Assignment Project Exam Help

So the union of these, $\overline{L_1} \cup \overline{L_2}$, is therefore regular, by the previous Theorem.

Email: tutorcs@163.com

Its complement, $\overline{\overline{L_1} \cup \overline{L_2}}$, must also be regular.

QQ: 749389476

But $\overline{\overline{L_1} \cup \overline{L_2}} = \overline{\overline{L_1}} \cap \overline{\overline{L_2}} = L_1 \cap L_2$.

https://tutorcs.com

So $L_1 \cap L_2$, must also be regular. \square

Closure properties of regular languages

Exercises

程序代写代做 CS编程辅导

- Prove that the class of regular languages is closed under **concatenation**.



$L_1 L_2 = \{xy : x \in L_1, y \in L_2\}$

- Prove that the class of regular languages is closed under **symmetric difference**.
(You can use the closure results we've already proved.)

WeChat: cstutorcs

Assignment Project Exam Help

$L_1 \triangle L_2 := \{\text{strings in } L_1 \text{ but not in } L_2, \text{ or in } L_2 \text{ but not in } L_1\}$

Email: tutorcs@163.com

QQ: 749389476

- Is the class of regular languages closed under taking **subsets**?
i.e., is a subset of a regular language necessarily regular?
- Is the class of regular languages closed under taking **supersets**?
i.e., is a superset of a regular language necessarily regular?

<https://tutorcs.com>

Circuits in Finite Automata

Definition

A **circuit** is a directed path which starts and ends at the same state.
The **length** of a circuit is the number of edges in the path.

程序代写代做 CS编程辅导



Observation

Take any Finite Automaton.

Take any string w with as many letters as there are states in that Finite Automaton.

WeChat: csnytuts

Then the path taken for input w must contain a circuit.

Assignment Project Exam Help

We can divide w up naturally into three parts $w = xyz$, where:

Email: tutores@163.com

x := the part before the circuit;

QQ: 749389476

y := the part that goes around the circuit;

https://tutores.com

z := the part after the circuit;

$w = \underline{a} \underline{baaa} \underline{abb}$
 $\quad \quad \quad x \quad y \quad z$


If w is accepted,
 then so are ...

$\underline{a} \underline{abb}$
 $\quad \quad \quad x \quad z$

$\underline{a} \underline{baaa} \underline{abb}$
 $\quad \quad \quad x \quad y \quad z$

$\underline{a} \underline{baaa} \underline{baaa} \underline{abb}$
 $\quad \quad \quad x \quad y \quad y \quad z$

...
 ...



$w = \underline{b}a\underline{b}b$
 $\quad \quad \quad y \quad z$
 $x = \varepsilon$

If w is accepted,
then so are ...


$\underline{b}b$
 $\quad \quad \quad z$

$\underline{b}a\underline{b}b$
 $\quad \quad \quad y \quad z$

$\underline{b}a\underline{b}a\underline{b}b$
 $\quad \quad \quad y \quad y \quad z$

...

...



Pumping Lemma

Theorem. (Pumping Lemma) 程序代写代做 CS编程辅导

Let L be an infinite regular language accepted by a FA with N states.

Then for all words $w \in L$ with $|w| \geq N$ letters,

there exist strings x, y, z , with $|xy| \leq N$ such that



► $w = xyz$

► $\text{length}(x) + \text{length}(y) \leq N$ WeChat: cstutorcs

► for all $i \geq 0$, $xy^iz \in L$, Assignment Project Exam Help
i.e.,

$xy^iz \in L$ Email: tutorcs@163.com

QQ: 749389476

Symbolically:

<https://tutorcs.com>

$$\forall w \in L : |w| \geq N \Rightarrow (\exists x, y, z : (w = xyz) \wedge (y \neq \varepsilon) \wedge (|x| + |y| \leq N) \wedge (\forall i \geq 0 : xy^iz \in L))$$

Pumping Lemma

Proof.

Take any word $w \in L$ with $\geq N$ letters.

By our earlier Observation on c FAs, the path taken by w must include a circuit.

Let

- x be the letters of w up to the first circuit.
- y be the letters corresponding to the circuit.
- z be the remaining letters of w .

We have:

- ▶ $w = xyz$ by construction.
- ▶ Since the circuit exists, $y \neq \epsilon$.
- ▶ $\text{length}(x) + \text{length}(y) \leq N$, since the FA reads xy without repeating any state.
- ▶ Since $w = xyz \in L$, and y starts and finishes at $\text{endState}(x)$, and z goes from $\text{endState}(x)$ to a Final State, we can repeat y any number of times (or none) and still we end up at the same Final State.



程序代写代做 CS编程辅导



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

https://tutorcs.com

Pumping Lemma: application

Consequence

Using the Pumping Lemma we

程序代写代做 CS编程辅导



there are non-regular languages.

Method

Assume L is regular.

Then, by Kleene's Theorem, it is recognised by some FA.

Let N be the number of states in this FA.

Choose a suitable word $w \in L$, of length $\geq N$.

Show that, for any $x, y \neq \varepsilon$, and z such that $w = xyz$ and $|xy| \leq N \dots$

\dots there exists $i \geq 0$ s.t. $xy^iz \notin L$.

Contradiction.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Compare quantifiers above with those in Pumping Lemma

Non-regular languages

HALF-AND-HALF:

$$L := \{a^n b^n : n \geq 0\} = \{\varepsilon, ab, aabb, aaabbb, \dots\}.$$

程序代写代做 CS编程辅导



Theorem.

L is not regular.

Proof. (by contradiction)

Assume that L is regular.

Let $N = \#$ states in an FA for it.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

Choose $w := a^{\lceil N/2 \rceil} b^{\lceil N/2 \rceil}$.

QQ: 749389476

$\overbrace{aaa \dots \dots \dots aa}^{\lceil N/2 \rceil \text{ letters}} \overbrace{bbb \dots \dots \dots bb}^{\lceil N/2 \rceil \text{ letters}}$

Observe that $|w| \geq N$.

<https://tutorcs.com>

Consider any $x, y \neq \varepsilon$, and z such that $w = xyz$ and $|xy| \leq N$.

Think: are $xz, xyz, xyyz, \dots, xy^N z, \dots$ all in L ?

Non-regular languages

Case 1: y is all a's.

Then $xyyz$ has more a's than xz , since $y \neq \varepsilon$.

So $xy^2z \notin L$.

程序代写代做CS编程辅导



$\overbrace{aaa \dots aa}^{\lceil N/2 \rceil \text{ letters}} \underbrace{bbb \dots bb}_{y}$

Case 2: y is all b's.

Then $xyyz$ has more b's than a's, since $y \neq \varepsilon$.

So $xy^2z \notin L$.

WeChat: cstutorcs

Assignment Project Exam Help

$aaa \dots aa \underbrace{bbb \dots bb}_y$

Case 3: y contains an ab.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

$aaa \dots aa \underbrace{bbb \dots bb}_y$

Then $xyyz$ has two occurrences of ab. This cannot happen for strings in L . So $xy^2z \notin L$.

In every possible case, we have found an i such that $xy^iz \notin L$.

This violates the conclusion of the Pumping Lemma.

Contradiction. \square

Non-regular languages

HALF-AND-HALF:

$L := \{a^n b^n : n \geq 0\} = \{\epsilon, ab, aabb, aaabbb, \dots\}.$

程序代写代做 CS编程辅导



Theorem.

L is not regular.

Proof. (by contradiction)

Assume that L is regular. Let $N = \#$ states in an FA for it.

Choose $w = a^N b^N$.

[No need for w to be of minimum length.]

Consider any $x, y \neq \epsilon$, and z such that $w = xyz \dots$

\dots and $|xy| \leq N$.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476 [Previous proof didn't use $|xy| \leq N$. Can it help?]

Think: are $xz, xyz, xyyz, \dots, xyy^N z, \dots$ all in L ?

<https://tutorcs.com/>

How many cases now?

Non-regular languages

Just one case: y is all a's.

Consider $xyyz$.

It has more a's than b's, since $y \neq \epsilon$.

So $xy^2z \notin L$.

This violates the conclusion of the Pumping Lemma.

Contradiction. \square

程序代写代做 C/C++ 编程辅导




WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>



Non-regular languages

程序代写代做 CS编程辅导

$\text{EQUAL} := \{ \text{all words which have an equal number of a's and b's} \}$
 $= \{ \epsilon, ab, bab, abba, baba, \dots \}$



Theorem.

EQUAL is not regular.

WeChat: cstutorcs

Proof. Assume EQUAL is regular.

Observe:

Assignment Project Exam Help

$\text{HALF-AND-HALF} := \{ a^n b^n : n \geq 0 \} = \text{EQUAL} \cap a^* b^*.$

Email: tutorcs@163.com

This implies that HALF-AND-HALF is also regular, since the language defined by $a^* b^*$ is regular, and regular languages are closed under intersection.

QQ: 749389476

<https://tutorcs.com>

But we have just seen that HALF-AND-HALF is non-regular.

This is a contradiction.

So our initial assumption, that EQUAL is regular, is wrong.

Therefore EQUAL is non-regular. \square

Non-regular languages

程序代写代做 CS编程辅导

PALINDROME $:=$ { all the strings which are the same if they are spelt backwards }
 $=$ $\{\epsilon, a, b, a, b, a, a, aba, bab, bbb, \dots\}$



Theorem.

PALINDROME is non-regular. WeChat: cstutorcs

Proof. (by contradiction)

Assume PALINDROME is regular. Email: tutorcs@163.com

Then there exists a FA with N states which accepts PALINDROME.

QQ: 749389476

Choose $w = a^N b a^N$.

<https://tutorcs.com>


$\overbrace{aaa \dots \dots \dots aa}^{N \text{ letters}} b \overbrace{aaa \dots \dots \dots aa}^{N \text{ letters}}$

Non-regular languages

Consider all strings $x, y \neq \varepsilon$, and z such that

► $w = xyz$,

► $\text{length}(x) + \text{length}(y) \leq N$




Consider xyy^2z .

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476



Since $y \neq \varepsilon$, the solitary b in w is *more than* half-way along xy^2z .

So xy^2z is not a palindrome.

<https://tutorcs.com>

This contradicts the conclusion of the Pumping Lemma applied to PALINDROME.

So our initial assumption, that PALINDROME is regular, is wrong.

Therefore PALINDROME is not regular.



Revision

{ all languages }

程序代写代做 CS编程辅导



{ regular languages }

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

Reading: Sipser, Ch. 1. <https://tutorcs.com>

- ▶ closure properties: pp. 58–63.
- ▶ Pumping Lemma, non-regular languages: pp. 77–82.