

程序代写代做 CS编程辅导

FIT2014 Theory of Computation



Lecture 4

Proofs

WeChat: cstutorcs

Assignment Project Exam Help

slides by Graham Farr

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

<https://tutorcs.com>

This material has been reproduced and communicated to you by or on behalf of Monash University  
in accordance with s113P of the Copyright Act 1968 (the Act).

The material in this communication may be subject to copyright under the Act.

Any further reproduction or communication of this material by you may be the subject of copyright protection under the Act.

Do not remove this notice.

# Overview

- ▶ Finding proofs
- ▶ Proof by construction
- ▶ Proof by cases
- ▶ Proof by contradiction
- ▶ Proof by induction
  - ▶ inductive basis
  - ▶ inductive step
  - ▶ conclusion

程序代写代做 CS编程辅导



WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

# Proof (recap)

程序代写代做 CS编程辅导

- ▶ A step-by-step argument establishes, logically and with certainty, that something is true.
  - ▶ Should be verifiable.
  - ▶ Must be finite.
  - ▶ Every statement must be:
    - ▶ something you already know at the start
      - ▶ a definition
      - ▶ an axiom
      - ▶ a previously-proved theorem
- or
- ▶ a logical consequence of some conjunction of previous statements.



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutores@163.com

QQ: 749389476

<https://tutorcs.com>

## Proof (recap)

程序代写代做 CS编程辅导

If you've previously established  
and also that  $P \Rightarrow Q$   
then you can deduce  $Q$ .  
(*modus ponens*)



WeChat: cstutorcs

Exercise in Boolean algebra: Prove that  $(P \wedge (P \Rightarrow Q)) \Rightarrow Q$  is a tautology.

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

If you've previously established all of  $P_1, P_2, \dots, P_n$   
and also that  $(P_1 \wedge P_2 \wedge \dots \wedge P_n) \Rightarrow Q$   
then you can deduce  $Q$ .

QQ: 749389476

<https://tutorcs.com>

## Finding proofs

程序代写代做 CS编程辅导

There is no systematic method for finding proofs for theorems.

There are deep theoretical results for this.

(Gödel, 1931; Church, 1936; Turing, 1936)



Discovering proofs is an art as well as a science. It requires

- ▶ skill at logical thinking and reasoning
- ▶ understanding the objects you're working with
- ▶ practice, experience
- ▶ play, exploration
- ▶ creativity and imagination
- ▶ perseverance

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

## Finding proofs: general advice

程序代写代做 CS编程辅导



To prove subset relations,  $A \subseteq$  (where  $A$  and  $B$  are sets):

1. Take a general member of  $A$  and give it a name. e.g., “Let  $x \in A$ ”
2. Use the definition of  $A$  to say something about  $x$ .
3. Follow through the logical consequences of that,
4. ... aiming to prove that  $x$  also satisfies the definition of  $B$ .

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

See, e.g., Lecture 1, slide 30, proof that  $\text{DOUBLEWORD} \subseteq \text{EVEN-EVEN}$ .

See also: Tutorial 1, exercise 1. QQ: 749389476

<https://tutorcs.com>

## Finding proofs: general advice

程序代写代做 CS编程辅导

To prove **set** equality,  $A = B$  (where  $A$  and  $B$  are **sets**):

1. Prove  $A \subseteq B$
2. Prove  $A \supseteq B$



WeChat: cstutorcs

To prove **numerical** equality,  $A = B$  (where  $A$  and  $B$  represent **numbers**):

If algebra can transform  $A$  to  $B$ , then that's good;

but if not:

1. Prove  $A \leq B$
2. Prove  $A \geq B$

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>

Assignment Project Exam Help

# Types of proofs

- ▶ Proof by construction
- ▶ Proof by cases
- ▶ Proof by contradiction
- ▶ Proof by induction

This list is not exhaustive.

Proofs can be quite individual in character and hard to classify, although many will follow one of the above patterns.

Many proofs are a mix of these types.

程序代写代做 CS编程辅导



WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>



# Proof by construction

程序代写代做 CS编程辅导

...also known as:



## Proof by example

- ▶ can be used where the theorem asserts the existence of some object with a specific property just give the example, show it has the property.
- ▶ BUT: an *illustration* is NOT a *proof*.
- ▶ So, if your example merely illustrates the idea of a proof, then it is not, itself, a proof (although it might still be useful in illustrating a proof).
- ▶ Recall Lecture 1: English has a palindrome.

<https://tutorcs.com>

# Proof by cases

程序代写代做 CS编程辅导

... also known as:



## Proof by exhaustion

or (if lots of cases) “brute force”

WeChat: cstutorcs

► identify a number of different cases which cover all possibilities

Assignment Project Exam Help

► Prove the theorem for each of these cases.

► Recall Lecture 1:

Email: tutorcs@163.com

Every English word has a vowel or a “y”

QQ: 749389476

<https://tutorcs.com>

# Proof by contradiction

程序代写代做 CS编程辅导



(also known as "reductio ad absurdum")

- ▶ Start by assuming the negation of the statement you want to prove.
- ▶ Deduce a contradiction.
- ▶ Therefore, the statement must be true.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

# Proof by contradiction

程序代写代做 CS编程辅导

## Theorem.

The statement “This statement



is not a proposition.

## Proof.

Assume that it is a proposition. **WeChat: cstutorcs**

Then it must be either true or false.

If it is true, then it is false.

**Assignment Project Exam Help**

If it is false, then it is true.

**Email: tutorcs@163.com**

So, it is false if and only if it is true.

This is a contradiction.

**QQ: 749389476**

So our assumption, that the statement is a proposition, must be false.

**<https://tutorcs.com>**



# Proof by contradiction

“Every positive integer was one of his personal friends.”

— J. E. Littlewood on Srinivasa Ramanujan, quoted by G. H. Hardy, *Srinivasa Ramanujan* (obituary), *Proceedings of the London Mathematical Society* (1921) xl–lviii. See p. lvii.

## Theorem.

Every natural number is interesting.



## Proof.

Assume that not every natural number is interesting.

So, there exists at least one uninteresting number.

Therefore there exists a *smallest* uninteresting number.

But that number must be interesting, by virtue of having this special property of being the smallest of its type.

This is a contradiction, as this number is uninteresting.

Therefore our original assumption was wrong.

Therefore every natural number is interesting.  $\square$

程序代写代做 CS编程辅导

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutormcs@163.com

QQ: 749389476

https://tutorcs.com



Srinivasa Ramanujan  
(1887–1920)

<https://mathshistory.st-andrews.ac.uk/Biographies/Ramanujan/>

See, e.g., Ch. 14 (Fallacies), in: Martin Gardner, *The Scientific American Book of Mathematical Puzzles and Diversions*, Simon & Schuster, New York, 1959.

# Proof by contradiction

## Comments:

That “theorem” and “proof” is just an informal argument, as the meaning of “interesting” is imprecise and so is “proof”. But it illustrates the structure of a proof by contradiction.

程序代写代做 CS编程辅导



It also illustrates the point that if you know a set of objects is nonempty, then you can choose an element of *smallest* size in the set.

Often, the smallest object in a set may have special properties that can help you go further in the proof.

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

Can you always choose an object of *largest* size in a nonempty set?

QQ: 749389476

<https://tutorcs.com>

Is every integer interesting?

Would the above proof still work, if applied to the set of all integers?

## More proofs

程序代写代做 CS编程辅导

Recall De Morgan's Laws:



$$\neg(P \wedge Q) = \neg P \vee \neg Q$$

WeChat: cstutorcs

We proved these using truth tables.

Assignment Project Exam Help

But, how to prove its extended form? ...

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

For all  $n$ :

QQ: 749389476

$$\neg(P_1 \vee \dots \vee P_n) = \neg P_1 \wedge \dots \wedge \neg P_n$$


<https://tutorcs.com>

## More proofs

程序代写代做 CS编程辅导

### Theorem.

For all  $n$ :


$$\neg(P_1 \vee \dots \vee P_n) = \neg P_1 \wedge \dots \wedge \neg P_n$$

### First proof:

Left-Hand Side is True

if and only if  $P_1 \vee \dots \vee P_n$  is False

if and only if  $P_1, \dots, P_n$  are all False

if and only if  $\neg P_1, \dots, \neg P_n$  are all True

if and only if Right-Hand Side is True

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Let's try for a different proof, using De Morgan's Law.





## More proofs

### Theorem.

For all  $n$ :

$$\neg(P_1 \vee \dots \vee P_n) = \neg P_1 \wedge \dots \wedge \neg P_n$$

程序代写代做 CS编程辅导



**Second proof** (*attempt*):

$$\begin{aligned} & \neg(P_1 \vee \dots \vee P_n) \\ &= \neg((P_1 \vee \dots \vee P_{n-1}) \vee P_n) \quad (\text{Justifying } \dots) \\ &= \neg(P_1 \vee \dots \vee P_{n-1}) \wedge \neg P_n \quad (\text{by De Morgan's Law}) \\ &= \dots \text{ and so on and so on } \dots \\ &= \neg P_1 \wedge \dots \wedge \neg P_n \quad \text{Q.E.D.??} \end{aligned}$$

WeChat: cstutorcs

Assignment Project (Justing Help)

Email: tutores@163.com

QQ: 749389476

<https://tutorcs.com>

Good try, but reader has to infer how to fill the gap.

It's shorthand for a “proof” whose length depends on  $n$ .

But we can turn its main idea into a proper proof.

# Proof by mathematical induction

Suppose you want to prove that a statement  $S(n)$  holds for every natural number  $n$ .



## Principle of Mathematical Induction

IF  $S(1)$  is true *(inductive basis)*  
AND  $\forall k: \text{if } S(k) \text{ is true, then } S(k+1) \text{ is true}$  *(inductive step)*

*inductive hypothesis*  
Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

THEN

$\forall n: S(n)$  is true.

<https://tutorcs.com>

$S(1), \dots, S(n), S(n+1), \dots$

# Proof by mathematical induction

## Theorem.

For all  $n$ :

$$\neg(P_1 \vee \dots \vee P_n) = \neg P_1 \wedge \dots \wedge \neg P_n$$

程序代写代做 CS编程辅导



**Second proof:** We prove it by induction on the # of propositions.

*Inductive basis:*

It is trivially true when we have just one proposition.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

*Inductive step:*

Suppose it's true for  $k$  propositions:

QQ: 749389476

<https://tutorcs.com>

$$\neg(P_1 \vee \dots \vee P_k) = \neg P_1 \wedge \dots \wedge \neg P_k$$

(This our Inductive Hypothesis. We will use it later.)

# Proof by mathematical induction

(continued)

程序代写代做 CS编程辅导

We have:



$$\begin{aligned} & \neg(P_1 \vee \dots \vee P_{k+1}) \\ &= \neg((P_1 \vee \dots \vee P_k) \vee P_{k+1}) \quad \text{(just grouping ...)} \\ &= \neg(P_1 \vee \dots \vee P_k) \wedge \neg P_{k+1} \quad \text{(by De Morgan's Law)} \\ &= \neg P_1 \wedge \dots \wedge \neg P_k \wedge \neg P_{k+1} \quad \text{(by Inductive Hypothesis)} \end{aligned}$$

Assignment Project Exam Help  
Email: tutorcs@163.com

*Conclusion:*

So, by the Principle of Mathematical Induction, it's true for any number of propositions.

QQ: 749389476

<https://tutorcs.com>



# Proof by mathematical induction

## Theorem.

For all  $n$ :

程序代写代做 CS编程辅导



$$n = \frac{n(n+1)}{2}.$$

**Proof:** We prove it by induction

*Inductive basis:*

WeChat: cstutorcs

When  $n = 1$ , LHS = 1 and RHS =  $\frac{1(1+1)}{2} = 1$ . ✓

Assignment Project Exam Help

*Inductive step:*

Email: tutorcs@163.com

Suppose it's true for  $n = k$ :

QQ: 749389476

$$1 + \dots + k = k(k+1)/2$$

<https://tutorcs.com>

We will deduce that it's true for  $n = k + 1$ .

$$1 + \dots + (k+1) = (1 + \dots + k) + (k+1) \quad (\text{preparing to use the inductive hypothesis})$$

## Proof by mathematical induction

程序代写代做 CS编程辅导

$$\begin{aligned} 1 + \cdots + (k+1) &= (1 + \cdots + k) + (k+1) && \text{(preparing to use the inductive hypothesis)} \\ &= k(k+1)/2 + (k+1) && \text{(by the Inductive Hypothesis)} \\ &= (k+1)(k/2 + 1) && \text{(algebra ...)} \\ &= (k+1)(k/2 + 1) \\ &= (k+1)(k+2)/2 \\ &= (k+1)((k+1)+1)/2 \end{aligned}$$



WeChat: cstutorcs

Assignment Project Exam Help

This is just the equation in the Theorem, for  $n = k+1$  instead of  $k$ .  
So the inductive step is now complete. ✓

QQ: 749389476

*Conclusion:*

<https://tutorcs.com>

Therefore, by the Principle of Mathematical Induction, the equation holds for all  $n$ . □

# Proof by mathematical induction

Alternatively, we could make the inductive step go from  $n = k - 1$  to  $n = k$ , instead of from  $n = k$  to  $n = k + 1$ .

程序代写代做CS编程辅导



**Slightly different proof:**

We prove it by induction on  $n$ .

*Inductive basis:*

When  $n = 1$ , LHS = 1 and RHS =  $1 + 1/2 = 3/2$ . ✓

WeChat: cs\_tutors

*Inductive step:*

Suppose it's true for  $n = k - 1$  where  $k \geq 2$ .

Assignment Project Exam Help

Email: tutorcs@163.com

$$1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{k-1} = (k-1)k/2$$

QQ: 749389476

We will deduce that it's true for  $n = k$ .

https://tutorcs.com

$$1 + \dots + k = (1 + \dots + (k-1)) + k \quad (\text{preparing to use the inductive hypothesis})$$

# Proof by mathematical induction

程序代写代做 CS编程辅导

$$\begin{aligned} 1 + \cdots + k &= (1 + \cdots + (k-1)) + k && \text{(preparing to use the inductive hypothesis)} \\ &= (k-1)k/2 + k && \text{the Inductive Hypothesis} \\ &= k(k-1)/2 + k && \text{(algebra ...)} \\ &= k((k-1)/2 + 1) \\ &= k(k+1)/2 \end{aligned}$$



WeChat: cstutorcs

This is just the equation in the Theorem, for  $n = k$  instead of  $k-1$ .

So the inductive step is now complete. ✓

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

*Conclusion:*

QQ: 749389476

Therefore, by the Principle of Mathematical Induction, the equation holds for all  $n$ .  $\square$

<https://tutorcs.com>



# Proof by mathematical induction

程序代写代做 CS编程辅导

## Exercise:

Prove by induction that, for a



$$1^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}.$$

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

Something to think about:

the relationship between induction and recursion

<https://tutorcs.com>

# Proof by mathematical induction

程序代写代做 CS编程辅导

Contrast with “induction” in statistics, which is the process of drawing general conclusions from data.



Statistical induction is typically used in situations where there is some randomness in the data.

WeChat: cstutorcs

Assignment Project Exam Help

Statistical induction cannot be used as a step in a mathematical proof.

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

Mathematical induction is a rigorous and very powerful tool for proofs in mathematics and computer science.

QQ: 749989476

<https://tutorcs.com>

### *Practise doing proofs*

- ▶ tutorial sheets, textbooks,



Sipser, pp. 22–25.

WeChat: cstutorcs

For more about Srinivasa Ramanujan, see:

- ▶ <https://mathshistory.st-andrews.ac.uk/Biographies/Ramanujan/>
- ▶ R Kanigel, *The Man Who Knew Infinity: A Life of the Genius Ramanujan*, Washington Square Press, New York, 1991.
- ▶ *The Man Who Knew Infinity*, feature film, 2015.
- ▶ film review: G Farr, *The Man Who Knew Infinity: inspiration, rigour and the art of mathematics*, *The Conversation*, 24 May 2016.

<https://theconversation.com/the-man-who-knew-infinity-inspiration-rigour-and-the-art-of-mathematics-59520>