

# MSML/DATA 603 MIDTERM EXAM

---

程序代写代做 CS编程辅导

Closed book, closed notes

No calculator or electronic devices



Please, turn off cell phones & smart phones

No speaking or whispering to classmates during exam

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

Date: October 26, 2021

QQ: 749389476

HONOR PLEDGE:

<https://tutorcs.com>

*I pledge on my honor that I have not given or received any unauthorized assistance on this examination. I pledge that I have not intentionally used or attempted to use unauthorized materials or information to assist me in this examination, and I pledge on my honor that I have not looked at or read anything from any classmate's exam papers or scrap-material sheets.*

SAMPLE

---

Student's name and UID (required)

(1) [3 points] Consider a two-label Bayesian Classification case where: all features are statistically independent, their likelihoods are Gaussian, and the features have been scaled to have the same variance. Furthermore, the priors are all the same. The mean of the likelihood for  $\omega_1$  is  $\vec{\mu}_1 = [0 \ 3]^T$  and the mean of the likelihood for  $\omega_2$  is  $\vec{\mu}_2 = [6 \ 9]^T$ . A new instance is studied and its feature vector  $\vec{x} = [4 \ 6]^T$ . (a) What classification would our Bayesian classifier apply to this new instance? (b) Please, justify your response in question (a). (c) Please, sketch the decision boundary in the feature space.

(a) Answer:  $\omega_2$

(b) Answer: Because the options underlined above, this is a "minimum distance classifier".

WeChat: cstutorcs

Wanda's features are closer to  $\vec{\mu}_2$ .

## Assignment Project Exam Help


You could just draw the boundary to get full credit, but if you want to solve it

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

(c) Answer:

QQ: 749389476

<https://tutorcs.com>



$$g_1(\vec{x}) = -\| \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} - \begin{bmatrix} 0 \\ 3 \end{bmatrix} \|^2$$

$$g_2(\vec{x}) = -\| \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} - \begin{bmatrix} 6 \\ 9 \end{bmatrix} \|^2$$

$$g_1(\vec{x}) = g_2(\vec{x}) \text{ at all } \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \text{ where}$$

$$\{x_1^2\} + \{x_2^2 - 6x_2 + 9\} = \{x_1^2 - 12x_1 + 36\} + \{x_2^2 - 18x_2 + 81\}$$

$$12x_2 = -12x_1 + 108$$


$$\boxed{x_2 = -x_1 + 9}$$

(equation of line)

(Draw it or derive it, either way you get full credit.)

decision boundary is a line that is equal distance from each  $\mu$

(2) [3 points] Consider the Bayesian Decision problem we saw in class: viewing the likelihoods for both labels,  $\omega_1$  and  $\omega_2$ , below, please, (a) provide the expectation of the likelihood for  $\omega_2$  (reading the graphs is not easy, so you don't need to be exact) (b) what can you say about the covariance matrices for both likelihoods? (c) Estimate which one seems to have larger variance(s),  $\omega_1$  or  $\omega_2$ ?



WeChat: cstutores

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

Answer (a):  $\vec{\mu}_2 \approx \begin{bmatrix} 2 \\ -2 \end{bmatrix}$

<https://tutorcs.com>

Answer (b): Features are independent, so covariance matrices are

diagonal. Features do not have the same variances (not scaled).

For  $\omega_1$ ,  $x_2$ 's variance  $>$   $x_1$ 's variance. For  $\omega_2$ ,  $x_1$ 's variance  $>$   $x_2$ 's variance.

Answer (c):  $\omega_2$  seems to have smaller variances, since boundary wraps closer to  $\vec{\mu}_2$ .

(3) [2 points] Please, recall our discussion in lecture and (a) explain the difference between a Regression Problem and a Classification Problem, and, (b) explain the difference between Instance Based Learning and Model Based Learning.

程序代写代做 CS编程辅导

Answer (a):

Regression: predict a real value from a



uous interval.

classif

assify an item into exactly

e of several (countable)

categories.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

Answer (b):

Instance based: basis for decision is empirical

QQ: 749389476

and not formulas

<https://tutorcs.com>

Model based: create a model formula to

predict label

(4) [3 points] Consider the 2-label classification case we saw in lecture in reference to admitting a student to "X" Medical School. We have training instances that each include two attributes: (i) undergraduate GPA and (ii) grade in organic chemistry. Assume the following:

- all instances (applicant students' grades) are independent from each other,
- all instances within a class ("admitted" or  $\omega_2$  = "not admitted") are identically distributed,
- the likelihood function for each class is Gaussian with pdf  $\mathcal{N}(\vec{\mu}_1, \Sigma_1)$  and  $\mathcal{N}(\vec{\mu}_2, \Sigma_2)$ , respectively.

(a) Given the information above, what method did we use to estimate  $\vec{\mu}_1, \Sigma_1, \vec{\mu}_2$ , and  $\Sigma_2$ ?

Maximum Likelihood estimation

WeChat: cstutorcs

(b) The training instances are provided in the Table 1, here. Using the method you named in (a), please find the estimated for the four parameters.

Instance number	Label	undergraduate GPA	grade in organic chemistry
1	Admitted	4	3
2	Admitted	3	2
3	Not Admitted	2	1
4	Not Admitted	1	3

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

[It's OK to leave answers as fractions, such as for example 7/4]

$$\vec{\mu}_1 = \frac{1}{2} \left\{ \begin{bmatrix} 4 \\ 3 \end{bmatrix} + \begin{bmatrix} 3 \\ 2 \end{bmatrix} \right\} = \begin{bmatrix} 3.5 \\ 2.5 \end{bmatrix}$$

$$\begin{aligned} \Sigma_1 &= \frac{1}{2} \left\{ \left( \begin{bmatrix} 4 \\ 3 \end{bmatrix} - \begin{bmatrix} 3.5 \\ 2.5 \end{bmatrix} \right) \cdot \left( \begin{bmatrix} 4 \\ 3 \end{bmatrix} - \begin{bmatrix} 3.5 \\ 2.5 \end{bmatrix} \right)^T + \left( \begin{bmatrix} 3 \\ 2 \end{bmatrix} - \begin{bmatrix} 3.5 \\ 2.5 \end{bmatrix} \right) \cdot \left( \begin{bmatrix} 3 \\ 2 \end{bmatrix} - \begin{bmatrix} 3.5 \\ 2.5 \end{bmatrix} \right)^T \right\} \\ &= \frac{1}{2} \left\{ \begin{bmatrix} 0.5 \\ 0.5 \end{bmatrix} \begin{bmatrix} 0.5 & 0.5 \end{bmatrix} + \begin{bmatrix} -0.5 \\ -0.5 \end{bmatrix} \begin{bmatrix} -0.5 & -0.5 \end{bmatrix} \right\} \\ &= \frac{1}{2} \left\{ \begin{bmatrix} 0.25 & 0.25 \\ 0.25 & 0.25 \end{bmatrix} + \begin{bmatrix} 0.25 & 0.25 \\ 0.25 & 0.25 \end{bmatrix} \right\} = \begin{bmatrix} \frac{1}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} \end{bmatrix} \end{aligned}$$

$$\vec{\mu}_2 = \frac{1}{2} \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix} + \begin{bmatrix} 1 \\ 3 \end{bmatrix} \right\} = \begin{bmatrix} 1.5 \\ 2 \end{bmatrix} = \begin{bmatrix} 3/2 \\ 2 \end{bmatrix}$$

$$\begin{aligned} \Sigma_2 &= \frac{1}{2} \left\{ \left( \begin{bmatrix} 2 \\ 1 \end{bmatrix} - \begin{bmatrix} 1.5 \\ 2 \end{bmatrix} \right) \cdot \left( \begin{bmatrix} 2 \\ 1 \end{bmatrix} - \begin{bmatrix} 1.5 \\ 2 \end{bmatrix} \right)^T + \left( \begin{bmatrix} 1 \\ 3 \end{bmatrix} - \begin{bmatrix} 1.5 \\ 2 \end{bmatrix} \right) \cdot \left( \begin{bmatrix} 1 \\ 3 \end{bmatrix} - \begin{bmatrix} 1.5 \\ 2 \end{bmatrix} \right)^T \right\} \\ &= \frac{1}{2} \left\{ \begin{bmatrix} 0.5 \\ -1 \end{bmatrix} \begin{bmatrix} 0.5 & -1 \end{bmatrix} + \begin{bmatrix} -0.5 \\ 1 \end{bmatrix} \begin{bmatrix} -0.5 & 1 \end{bmatrix} \right\} \end{aligned}$$



$$\sum_2 = \frac{1}{2} \left\{ \begin{bmatrix} 0.25 & -0.5 \\ -0.5 & 1 \end{bmatrix} + \begin{bmatrix} 0.25 & -0.5 \\ -0.5 & 1 \end{bmatrix} \right\} = \begin{bmatrix} 0.25 & -0.5 \\ -0.5 & 1 \end{bmatrix}.$$

程序代写代做 CS编程辅导



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

(5) [1 point] From what we saw in class about Linear Discriminant Functions, if we have a three-label classification case ( $\omega_1$  = "good email",  $\omega_2$  = "phishing email", and  $\omega_3$  = "ransomware email"), and we are given the weight vectors for the linear discriminant functions (Note: these weight vectors and feature vectors already include the bias integrated into them):

$$\vec{w}_1 = \begin{bmatrix} 0 \\ 2 \\ 2 \end{bmatrix} \text{ and } \vec{w}_2 = \begin{bmatrix} 4 \\ 0 \\ 2 \end{bmatrix} \text{ and } \vec{w}_3 = \begin{bmatrix} 1 \\ 5 \\ 0 \end{bmatrix} \text{ instance has the following feature vector: } \vec{x} = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}.$$

Please, determine what label our instance belongs to and predict for this new instance.



Answer:

$w_3$  "ransomware"

WeChat: cstutorcs

$$w_1: [0 \ 2 \ 2] \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} = 0 + 4 + 2 = 6$$

Assignment Project Exam Help

$$w_2: [4 \ 0 \ 2] \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} = 4 + 0 + 2 = 6$$

Email: tutorcs@163.com

$$w_3: [1 \ 5 \ 0] \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} = 1 + 10 + 0 = 11 \checkmark \text{ biggest discriminant function}$$

<https://tutorcs.com>

Good luck!!