

# 程序代写代做 CS编程辅导



Project 6  
Socialization & Matching  
Computational Social Science

## 1 Introduction

In this project, you will explore the question of whether college education causally affects political participation. Specifically, you will use replication data from [Who Matches? Propensity Scores and Bias in the Causal Effects of Education on Participation](#) by former Berkeley PhD students John Henderson and Sara Chatfield. Their paper is itself a replication study of [Reconsidering the Effects of Education on Political Participation](#) by Cindy Kam and Carl Palmer. In their original 2008 study, Kam and Palmer argue that college education has no effect on later political participation, and use the propensity score matching to show that pre-college political activity drives selection into college and later political participation. Henderson and Chatfield in their 2011 paper argue that the use of the propensity score matching in this context is inappropriate because of the bias that arises from small changes in the choice of variables used to model the propensity score. They use [genetic matching](#) (at that point a new method), which uses an approach similar to optimal matching to optimize Mahalanobis distance weights. Even with genetic matching, they find that balance remains elusive however, thus leaving open the question of whether education causes political participation.

You will use these data and debates to investigate the benefits and pitfalls associated with matching methods. Replication code for these papers is available online, but as you'll see, a lot has changed in the last decade or so of data science! Throughout the assignment, use tools we introduced in lab from the [tidyverse](#) and the [MatchIt](#) packages. Specifically, try to use dplyr, tidyr, purrr, stringr, and ggplot instead of base R functions. While there are other matching software libraries available, MatchIt tends to be the most up to date and allows for consistent syntax.

## 2 Data

The data is drawn from the [Youth-Parent Socialization Panel Study](#) which asked students and parents a variety of questions about their political participation. This survey was conducted in several waves. The first wave was in 1965 and

# 程序代写代做 CS编程辅导

established the baseline pre-treatment covariates. The treatment is whether the student attended college or not. The outcome is an index that calculates the number of political activities the student engaged in after 1965. Specifically, the key variables

- **college**. Whether the student attended college or not. 1 if the student attended college between 1965 and 1973, 0 otherwise.
- **political**. Outcome variable measuring the number of political activities the student participated in. Additive combination of whether the student voted in 1972 or 1980 (`student.vote`), attended a campaign rally or meeting (`student.meeting`), wore a campaign button (`student.button`), donated money to a campaign (`student.money`), communicated with an elected official (`student.communicate`), attended a demonstration or protest (`student.demonstrate`), was involved with a local community event (`student.community`), or some other political participation (`student.other`).

Otherwise, we also have covariates measured for survey responses to various questions about political attitudes. We have covariates measured for the students in the baseline year, covariates for their parents in the baseline year, and covariates from follow-up surveys. Be careful here. In general, post-treatment covariates will be clear from the name (i.e. `student_1973Married` indicates whether the student was married in the 1973 survey). Be mindful that the baseline covariates were all measured in 1965, the treatment occurred between 1965 and 1973, and the outcomes are from 1973 and beyond. We will distribute the Appendix from Henderson and Chatfield that describes the covariates they used, but please reach out with any questions if you have questions about what a particular variable means.

## 3 Randomization

Matching is usually used in observational studies to approximate random assignment to treatment. But could it be useful even in randomized studies? To explore the question do the following:

1. Generate a vector that randomly assigns each unit to either treatment or control.
2. Choose a baseline covariate for either the student or parent (note that a binary covariate is probably best for this exercise).
3. Visualize the distribution of the covariate by treatment/control condition. Are treatment and control balanced on this covariate?
4. Simulate the first 3 steps 10,000 times and visualize the distribution of treatment/control balance across the simulations.

# 程序代写代做 CS编程辅导

## 3.1 Questions

1. Why does independent assignment and baseline covariates not guarantee independent assignment and baseline covariates?

## 4 Propensity Score Matching

### 4.1 Overview

Select covariates that you think best represent the “true” model predicting whether a student chooses to attend college, and estimate a propensity score model to calculate the Average Treatment Effect on the Treated (ATT). Plot the balance of the covariates. Report the balance of the p-scores across both the treatment and control groups, and using a threshold of standardized mean difference of p-score  $\leq .1$ , report the number of covariates that meet that balance threshold.

### 4.2 Simulations

Henderson/Chatfield argue that an improperly specified propensity score model can actually *increase* the bias of the estimate. To demonstrate this, they simulate 800,000 different propensity score models by choosing different permutations of covariates. To investigate their claim, do the following:

- Using as many simulations as is feasible (at least 10,000 should be ok, but more is better!), randomly select the number of and the choice of covariates for the propensity score model.
- For each run, store the ATT, the proportion of covariates that meet the standardized mean difference  $\leq .1$  threshold, and the mean percent improvement in the standardized mean difference. You may also wish to store the entire models in a list and extract the relevant attributes as necessary.
- Plot all of the ATTs against all of the balanced covariate proportions. You may randomly sample or use other techniques like transparency if you run into overplotting problems. Alternatively, you may use plots other than scatterplots, so long as you explore the relationship between ATT and the proportion of covariates that meet the balance threshold.
- Finally choose 10 random models and plot their covariate balance plots (you may want to use a library like [gridExtra](#) to arrange these)

**Note:** There are lots of post-treatment covariates in this dataset (about 50!)! You need to be careful not to include these in the pre-treatment balancing. Many of you are probably used to selecting or dropping columns manually, or positionally. However, you may not

# 程序代写代做 CS编程辅导

always have a convenient arrangement of columns, nor is it fun to type out 50 d...es. Instead see if you can use dplyr 1.0.0 function... drop post-treatment variables ([here](#) is a useful ti



## 4.3 Q

1. How... resulted in models with a higher proportion... Do you have any concerns about this?
2. Analyze the distribution of the ATTs. Do you have any concerns about this distribution?
3. Do your 10 randomly chosen covariate balance plots produce similar numbers on the same covariates? Is it a concern if they do not?

## 5 Matching Algorithm of Your Choice

### 5.1 Simulate Alternative Model

Henderson/Chatfield propose using genetic matching to learn the best weights for Mahalanobis distance matching. Choose a matching algorithm other than the propensity score (you may use genetic matching if you wish, but it is also fine to use the greedy or optimal algorithms we covered in lab instead). Repeat the same steps as specified in Section 4.2 and answer the following questions:

### 5.2 Questions

1. Does your alternative matching method have more runs with higher proportions of balanced covariates?
2. Use a visualization to examine the change in the distribution of the percent improvement in balance in propensity score matching vs. the distribution of the percent improvement in balance in your new method. Which did better? Analyze the results in 1-2 sentences.

**Optional:** Looking ahead to the discussion questions, you may choose to model the propensity score using an algorithm other than logistic regression and perform these simulations again, if you wish to explore the second discussion question further.

# 程序代写代做 CS编程辅导

## 6 Discussion Questions

1. Why is it important to do matching even if we have a randomized or a controlled experiment?
2. The propensity score is estimated using a logistic regression. Given what we know about the reliability of treatment. Given what we know about the propensity score, do you think there might be advantages to using other algorithms (decision trees, bagging/boosting forests, ensembles, etc.) to estimate propensity scores instead?

WeChat: cstutorcs

Assignment Project Exam Help

Email: [tutorcs@163.com](mailto:tutorcs@163.com)

QQ: 749389476

<https://tutorcs.com>