

CDMA with Noise Solution 程序代写代做 CS编程辅导

In the class, we have considered CDMA in an ideal system. However, in reality, we have much more complicated scenario: Users will experience noise.

Consider the scenario where the receiver. The chipping sequence is $(-1 -1 -1 +1 -1 +1 +1 +1)$. Suppose the signal sent will be $(-1 -1 -1 +1 -1 +1 +1 +1)$. In the channel, noise is added. The received signal will be $(-1+n_1 -1+n_2 -1+n_3 +1+n_4 -1+n_5 +1+n_6 +1+n_7 +1+n_8)$. These terms. They are independently normally distributed with zero mean and σ^2 variance. Formally, $n_i \sim N(0, 1)$. You should know the normal distribution in a previous class.

After the computing, the receiver, what “value” does the receiver derive? If the value is smaller than 0, it is decoded as -1, otherwise, it is decoded as 1. Use the provided table to find the probability that it is wrongly decoded as -1.

The tail probability (Q function) of a standard normal distribution is given in the attached [q_function.pdf](#).

WeChat: cstutorcs


Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

$Q(a)$, the shaded area



(1) Inner product/M: $R=1+(-n_1-n_2-n_3-n_4+n_5+n_5+n_7+n_8)/8$

$n_i \sim N(0, 1)$.

$-n_i \sim N(0, 1)$.

$n_i/8 \sim N(0, 1/64)$.

$$-n_i/8 \sim N(0, 1/64).$$

程序代写代做 CS编程辅导

This is because if n is a Gaussian random variable with variance σ^2 , $a*n$ is a Gaussian random variable with variance $a^2\sigma^2$.

$$R \sim N(1, 1/8) = (\mu_0, c$$

Two independent Gaussian random variables is still a Gaussian random variable. See http://en.wikipedia.org/wiki/Sum_of_normally_distributed_random_variables



$$Y \sim N(\mu_X, \sigma_X^2)$$

$$Y \sim N(\mu_Y, \sigma_Y^2)$$

$$Z = X + Y,$$

WeChat: cstutorcs

$$Z \sim N(\mu_X + \mu_Y, \sigma_X^2 + \sigma_Y^2).$$

$$P(R < 0) = Q\left(\frac{|0 - \mu_0|}{\sigma_0}\right) = Q(2.82) \approx 2.401 \times 10^{-3}$$

Assignment Project Exam Help

It follows the definition of Q function, see <https://en.wikipedia.org/wiki/Q-function>

If Y is a Gaussian random variable with mean μ and

variance σ^2 , then $X = \frac{Y - \mu}{\sigma}$ is standard normal

and QQ: 749389476

$$P(Y > y) = P(X > x) = Q(x)$$

<https://tutorcs.com>

$$\text{where } x = \frac{y - \mu}{\sigma}.$$