

程序代写代做 CS编程辅导



Introduction

Theory of Computation

Lecture 2

WeChat: cstutorcs

Ruth Urner
Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476
January 11, 2023

<https://tutorcs.com>

程序代写代做 CS编程辅导



Basic mathematical notation and terminology

Assignment Project Exam Help
Reading: TTC Section 0.2

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Sets

程序代写代做 CS编程辅导



Set-theory is the foundation of mathematics.

We will start with the set of natural numbers as a basic given set to start with:

Assignment, Project, Exam Help

Email: tutorcs@163.com

Further, we will assume the existence of the empty set: \emptyset

<https://tutorcs.com>

程序代写代做 CS编程辅导

Sets



Set-theory is of mathematics.

We will start with the set of natural numbers as a basic given set to start with:

$\mathbb{N} = \{0, 1, 2, 3, \dots\}$

Assignment Project Exam Help

Further, we will assume the existence of the empty set: \emptyset

Email: tutorcs@163.com

\emptyset does not contain any elements

\emptyset is a ~~subset~~ \emptyset set

<https://tutorcs.com>

Elements of sets

A set consists of elements, denoted by the \in -relation:



$a \in A$

means a is an element of A .

To state that a is not an element of set A , we use the notation
WeChat: cstutorcs

$a \notin A$
Assignment Project Exam Help

Important properties
Email: tutorcs@163.com

- Sets are not ordered and contain each element only once!
QQ: 749389476
- The empty set has no elements.

<https://tutorcs.com>

程序代写代做 CS 编程辅导

Elements of sets



A set consists

denoted by the \in -relation:

$a \in A$

means a is an element of A .

To state that a is not an element of set A , we use the notation

WeChat: cstutorcs

$a \notin A$

Important properties:

- Sets are not ordered and contain each element only once!

$$\{2, 6, 4\} = \{3, 4, 15\} = \{4, 6, 23\} = \{6, 4, 2, 23\}$$

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Subsets

程序代写代做 CS编程辅导

Using the element relation, we can define what subsets are:

- $A \subseteq B$



can define what subsets are:

- $A \subsetneq B$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

- $A = B$

<https://tutorcs.com>

程序代写代做 CS编程辅导

Subsets

Photos Select Text Draw Stamp Spotlight Eraser Format Undo Auto Clear Share

Which can open on your device? Recording On



Using the elements of set B , we can define what subsets are:

- $A \subseteq B$ "A is a subset of B "
 $\{1, 2, 4\} \subseteq \{1, 2, 3\}$
 $\{1, 2, 4\} \subseteq \mathbb{N}$, $\{n \in \mathbb{N} | n \text{ even}\} \subseteq \mathbb{N}$
- $A \subsetneq B$ "A is a proper subset of B "
 $\{1, 2, 4\} \subsetneq \{1, 2, 3\}$
 $\{1, 2, 4\} \subsetneq \mathbb{N}$, $\{n \in \mathbb{N} | n \text{ even}\} \subsetneq \mathbb{N}$
- $A = B$ "A and B are equal as sets"
 $\{1, 2, 4, 2\} = \{4, 2, 1\}$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

$\{1, 2, 4, 2\} = \{4, 2, 1\}$

<https://tutorcs.com>

程序代写代做 CS 编程辅导

$A \subseteq B$

\subset symbol used to indicate that A is a proper subset of B

By default, we assume that all our sets are subsets of the natural numbers. If we assume a universe $U = \mathbb{N}$. Then we can define



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

How to define sets

程序代写代做 CS编程辅导

Here are some tools for defining sets:

1. Make a list of the elements

- ▶ Set of all students in this class
- ▶ Set of odd natural numbers smaller than 10: {1, 3, 5, 7, 9}



Problem: this technique fails for large or infinite sets

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

Problem: sometimes we don't know a precise defining characteristic

3. Inductive definition

- ▶ We'll see how to do this later

<https://tutores.com>

How to define sets – a warning

程序代写代做 CS编程辅导

Warning:

When we define sets, it is need to specify from which universe (that is a possibly much larger ground set, for example the natural numbers) the elements of our set should be taken from!



Example:

Assignment Project Exam Help

- Odd natural numbers $\{n \in \mathbb{N} \mid n \text{ is not divisible by } 2\}$
- Interval on the real line $\{x \in \mathbb{R} \mid 2 \leq x \leq 4\}$

Email: tutorecs@163.com

QQ: 749389476

Otherwise we can fall into Russell's paradox...!

<https://tutorecs.com>

Russel's paradox

程序代写代做 CS编程辅导

Consider the following definition of a set:



$$\{r \mid r \notin r\}$$

That is, the set R contains all those sets that do not contain themselves as an element.

Question: Is R an element of R ? Assignment Project Exam Help

Email: tutorcs@163.com
Now, $R \in R$ implies that $R \notin R$, and vice versa ($R \notin R$ implies that $R \in R$)— a contradiction.

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

Russel's paradox



Consider the set  definition of a set:

$$R = \{r | r \notin r\}$$

That is, the set R contains all those sets that do not contain themselves as an element.

WeChat: cstutorcs

Question: Is R an element of R ?

Assignment Project Exam Help

Now, $R \in R$ implies that $R \notin R$, and vice versa ($R \notin R$ implies that $R \in R$)—a contradiction.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Set operations

程序代写代做 CS编程辅导

By default, we will assume that all our sets are **subsets of the natural numbers**. We assume a universe $U = \mathbb{N}$. Then we can define:



- The **set-difference** $A \setminus B$ of two sets A and B

WeChat: cstutorcs

Assignment Project Exam Help

- The **intersection** $A \cap B$ of two sets A and B

Email: tutorcs@163.com
QQ: 749389476

<https://tutorcs.com>

- The **union** $A \cup B$ of two sets A and B

程序代写代做 CS编程辅导

Set operations



By default, we assume that all our sets are **subsets** of the **natural numbers**. We assume a universe $U = \mathbb{N}$. Then we can define

- The **set-difference** $A \setminus B$ of two sets A and B

$A \setminus B$ WeChat: cstutorcs

$$A = \{1, 3, 5, 7\}, B = \{1, 2, 3\}$$

$$A \setminus B = \{5, 7\}$$

always:

$$A \setminus B \subseteq A$$

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

Set operations



By default, we assume that all our sets are **subsets** of the **natural numbers**. If we assume a universe $U = \mathbb{N}$. Then we can define

- The **set-difference** $A \setminus B$ of two sets A and B

In general, **WeChat: cstutorcs**
 $A = \{1, 3, 5, 7\}, B = \{1, 2, 3\}$
 $A - B = \{5, 7\}$

Assignment Project Exam Help

Q: Is it possible to have $A - B = B - A$ for some sets A and B ?

A: Yes: **QQ: 749389476**
Or if $A = B = \{1, 2\}$, then $A - B = B - A = \emptyset$.

<https://tutorcs.com>

程序代写代做 CS 编程辅导

Set operations



By default, we assume that all our sets are **subsets** of the **natural numbers**. We assume a universe $U = \mathbb{N}$. Then we can define

- The set difference $A \setminus B$ of two sets A and B :
 $A = \{1, 3, 5, 7\}$, $B = \{1, 2, 3\}$
 $A \setminus B = \{5, 7\}$

WeChat: cstutorcs

- The intersection of two sets A and B :

$A \cap B$ is the set of all elements that are members of both A and B .

$A \cap B = \{n \in \mathbb{N} \mid n \in A \text{ and } n \in B\}$
QQ: 749389476
or universe

<https://tutorcs.com>

程序代写代做 CS编程辅导

Set operations



- The set-difference $A \setminus B$ of two sets A and B

WeChat: cstutorcs

- The intersection $A \cap B$ of two sets A and B

$$A = \{1, 3, 5, 7\}, B = \{1, 2, 3\}$$
$$A \cap B = \{1, 2, 3\}$$

Email: tutores@163.com

- The union $A \cup B$ of two sets A and B

$$A \cup B \text{ is a set of all elements in } A \text{ and all elements in } B. A \cup B = \{n \in \mathbb{N} \mid n \in A \text{ or } n \in B\}$$

QQ: 749389476
<https://tutorcs.com>

More set operations

程序代写代做 CS编程辅导

By default, we will assume that all our sets are **subsets of the natural numbers**.
can define:

- The **complement** of a set A

WeChat: cstutorcs

- The **(cartesian) product** of two sets A and B

Email: tutorcs@163.com

- The set A^k of ~~k tuples~~ of elements of A

<https://tutorcs.com>



程序代写代做 CS编程辅导

More set operations



By default, we assume that all our sets are **subsets of the natural numbers**. We can define:

- The **complement** of a set A is the set of all elements in the universe that are not in A .

$$A = \{n \in \mathbb{N} \mid n \text{ divisible by } 23\} = \{\text{even numbers}\}$$

$$\overline{A} = \{n \in \mathbb{N} \mid n \text{ is not divisible by } 23\}$$

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

More set operations



WeChat: cstutorcs

- The (cartesian) product of two sets A and B

Assignment Project Exam Help

A and B.

$A = \{1, 3\}$, $B = \{1, 2, 3\}$

Email: tutorcs@163.com


$A \times B = \{(1, 1), (1, 2), (1, 3), (3, 1), (3, 2), (3, 3)\}$

QQ: 749389476, (7, 1), (7, 2), (7, 3)

<https://tutorcs.com>

程序代写代做 CS 编程辅导

More set operations



By default, we assume that all our sets are **subsets of the natural numbers**. We can define:

- The **complement** of a set A

WeChat: cstutorcs

- The **(cartesian) product** of $A \times B$ two sets A and B

Assignment Project Exam Help

- The set A^k of k -tuples A^k of elements of A

$$A = \{(1, 1), (1, 2), (2, 1), (2, 2)\}$$

$$A^3 = \{(1, 1, 1), (1, 1, 2), (1, 2, 1), (1, 2, 2), (2, 1, 1), (2, 1, 2), (2, 2, 1), (2, 2, 2)\}$$

$$\text{QQ: } 749389476 \dots (4, 4, 4)$$

<https://tutorcs.com>

More set operations

程序代写代做 CS编程辅导

- The power set $\mathcal{P}(A)$ of set A



WeChat: cstutorcs

Assignment Project Exam Help

- The symmetric difference $A \Delta B$ between two sets A and B

QQ: 749389476

Email: tutores@163.com

<https://tutorcs.com>

程序代写代做 CS 编程辅导

More set operations

You are screen sharing Stop Share

Draw Stamp Eraser Paint Undo Redo Clear Save

And can see what you share here Recording On

Taking:

- The **pow** is tea

$A = \{\}$

$P(A) = \{\emptyset, \{\}, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$

WeChat: cstutorcs

Assignment Project Exam Help

Notes

\emptyset is a member of the powerset of every set

Email: tutorcs@163.com

Set itself is always a member of its own powerset

Warning: QQ: 749389476, $\{1\} \notin A, \{1\} \in P(A)$

<https://tutorcs.com>

程序代写代做 CS 编程辅导

More set operations

- The ~~pow~~

$$\underline{A} = \underline{\underline{2, 3}}$$



set A

$$B = \{1, 2, 3\}$$

$$A \Delta B = \{1, 5, 7\}$$

$$A \Delta B$$



WeChat: cstutorcs

A_nB

- The symmetric difference $A \Delta B$ between two sets A and B

is the set of all elements in $A \cup B$ that are

not in $A \cap B$.

Email: tutorcs@163.com

$$A \Delta B = (A \cup B) \setminus (A \cap B)$$

QQ: 749389476

<https://tutorcs.com>

Sets, multi-sets, sequences

程序代写代做 CS编程辅导

- Recall: **Sets** contain elements only once and the order does not matter



- Multi-sets** can contain elements multiple times, but do not contain an order of their elements

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

- Sequences** or **tuples** are collections of elements in a fixed order.

<https://tutorcs.com>

程序代写代做 CS 编程辅导

Sets, multi-sets, sequences



- Recall: Sets can contain an element only once and the order does not matter.

• Multi-sets can contain elements multiple times, but do not contain an order of their elements.

$\{1, 1, 3, 5\}$ is not equal to $\{1, 3, 5\}$
as a multiset, but equal as sets

$\{1, 1, 3, 5\}$ is equal to $\{1, 3, 5\}$
as a multiset.

Assignment Project Exam Help
Email: tutores@163.com
QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

Sets, multi-sets, sequences



- Recall: Sets can contain an element only once and the order does not matter.

- Multi-sets can contain elements multiple times, but do not contain an order of their elements.

Assignment Project Exam Help

- Sequences or tuples are collections of elements in a fixed order.

Email: tutorcs@163.com

$(3, 1, 5)$ is a 3-tuple

$(1, 1, 1, 2, 2, 3, 4)$ is an 8-tuple

$(2i)_{i \in \mathbb{N}}$ is the sequence of even numbers
 $\subseteq \{2, 4, 6, 8, 10, \dots\}$

<https://tutorcs.com>

Relations

程序代写代做 CS编程辅导

Definition:

A relation R between two sets A and set B is a subset of $A \times B$.



WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

We sometimes use notation aRb to state that $(a, b) \in R$.

程序代写代做 CS 编程辅导

Relations

Definition:

A relation R



" \leq " is

and set B is a subset of $A \times B$.

over the natural numbers :

$$\{(0,1), (0,0), (1,1), (0,2), (1,2), \dots\}$$

WeChat: cstutorcs

$$A = \{1, 3, 5, 7, 9\}, B = \{1, 2, 3\}, \text{ then}$$

$$R = \{(1,1), (3,3)\} \text{ is a relation between } A \text{ and } B.$$

$$R' = \{(3,2), (3,1), (3,5), (7,3)\}$$

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

Relations



Definition:
A relation R is a set of ordered pairs (a, b) where a is in set A and b is in set B is a subset of $A \times B$.

$$2 \leq 4$$

5 \leq WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

We sometimes use notation R to state that $(a, b) \in R$.

QQ: 749389476

<https://tutorcs.com>

Relations

程序代写代做 CS编程辅导



- If R is a relation between A and itself, that is $R \subseteq A \times A$, we also call R a ~~binary relation~~ or a relation of arity 2 on A .
- A subset $R \subseteq A^k$ of k -tuples of elements of A , is also called a ~~k-ary relation~~, or a relation of arity k .

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Equivalence relations

程序代写代做 CS编程辅导



Definition:

A binary relation R on some set A is called an equivalence relation if it has the following properties

- reflexive
- symmetric
- transitive

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS 编程辅导

Equivalence relations

\leq as a

is \sim

$=$ is an



is transitive but not symmetric
an equivalence relation

Definition:

A binary relation R on some set A is called an **equivalence relation** if it has the following properties:

- reflexive

for every $a \in A$, $(a, a) \in R$

- symmetric

if $(a, b) \in R$ for some $a, b \in A$, then
 $(b, a) \in R$

- transitive

if $(a, b) \in R$ and $(b, c) \in R$, then $(a, c) \in R$

$\equiv_{\text{mod} 5}$ (remainder modulo 5) is an equivalence

relation on the natural numbers

$6 \equiv_{\text{mod} 5} 11$, $6 \equiv_{\text{mod} 5} 21$, $11 \equiv_{\text{mod} 5} 21$

<https://tutorcs.com>

Talking: Ruth Urner

Functions

程序代写代做 CS编程辅导

Definition:

A **function** f from set A to set B is a relation between A and B that satisfies the following properties:



- For every $a \in A$, there exists a $b \in B$ such that $(a, b) \in f$
- If $(a, b) \in f$ and $(a, b') \in f$, then $b = b'$.

WeChat: cstutorcs
Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

Instead of $(a, b) \in f$, we usually use notation $f(a) = b$.

Alternatively, we also use the notation $f : a \mapsto b$.

<https://tutorcs.com>

程序代写代做 CS 编程辅导

Functions

i	f(i)
1	1
3	1
5	3
7	2

for functions,
we can use
such tables
to define
them

$$f(x) = |x|$$

$$g(x) = \sin(x)$$

Definition:

A function f is a relation between A and B that satisfies



Let B is a relation between A and B properties:

- For every $a \in A$, there exists a $b \in B$ such at $(a, b) \in f$
- If $(a, b) \in f$, then b is unique.

$$A = \{1, 3, 5, 7\}, B = \{1, 2, 3\}$$

$f: A \rightarrow B = \{ \text{Assignment Project Exam Help} \}$

$R = \{(1, 1), (5, 1), (1, 2), (7, 2)\}$ is not a function;

Instead of $(a, b) \in R$, we usually use notation $f(a) = b$. it violates both

Alternatively, we also use the notation $f: a \mapsto b$.

QQ: 749389476

<https://tutorcs.com>

properties (input 1
gets mapped to two outputs
and input 3 doesn't get
mapped to any output)

Functions – important properties

We use the notation 程序代写代做 CS编程辅导

$$f : A \rightarrow B$$

to state that f is a function from A to B .



We call A the domain of the function f and B the range of the function f .

WeChat: cstutorcs

Functions can be: Assignment Project Exam Help

- one-to-one: Email: tutorcs@163.com

We call a function $f : A \rightarrow B$ one-to-one if $f(a) = b$ and $f(a') = b$ for

some $b \in B$ implies that $a = a'$.

QQ: 749389476

- onto: <https://tutorcs.com>

We call a function $f : A \rightarrow B$ onto if for all $b \in B$ there exists an $a \in A$ with $f(a) = b$.