

程序代写代做 CS编程辅导



Introduction

Theory of Computation

Lecture 5

WeChat: cstutorcs

Ruth Urner
Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476
January 23, 2023

<https://tutorcs.com>

程序代写代做 CS编程辅导



Logic Reading: ITC Section 0.2
WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Boolean/Propositional logic

程序代写代做 CS编程辅导



Propositional logic means the way we reason about the **truth value** of **statements** (**propositions**).

A **propositional variable** is a variable (that stands for a statement) that can take on one of two values:

- **true** (we also use \top or 1 for true)
- **false** (we also use \perp or 0 for false)

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

A **truth assignment** is a function from a set P of propositional variables to the set of $\{0, 1\}$ (or $\{T, F\}$).

<https://tutorcs.com>

Boolean operations

程序代写代做 CS编程辅导

Given basic propositions p and q we can use **boolean operations** to derive complex statements (and their truth values):



- \neg
- \wedge
- \vee
- \rightarrow
- \leftrightarrow
- \oplus

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Truth values of composed statements

程序代写代做 CS编程辅导

We define the truth values of composed propositional statements by the following truth table:



WeChat: cstutorcs

| α | β | $(\neg\alpha)$ | $(\alpha \vee \beta)$ | $(\alpha \wedge \beta)$ | $(\alpha \rightarrow \beta)$ | $(\alpha \leftrightarrow \beta)$ | $(\alpha \oplus \beta)$ |
|----------|---------|----------------|-----------------------|-------------------------|------------------------------|----------------------------------|-------------------------|
| T | T | F | T | T | T | T | F |
| T | F | F | T | F | F | F | T |
| F | T | T | T | F | T | F | T |
| F | F | T | F | F | T | T | F |

Assignment Project Exam Help
Email: tutorcs@163.com
QQ: 749389476
<https://tutorcs.com>

Extending truth assignments to WFF – examples

程序代写代做 CS编程辅导

Say, we have propositional variables p and q , and a formula $\alpha = ((p \rightarrow q) \vee (q \rightarrow p))$.
To figure out the truth value of α under this assignment, we build a truthtable with one column for every element in a construction sequence of α as follows:



variables p and q , and a formula $\alpha = ((p \rightarrow q) \vee (q \rightarrow p))$.
a truth assignment v that sets p to T and q to F .

WeChat:tutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

We start with filling the first columns using the truth assignment v and then successively fill the other columns using the truth table for the connectives.

<https://tutorcs.com>

程序代写代做 CS编程辅导

Extending truth assignments to WFF – examples



Say, we have

$$\alpha = ((p \rightarrow q)$$

p and q to T

variables p and q , and a formula

and a truth assignment v that sets p to T and q to F

T and q to F

| <u>P</u> | <u>q</u> | $(p \rightarrow q)$ | $(q \rightarrow p)$ | $((p \rightarrow q) \vee (q \rightarrow p))$ | <u>T</u> |
|----------|----------|---------------------|---------------------|--|----------|
| T | F | T | F | T | T |
| F | T | F | T | T | F |

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

$\vee (\alpha) \dashv$
QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

Extending truth assignments to WFF – examples



Say, we have

$$\alpha = ((p \rightarrow q)$$

T and q to F

variables p and q , and a formula

and a truth assignment v that sets p to \perp

\perp and q to F

| p | q | $(p \rightarrow q)$ | $(q \rightarrow p)$ | $((p \rightarrow q) \vee (q \rightarrow p))$ |
|-----|-----|---------------------|---------------------|--|
| T | F | F | T | T |
| T | T | T | T | T |
| F | T | T | F | T |
| F | F | T | F | T |

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

$v(\alpha)$

QQ: 749389476

evaluates to T
under every truth assignment
 α is a tautology

<https://tutorcs.com>

Logical equivalence

程序代写代做 CS编程辅导



Definition: We call two propositional formulas (statements) α and β logically equivalent if they have the same truth table (that is, they evaluate to the same truth value under all truth assignments).

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Exercises

程序代写代做 CS编程辅导

Show that the following formulas α and β are logically equivalent:



- $\alpha = p$ and $\beta = (\neg(\neg p))$
WeChat: cstutorcs
- $\alpha = (p \rightarrow q)$ and $\beta = (\neg p) \vee q$
Assignment Project Exam Help
- $\alpha = (\neg(p \wedge q))$ and $\beta = ((\neg p) \vee (\neg q))$
Email: tutorcs@163.com
QQ: 749389476
- $\alpha = (\neg(p \vee q))$ and $\beta = ((\neg p) \wedge (\neg q))$
<https://tutorcs.com>

程序代写代做 CS 编程辅导

Exercises

Show that the



equivalent:

of formulas α and β are logically

If it rains then the street is wet.

- $\alpha = p$ and $\beta = (\neg(\neg p))$

If the street is not wet, I can
conclude that it's not
raining.

- $\alpha = (p \rightarrow q)$ and $\beta = ((\neg q) \rightarrow (\neg p))$

WeChat: cstutorcs
Assignment Project Exam Help

| P | $(p \rightarrow q)$ | $(\neg p)$ | $(q \rightarrow p)$ | $(\neg q)$ | $((\neg q) \rightarrow (\neg p))$ |
|---|---------------------|------------|---------------------|------------|-----------------------------------|
| T | T | F | F | F | T |
| F | F | T | T | T | F |
| T | F | F | T | F | T |
| F | T | T | F | T | F |

Email: tutorcs@163.com

QQ: 749389476 —

<https://tutorcs.com>

程序代写代做 CS 编程辅导

Exercises

Show that the



of formulas α and β are logically equivalent:

- $\alpha = p$ and $\beta = (\neg(\neg p))$

WeChat: cstutorcs

- $\alpha = (p \rightarrow q)$ and $\beta = ((\neg q) \rightarrow (\neg p))$

Exercise!

- $\alpha = (\neg(p \wedge q))$ and $\beta = ((\neg p) \vee (\neg q))$

Assignment Project Exam Help !
prove !

- $\alpha = (\neg(p \vee q))$ and $\beta = ((\neg p) \wedge (\neg q))$

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

More exercises

程序代写代做 CS编程辅导

Show that the following pairs of formulas α and β are logically equivalent:



- $\alpha = (p \vee q)$ and $\beta = ((\neg p) \wedge (\neg q))$

WeChat: cstutorcs

- $\alpha = (p \rightarrow q)$ and $\beta = ((\neg p) \vee q)$

Assignment Project Exam Help

- $\alpha = (\neg(p \rightarrow q))$ and $\beta = (\neg p \wedge (\neg q))$

Email: tutorcs@163.com
QQ: 749389476

<https://tutorcs.com>

- $\alpha = (p \oplus q)$ and $\beta = (\neg(p \leftrightarrow q))$

程序代写代做 CS 编程辅导

More exercises

Show that the
equivalent:



of formulas α and β are logically

- $\alpha = (p \vee q)$ and $\beta = ((\neg p) \wedge (\neg q))$

- $\alpha = (p \rightarrow q)$ and $\beta = (\neg p \vee q)$

- $\alpha = (\neg(p \rightarrow q))$ and $\beta = ((p \rightarrow q) \wedge (\neg q))$

- $\alpha = (p \leftrightarrow q)$ and $\beta = ((p \rightarrow q) \wedge (q \rightarrow p))$

- $\alpha = (p \oplus q)$ and $\beta = (\neg(p \leftrightarrow q))$

Exercise :

prove these
equivalences.

WeChat: estutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

More exercises

程序代写代做 CS编程辅导

Show that the following two formulas α and β are **not logically equivalent**:



- $\alpha = (p \vee q)$ and $\beta = (\neg(p \wedge q))$

WeChat: cstutorcs

- $\alpha = (p \rightarrow q)$ and $\beta = ((\neg p) \rightarrow (\neg q))$

Email: tutorcs@163.com

QQ: 749389476

- $\alpha = (p \oplus q)$ and $\beta = (p \vee q)$

<https://tutorcs.com>

程序代写代做 CS 编程辅导

More exercises

Show that the
equivalent:



of formulas α and β are not logically equivalent

- $\alpha = (p \vee q)$ and $\beta = (\neg p \wedge q)$

WeChat: cstutorcs

- $\alpha = (p \rightarrow q)$ and $\beta = ((\neg p) \rightarrow (\neg q))$

Assignment Project Exam Help

| P | q | $(p \rightarrow q)$ | $(\neg p)$ | $(\neg q)$ | $(\neg p) \rightarrow (\neg q)$ |
|---|---|---------------------|------------|------------|---------------------------------|
| T | T | T | F | F | T |
| T | F | F | F | T | F |
| F | T | T | T | F | T |
| F | F | T | T | T | T |

Email: tutores@163.com

QQ: 749389476

<https://tutorcs.com>



Talking: Ruth Urner

of formulas α and β are not logically equivalent

β is logically equivalent to $\gamma = (q \rightarrow p)$. Thus we now also show that α and β are not logically equivalent.

there is a truth assignment for which the truth values of the two formulas differ, hence they are not logically equivalent.

More exercises: Show the distributive laws

程序代写代做 CS编程辅导

Show that the following formulas α and β are logically equivalent:



- $\alpha = (p \wedge (q \vee r))$ and $\beta = ((p \wedge q) \vee (p \wedge r))$

WeChat: cstutorcs

Assignment Project Exam Help

- $\alpha = (p \vee (q \wedge r))$ and $\beta = ((p \vee q) \wedge (p \vee r))$

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

These two logical equivalences are called **distributive laws** of propositional logic.

程序代写代做 CS 编程辅导

More exercises: Show the distributive laws



Show that the
equivalent:

of formulas α and β are logically
equivalent:
*have identical evaluations for all
truth assignments*

- $\alpha = (p \wedge (q \vee r))$ and $\beta = ((p \wedge q) \vee (p \wedge r))$

| P | q | r | $(p \wedge q)$ | $(p \wedge r)$ | $(p \wedge (q \vee r))$ | $((p \wedge q) \vee (p \wedge r))$ |
|---|---|---|----------------|----------------|-------------------------|------------------------------------|
| T | T | T | T | T | T | T |
| T | T | F | F | F | F | F |
| T | F | T | F | F | F | F |
| T | F | F | F | F | F | F |
| F | T | T | T | T | T | T |
| F | T | F | F | F | F | F |
| F | F | T | F | F | F | F |
| F | F | F | F | F | F | F |

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

$= 2^3$
8 possible
truth
assignments
for 3
variables

Tautologies and contradictions

程序代写代做 CS编程辅导

We call a propositional formula a **tautology** if it evaluates to **true** under every truth assignment.



WeChat: cstutorcs

Assignment Project Exam Help

We call a propositional formula a **contradiction** if it evaluates to false under every truth assignment.

Email: cstutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

Tautologies and contradictions



We call a propositional formula a **tautology** if it evaluates to **true** under every truth assignment.

Example: $(P \rightarrow P)$

WeChat: cstutorcs

We call a propositional formula a **contradiction** if it evaluates to **false** under every truth assignment.

Example: $(P \wedge \neg P)$

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

Extending truth assignments to WFF – examples



Say, we have

$$\alpha = ((p \rightarrow q)$$

T and q to F

variables p and q , and a formula

and a truth assignment v that sets p to \perp

\perp and q to F

| p | q | $(p \rightarrow q)$ | $(q \rightarrow p)$ | $((p \rightarrow q) \vee (q \rightarrow p))$ |
|-----|-----|---------------------|---------------------|--|
| T | F | F | T | T |
| T | T | T | T | T |
| F | T | T | F | T |
| F | F | T | F | T |

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

$v(\alpha)$

QQ: 749389476

evaluates to T
under every truth assignment
 α is a tautology

<https://tutorcs.com>

First order/Predicate logic—motivation

程序代写代做 CS编程辅导

Assumptions:

- All people are mortal
- I am a person

(Sad) Conclusion:

- I am mortal



WeChat: cstutorcs

Question: Can propositional logic explain the pattern used in this example of reasoning?
Assignment Project Exam Help

Answer: No. Propositional logic can only relate truths or falsehood of statements as a whole. It does not provide as a means of reasoning about objects and properties that these objects may have.
Email: tutorcs@163.com
QQ: 749389476
<https://tutorcs.com>

First order/Predicate logic—motivation

Assumptions:

- All people are mortal
- I am a person

(Sad) Conclusion:

- I am mortal

程序代写代做 CS编程辅导



To explain this example of reasoning, we need means to refer to the inner structure of these statements (not just the statements as a whole). First order logic allows us to

WeChat: cstutorcs
Assignment Project Exam Help

- refer to **objects** (for example people)
- state that objects have certain **properties** (for example being mortal)
- make statements about **relationships** between objects
- **quantify** over objects (for example state that something holds **for all** (all people are mortal) objects or that **there exists** an object that has a certain property)

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

First order/Predicate logic–quantifiers

程序代写代做 CS编程辅导

In mathematical reasoning we make statements that **quantify** over objects, for example:



- **Definition:** A graph (V, E) is connected, if **for every** pair of vertices $v_1, v_2 \in V$ **there exists** a path from v_1 to v_2 .

WeChat: cstutorcs

- To express that there are infinitely prime numbers, we say:

For every prime number p , **there exists** a prime number p' with $p' > p$.

Email: tutorcs@163.com

QQ: 749389476

- If I want to show that some propositional formula is not a tautology, I need to show that **there exists** truth assignment to its variables, for which α evaluates to false.

<https://tutorcs.com>

First order/Predicate logic–quantifiers

程序代写代做 CS编程辅导

In general, I recommend you spell out mathematical statements and proofs in English language.



Sometimes, it is useful to have some shorthand for statements. We use

WeChat: cstutorcs

- the symbol \forall to stand for for all and
- the symbol \exists to stand for there exists.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

First order/Predicate logic–negating statements with quantifiers

程序代写代做 CS编程辅导

It is important to remember that when we **negate** statements in first order logic **the two quantifiers change roles**:



- $\neg(\forall x P(x))$ is equivalent to $\exists x (\neg P(x))$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

- $\neg(\exists x P(x))$ is equivalent to $\forall x (\neg P(x))$

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS 编程辅导

First order/Predicate logic—negating statements with quantifiers



It is important to note that, when we **negate** statements in first order logic the quantifiers \forall and \exists “change roles”:

- $\neg(\forall x P(x))$ is equivalent to $\exists x (\neg P(x))$

Example: WeChat: cstutorcs
having population larger than 1 000 000.

It's not true that all cities

population larger than 1000 000.

Email: tutorcs@163.com

There exist a city with less than

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS 编程辅导

First order/Predicate logic—negating statements with quantifiers



It is important to note that, when we **negate** statements in first order logic the symbols \forall and \exists “change roles”:

- $\neg(\forall x P(x))$ is equivalent to $\exists x (\neg P(x))$

WeChat: cstutorcs

Assignment Project Exam Help

- $\neg(\exists x P(x))$ is equivalent to $\forall x (\neg P(x))$

There does not exist a student in my class
that is confused.
All students in my class are not confused.

<https://tutorcs.com>

First order/Predicate logic–negating statements with quantifiers

程序代写代做 CS编程辅导

Example: The definition of continuity for a function $f : \mathbb{R} \rightarrow \mathbb{R}$ at a point $x_0 \in \mathbb{R}$ can be compactly stated as follows:



$$\forall \epsilon > 0 \exists \delta > 0 \forall x \in \mathbb{R} (|x - x_0| < \delta \rightarrow |f(x) - f(x_0)| < \epsilon)$$

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com


To prove that function $f : \mathbb{R} \rightarrow \mathbb{R}$ is **not continuous** in x_0 , we need to show:

$$\exists \epsilon > 0 \forall \delta > 0 \exists x \in \mathbb{R} (|x - x_0| < \delta \wedge |f(x) - f(x_0)| \geq \epsilon)$$

<https://tutorcs.com>

程序代写代做 CS编程辅导

First order/Predicate logic—negating statements with quantifiers



Example: The continuity of a function $f : \mathbb{R} \rightarrow \mathbb{R}$ at a point $x_0 \in \mathbb{R}$ can be defined as follows:

$$\forall \epsilon > 0 \exists \delta > 0 \quad |x - x_0| < \delta \rightarrow |f(x) - f(x_0)| < \epsilon$$



WeChat: cstutorcs

Assignment Project Exam Help

To prove that function $f : \mathbb{R} \rightarrow \mathbb{R}$ is not continuous in x_0 , we need to show:

$$\exists \epsilon > 0 \forall \delta > 0 \exists x \in \mathbb{R} \quad (|x - x_0| < \delta \wedge |f(x) - f(x_0)| \geq \epsilon)$$

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导



Mathematical statements and proofs

Reading: WeChat: cstutorcs 0.3 andd 0.4

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Basic elements of mathematical text

程序代写代做 CS编程辅导

Definition

A statement that clearly defines an object/structure/concept based on previously defined terms.



Theorem

WeChat: cstutorcs

A statement that has been proven to be true.

Assignment Project Exam Help

Proof

Email: tutorcs@163.com

A clean, deductive argument for why a statement is true.

QQ: 749389476

Lemma

<https://tutorcs.com>

A “helper theorem”, typically only stated as a step in a proof of some theorem.

Examples of what not to do..

程序代写代做 CS编程辅导

Definition
A statement defines an object/structure/concept based on previous terms.

Warning: definitions can not be circular!

WeChat: cstutorcs

Example: Define $x > y$ as "x-y is positive"

Assignment Project Exam Help

Define x is positive as $x > 0$

Email: tutores@163.com

QQ: 749389476

<https://tutorcs.com>

Examples of what not to do..

程序代写代做 CS编程辅导

Definition
A statement defines an object/structure/concept based on previous terms.

Counter-example to the claim:
9 is odd
and
 $9 = 3 \cdot 3$ is not prime.
Thus the claim is false.

Theorem WeChat: cstutorcs
A statement that has been proven to be true.

Proof A clean, deductive argument for why a statement is true.

~~Email: tutorcs@163.com~~

What not do: proof by example

Example: ~~Claim: "All odd natural numbers are prime"~~

~~"Proof" (by example): 1, 3, 5, 7, ...~~

To refute a statement by giving a counter-example is a sufficient proof.

<https://tutorcs.com>

Some comments

- Understanding a statement is not the same as understanding why the statement is true (or false). The first step in attempting to prove a statement, is always **sure you understand the statement fully.**
- When you attempt something, I recommend to always first develop an **intuition** about the statement and what may be the proof. Eg, first come up with some simple **examples to illustrate the statement**, then **develop an intuition** for why the statement is true, then **develop a proof** for it.



WeChat: estidores

Assignment Project Exam Help

- Understanding whether a proof is correct and complete, is an important skill. It's **important** that you learn to evaluate whether your own proofs are correct.

QQ: 749389476

- If you want to prove a statement, you need to **provide a general argument**. If you want to disprove a statement, you need to present a **counter-example**.
- Learning to prove mathematical statements is a skill that develops with practice. **Be patient with yourself :)**

Email: tutorcs@163.com

https://tutorcs.com

Types of proofs—constructive proofs

程序代写代做 CS编程辅导



If the statement that we are aiming to proof is a claim about **WeChat: cstutorcs** **existence** of some object, then often we can prove the statement by constructing such an object. **Assignment Project Exam Help**

Email: **tutorcs@163.com**

QQ: **749389476**

<https://tutorcs.com>

Types of proofs—constructive proof example

Definition

程序代写代做 CS编程辅导

For a natural number $k \in \mathbb{N}$, we call a graph $G = (V, E)$ a **k -regular graph** if every vertex in V has degree k .



Theorem

For every even natural number $n \geq 4$, there exists a 3-regular graph with n vertices.

Assignment Project Exam Help

Email: tutorcs@163.com

Proof

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS 编程辅导

Types of proofs-construction

Definition

For a natural
k-regular graph



we call a graph $G = (V, E)$ a
k-regular graph if every vertex in V has degree k .



↑
not a k-regular
graph
for any k

for claims of existence,
often "proof by
construction" is useful
technique

Theorem

For every even natural number $n \geq 4$, there exists a 3-regular graph with n vertices.

WeChat: cstutorcs

proof (by construction).

Let $n \geq 4$ be an even natural number.

vertex set $V = [n] = \{1, 2, \dots, n\}$.


We define the following edges:

$$E = \{\{i, i+1\} \mid i \in \{1, \dots, n-1\}\} \cup \{\{n, 1\}\}$$

$$\cup \left\{ \left\{ i, \frac{n}{2} + i \right\} \mid i \in \{1, 2, \dots, \frac{n}{2}\} \right\}$$

This results in a graph where every vertex has degree 3.

<https://tutorcs.com>



Types of proofs—“by way of contradiction”

程序代写代做 CS编程辅导



Sometimes, in order to prove that some statement is true, we assume that the statement is false and then show that this assumption leads to a contradiction. This in turn implies that the statement is true.

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS编程辅导

Types of proofs—“by way of contradiction”



Scalable Vector Graphics (SVG) Editor
Save As... Share Recordings

Saved as PNG Show in Folder

Talking: Ruth Urner

Sometimes, in order to prove that some statement is true, we assume that the statement is false and then show that this assumption leads to a contradiction. This, in turn, implies that the statement is true.

To prove statement P , we assume $(\neg P) \Rightarrow F$ is a true statement.

$(\neg P) \Rightarrow F$ is a true statement.
Email: tutorcs@163.com

This then lets us conclude that $(\neg P)$ must be false, and thus the statement P must be true.

QQ: 749389476

<https://tutorcs.com>

Types of proofs— examples of proofs “by way of contradiction”

程序代写代做 CS编程辅导

Theorem

$\sqrt{2}$ is not a rational



Proof

See textbook.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Types of proofs— examples of proofs “by way of contradiction”

程序代写代做 CS编程辅导

Theorem

There are infinitely many prime numbers.



Proof

WeChat: cstutorcs

Assignment Project Exam Help


Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

程序代写代做 CS 编程辅导

Types of proofs—examples of proofs “by way of contradiction”



Theorem

There are infinitely many numbers.

Proof

B.w.o.c. (by way of contradiction), let's assume there are only finitely many primes, that is that there are only finitely, say $n \in \mathbb{N}$, many prime numbers $p_1, p_2, p_3, \dots, p_n$.
(and let's assume $p_i > 1$ for all i).

Now, let's consider the number

$$N = (p_1 \cdot p_2 \cdot p_3 \cdots \cdot p_n) + 1$$

We have $N \neq p_i$ for all i , and none of the p_i divides

N . Thus N must be a new prime number, a contradiction

+ to the assumption that there are only n primes

<https://tutorcs.com>

Talking: Ruth Urner

Thus the assumption was false, and therefore there aren't infinitely many prime numbers.

Types of proofs— examples of proofs “by way of contradiction”

程序代写代做 CS编程辅导

Theorem

The set \mathbb{R} of real numbers is not countable.



Proof

We'll prove this in the tutorial on Friday.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Types of proofs—proof by (structural) induction

程序代写代做 CS编程辅导



We can use a proof by induction, if we want to prove a statement about elements of a set that is (or can be) defined inductively.
Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Inductive definition of sets – motivating example

程序代写代做 CS编程辅导

Say, I'd like to define all (biological) relatives of mine
(living and dead ones)



- I can not make

(I don't know them all, especially not those that lived a thousand years ago..)

WeChat: cstutorcs

- I can not give a precise characteristic

Assignment Project Exam Help

(Maybe I could if I was a biologist, but I am not..)

- But I know some operations that will allow me to get from me to all of them!

QQ: 749389476

The idea is to start with me, and consider everyone that can be reached by successively considering all children and all parents of previously reached people.

<https://tutorcs.com>

Inductive definition of sets – motivating example

程序代写代做 CS编程辅导



Here is a more formal way of defining all my biological relatives:

Consider the following three components:

1. **Universe:** all people
2. **Core set:** me
3. **Operations:** parent-of, child-of

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

The set of all my relatives: Start with me, and successively apply the operations parent-of and child-of. The set of all my relatives are all people that can be “reached” this way.

QQ: 749389476

<https://tutorcs.com>

Inductive definition of sets – general pattern

程序代写代做 CS编程辅导

An inductive definition consists of

1. A universe set 
 2. A core set $C \subseteq U$
 3. A finite set $O = \{o_1, o_2, \dots, o_n\}$ of operations from
 $o_i : U^{r_i} \rightarrow U$ for some arities $r_i \in \mathbb{N}$
- WeChat: cstutorcs
Assignment Project Exam Help

Email: tutorcs@163.com

We define $\mathcal{I}(U, C, O)$ as the set of elements that we obtain by
starting with the core set and putting all those elements of U into
 $\mathcal{I}(U, C, O)$ that one can reach by successively applying the
operations in O .

QQ: 749389476

<https://tutorcs.com>

Structural induction—general definition

程序代写代做 CS编程辅导

Consider some induction hypothesis on a defined set $\mathcal{A} = \mathcal{I}(U, C, O)$. To show that all elements of \mathcal{A} satisfy a certain property \mathcal{P} we prove the following:



Base case Show that all elements $c \in C$ of the core set satisfy the property \mathcal{P} .

WeChat: cstutorcs

Induction hypothesis Assume that some $a_1, a_2, \dots, a_n \in I(U, C, O)$ satisfy the property (where n is the largest arity of the operations in O).

Assignment Project Exam Help

Emails tutorcs@163.com

Induction step Show that for all operation $o_i \in O$, if the induction hypothesis holds, then the property also holds for

QQ: 749389476
<https://tutorcs.com>

$o_i(a_1, a_2, \dots, a_{r_i})$.

Structural induction

程序代写代做 CS编程辅导



- When proving something by (structural) induction, it is very important that you clearly state the hypothesis and make it clear to yourself where in the induction step you are actually using it. If it is not clear where you use it, there is likely something wrong with your proof..!

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Structural induction-example

程序代写代做 CS编程辅导

Game with cups

We consider three cups on a table as follows:



∩ U

(That is, two upright and the middle one upside down.)

WeChat: cstutorcs

- We can now play with the cups by, at each step, flipping exactly two of them
- Eg, flipping the two left ones results in

Email: tutorcs@163.com

QQ: 749389476

Question: Can we, by repeatedly flipping two cups, end up with all cups upright
<https://tutorcs.com>

U U U?

Structural induction-example

程序代写代做 CS编程辅导



First, we note that we can define the set of all reachable cup-configurations as a recursively defined set:

- **Universe:** $U_c = \text{All ways to place three cups on the table.}$
WeChat: cstutorcs
(Question for you: How big is this universe?)
- **Coreset:** The initial configuration, $C_c = \{\text{UUU}\}$
- **Operations:** $O_c = \{\text{flip-left-two, flip-outer-two, flip-right-two}\}$
Email: tutorcs@163.com

Question: Is $\text{UUU} \in \mathcal{I}(U_c, C_c, O_c)$?

QQ: 749389476
<https://tutorcs.com>

程序代写代做 CS 编程辅导

Structural induction-example



First, we note that we want to define the set of all reachable cup-configurations. This is an inductively defined set:

- **Universe:** $U_c = \text{All ways to place three cups on the table.}$

(Question for you: How many ways are there?)

- **Coreset:** The initial configuration, $C_c = \{U \cap U\}$

- **Operation:** $O_c = \{\text{flip-left-two}, \text{flip-outer-two}, \text{flip-right-two}\}$

→ This defines (inductively) the set of all reachable

Question: Is $|U| \leq |L^*(U, C_c, O_c)|?$ (This is a game.)

Email: tutorcs@163.com

→ No!

|| we'll prove that the number //
of upside cups is always even. ||

<https://tutorcs.com>



Structural induction-example

程序代写代做 CS编程辅导



Conjecture: It is not possible to get all cups upright..

We will prove the following property by induction:
In all reachable states, the number of upright cups is even.
Assignment Project Exam Help

Email: tutorcs@163.com

Since $\cup\cup\cup$ has an odd number of upright cups, this will imply
QQ: 749389476
that this state is not reachable.

<https://tutorcs.com>

程序代写代做 CS 编程辅导

Structural induction-example

2) if we flip two up cups, then XYZ must have had two up (since now) plus we end up with zero up-cups.
3) if we flip two down cups, XYZ must have had one up and one up with 2 up cups.



Proof by

Base case



Initial configuration
XYZ, there are two (i.e even number of) up-cups.

WeChat: cstutorcs

Induction hypothesis: Assume XYZ is a sequence of n up-cups and n down-cups of up-cups.

Induction step: If we flip two up XYZ, there

are three cases: 1) if we flip one up and one down, the number of up-cups stays the same.

QQ: 749389476

<https://tutorcs.com>

The base property is maintained under the operation, and therefore holds for all subsequent configurations by induction.

Structural induction-example

程序代写代做 CS编程辅导



Property: The number of up-cups is even.

Proof by induction WeChat: cstutorcs

Base case In the initial configuration $U_0 C_0 O_0$ the property holds
(2 cups are up, which is even).

Email: tutorcs@163.com

Induction hypothesis Assume that for some configuration

QQ: 749389476

$XYZ \in \mathcal{I}(U_c, C_c, O_c)$ the number of up-cups is even.

<https://tutorcs.com>

Structural induction-example

程序代写代做 CS编程辅导

Induction step First, note that if the number of up-cups in XYZ is even, it is either 0 or 2 (since 0 and 2 are the only even numbers smaller than 3). This observation motivates the following case distinction:



Case 1: It is 0 Then flipping two cups results in 2 up-cups, which is even again.

WeChat: cstutorcs

Case 2: It is 2 Then we either flip the two up-cups in XYZ or we flip one up-cup and one down-cup. In the first case, we end up with 0 up-cups, which is even, in the second case, we maintain 2 up-cups.

Email: tutorcs@163.com

Thus in all cases, the number of up-cups in $\text{flip-left-two}(XYZ)$, $\text{flip-outer-two}(XYZ)$, $\text{flip-right-two}(XYZ)$ is even again.

<https://tutorcs.com>

Question for you: Where did we use the induction hypothesis?

Types of proofs—proof by induction

程序代写代做 CS编程辅导



When we use a structural induction proof for the set of natural numbers, we often simply call it “proof by induction”
Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>

Example of proof by induction for natural numbers

Theorem

程序代写代做 CS编程辅导

For a natural number $n \geq 1$, we let $S(n)$ denote the sum of the natural numbers up to n . Then the following equality holds:



$$\frac{1}{2} \cdot n \cdot (n + 1)$$

Proof

This is part of exercise 0.11, and we'll prove it in the Tutorial on Friday.

WeChat: cstutorcs

Assignment Project Exam Help

Email: tutorcs@163.com

QQ: 749389476

<https://tutorcs.com>