

# Bang-Bang principle for controls in a simplex

Alexander Gudev

**Definition.** Let  $I \subset \mathbb{R}$  be an interval,  $A \in L_1(I, M_{n \times n})$ . *Fundamental matrix* of the linear system

$$\dot{x}(t) = A(t)x(t) \quad (1)$$

is any matrix function  $\Phi : I \rightarrow M_n$ , whose columns are linearly independent solutions to the system.

*Remark 1.* All solutions are expressible in terms of  $\Phi$ ,

$$x(t) = \sum_i \alpha_i \Phi_i(t) = \Phi(t)\alpha, \quad \alpha \in \mathbb{R}^n.$$

Provided  $\det \Phi(t) \neq 0$  for all  $t \in I$ , each column of  $\Phi(t)$  separately is a solution to (1) iff  $\dot{\Phi}(t) = A(t)\Phi(t)$ . That is, the fundamental matrices of (1) are determined by

$$\dot{\Phi}(t) = A(t)\Phi(t).$$

Everywhere below we assume  $\Phi(t_0) = E$ ; then the vector  $\alpha$  above is actually the initial condition  $\alpha = x(t_0)$ .

In this sense the matrix  $\Phi(t)$  „walks“ the solution *forward* as time passes from  $t_0$  to  $t$ .

*Remark 2.* It is natural to expect  $\Phi^{-1}(t)$  to walk the solution *backward*, i.e. to solve the system with  $(-A)$  (opposite velocity  $\dot{x}$ ). Indeed, differentiating  $\Phi^{-1}\Phi = E$  we get  $\frac{d\Phi^{-1}}{dt}\Phi + \Phi^{-1}\dot{\Phi} = 0$ , from where  $\frac{d\Phi^{-1}}{dt}\Phi = -\Phi^{-1}A\Phi$  and

$$\frac{d\Phi^{-1}}{dt} = -\Phi^{-1}A. \quad (2)$$

*Remark 3.* Below at each point in time  $t \in I$  the state of the system is a vector  $x(t) \in \mathbb{R}^n$ , and the control is a vector  $u(t) \in \mathbb{R}^m$ .

**Problem.** Let  $A \in L_1(I, M_{n \times n})$  and  $B \in L_1(I, M_{n \times m})$ . Solve the system

$$\dot{x}(t) = A(t)x(t) + B(t)u(t)$$

in terms of the fundamental matrix  $\Phi(t)$  of  $\dot{x}(t) = A(t)x(t)$ .

**Solution** (wiki books: Control Systems/Linear System Solutions). Rewrite  $\dot{x} - Ax = Bu$ , multiply by  $\Phi^{-1}(t)$ , and integrate:

$$\begin{aligned} \Phi^{-1}\dot{x} + \Phi^{-1}(-A)x &= \Phi^{-1}Bu \\ \frac{d}{dt}(\Phi^{-1}x) &= \Phi^{-1}Bu \quad (\text{by (2)}) \\ \Phi^{-1}x &= x(t_0) + \int_{t_0}^t \Phi^{-1}(s)B(s)u(s) ds, \end{aligned}$$

from where

$$x(t) = \Phi(t)x(t_0) + \Phi(t) \int_{t_0}^t \Phi^{-1}(s)B(s)u(s) \, ds.$$

Below for convenience we will assume  $x(t_0) = 0$ , in which case  $x(t) = \int_{t_0}^t \Phi^{-1}(s)B(s)u(s) \, ds$ .

**Theorem** (Принципът Бум-трас). *Let  $C \subset \mathbb{R}^m$  be a convex compact (allowed control values). For the system*

$$\begin{aligned} \dot{x} &= A(t)x(t) + B(t)u(t) \\ x(0) &= \vec{0} \end{aligned}$$

*every point accessible with some control  $u : I \rightarrow C$  in time  $t$  is accessible with an extremal control  $u^0 : I \rightarrow \text{Ext } C$  as well.*

We won't prove it in full generality, but only in the special case when the allowed control values form the unit ball (Holmes) or a simplex (Hermes).

**Theorem 1** (Holmes). *Let  $Y \in L_1^{loc}(I, M_{n \times m})$  have columns  $Y_i$ . For each control  $u \in \Psi$  where*

$$\begin{aligned} \Psi &:= L^\infty(I, \{\|x\|_\infty \leq 1\}) = (\Psi')^m \\ \Psi' &:= L^\infty(I, [-1, 1]), \end{aligned}$$

*there exists  $v \in \Psi$  with  $\text{Im } v \subseteq \{-1, 1\}^m$  s.t.  $\int_I Y \cdot u = \int_I Y \cdot v$  (here  $Y \cdot u$  denotes matrix-vector multiplication).*

*Proof.* We want  $\int_I Y \cdot u = \int_I Y \cdot v$ , but with  $|v_j(t)| \stackrel{\text{a.e.}}{=} 1$ . For  $j = 1, \dots, m$  we are looking for a partition  $I = B_j \cup \bar{B}_j$  with  $\begin{matrix} v_j|_{B_j} \equiv +1 \\ v_j|_{\bar{B}_j} \equiv -1 \end{matrix}$ . Equivalently,

$$\begin{aligned} \int_I u_j Y_j &= \int_I v_j Y_j = \int_{B_j} Y_j - \int_{\bar{B}_j} Y_j \\ &= \int_{B_j} Y_j - \left( \int_I Y_j - \int_{B_j} Y_j \right) \\ &= 2 \int_{B_j} Y_j - \int_I Y_j, \end{aligned}$$

that is,

$$\int_{B_j} Y_j = \frac{1}{2} \int_I (u_j + 1) Y_j. \quad (3)$$

We want to prove that to each  $u_j \in \Psi'$  there corresponds a measurable  $B_j \subseteq I$ , satisfying the last equality. To express more clearly the relationship between the given  $u_j$  and the  $B_j$  looked for, we introduce the mappings, defining their part in the respective sides in the equality.

- For  $B_j$  this is the vector measure

$$\vec{\mu}_j(E) := \int_E Y_j, \quad E \in \mathcal{L}_I,$$

where  $\mathcal{L}_I = \{E \subset I : E \text{ is measurable}\}$ .

- For  $u_j$  this is the affine operator  $T_j : \Psi' \rightarrow \mathbb{R}^n$ ,

$$T_j(y) := \frac{1}{2} \int_I (y(t) + 1) d\vec{\mu}_j, \quad y \in \Psi'.$$

$T_j$  is  $w^*$ -continuous, since for its linear part

$$L_\infty \ni y \mapsto \int_I y Y_j \leq \|y\|_\infty \vec{\mu}_j(I) \xrightarrow{\|y\| \rightarrow \infty} 0.$$

Thus the equality (3) can be expressed with the formula  $\vec{\mu}_j(B_j) = Tu_j$ , and the statement we want to prove becomes the inclusion

$$T_j \Psi' \subset \vec{\mu}_j(\mathcal{L}_I).$$

By Lyapunov's theorem the RHS  $\vec{\mu}_j(\mathcal{L}_I)$  is a convex compact, and since  $\Psi'$  is obviously convex and  $T_j$  is linear,  $T_j \Psi'$  is convex as well. Then it suffices to show  $T_j(\text{Ext } \Psi') \subset \vec{\mu}_j(\mathcal{L}_I)$ .

Let  $u \in \text{Ext } \Psi'$ , then  $u = \chi_E$  for some  $E \subset I$  and

$$\begin{aligned} T_j u &= \frac{1}{2} \int_I (\chi_E + 1) Y_j \\ &= \int_E Y_j + \frac{1}{2} \int_{\overline{E}} Y_j \\ &= \vec{\mu}_j(E) + \frac{1}{2} \vec{\mu}_j(\overline{E}). \end{aligned}$$

Let  $\overline{E}' \subset \overline{E}$  have  $\vec{\mu}_j(\overline{E}') = \frac{1}{2} \vec{\mu}_j(\overline{E})$ . Then

$$T_j u = \vec{\mu}_j(E \cup \overline{E}').$$

□

**Definition 1.** From here on we fix a closed bounded interval  $I$ , the convex compact

$$\mathcal{C} := \{x \in \mathbb{R}_{\geq 0}^m : \sum_i x_i = 1\},$$

as well as a matrix-valued function  $Y \in L_1(I, M_{n \times m})$  (in the system above,  $Y(s) := \Phi^{-1}(s)B(s)$ ).

*Remark.*  $\mathcal{C}$  is actually a **simplex**, since for  $f_j \in (\mathbb{R}^m)^\star, j = 0, \dots, m$ ,

$$\begin{aligned} f_0(x_1, \dots, x_m) &= \sum_i x_i \\ f_i(x_1, \dots, x_m) &= x_i, \quad i = 1, \dots, m \end{aligned}$$

we can express  $\mathcal{C} = \bigcap_{i=1}^m f_i^{-1}[0, \infty) \cap f_0^{-1}(1)$ , that is,  $\mathcal{C}$  is an intersection of closed half-spaces.

Since  $\mathcal{C}$  is a simplex,  $\text{Ext } \mathcal{C}$  consists of  $m$  **corners** we denote by  $r_1, \dots, r_m$ .

In the proof of the Bang-bang principle later we'll refer to the following separability proposition:

**Lemma 1.** Let  $E \subset \mathbb{R}$  be a non-null set and  $u(t) \in \mathbb{R}^m \setminus \mathcal{C}$  a.e. in  $E$ . Then there exists a vector  $\eta \in \mathbb{R}^m$ , separating strictly  $\mathcal{C}$  from  $u(t)$  a.e. in a non-null  $E' \subset E$ , that is,

$$\exists \varepsilon, B > 0, \eta \in \mathbb{R}^m : \begin{cases} \langle \eta, x \rangle \geq B & \forall x \in \mathcal{C} \\ \langle \eta, u(t) \rangle \leq B - \varepsilon & \forall t \in E' \end{cases}$$

*Proof.* According to the previous remark we can represent with a countable union

$$\begin{aligned} \mathbb{R}^m \setminus \mathcal{C} &= \bigcup_{i=1}^{m+2} g_i^{-1}(-\infty, a_i) \\ &= \bigcup_{i=1}^{m+2} \bigcup_{k \in \mathbb{N}} g_i^{-1}(-\infty, a_i - 1/k], \end{aligned}$$

where  $g_i \in (\mathbb{R}^m)^*$ ,  $a_i \in \mathbb{R}$ ,  $i = 1, \dots, m+2$ . Then the image under  $u$  of some non-null  $E'$  must lie in  $g_i^{-1}(-\infty, a_i - \frac{1}{k}]$  for some  $k \in \mathbb{N}$ . Let  $\eta \in \mathbb{R}^m$  be the respective (by Riesz' theorem) vector to the functional  $g_i$  (for which  $\langle \eta, \cdot \rangle = g_i(\cdot)$ ). Then

$$\begin{aligned} \forall x \in \mathcal{C} : \langle \eta, x \rangle &= g_i(x) \geq a_i \\ \forall t \in E' : \langle \eta, u(t) \rangle &= g_i(u(t)) \leq a_i - \frac{1}{k}. \end{aligned}$$

□

**Definition.** We denote the sets of the *allowed* and the *extremal* controls, respectively:

$$\begin{aligned} \Psi &:= L_\infty(I, \mathcal{C}) \subset L_\infty(I, \mathbb{R}^m) \\ \Psi_e &:= L_\infty(I, \text{Ext } \mathcal{C}) \subseteq \Psi. \end{aligned}$$

*Remark.* From here on we'll talk about the space  $L_\infty(I, \mathbb{R}^m)$ , which, taken as the dual of  $L_1(I, \mathbb{R}^m)$  by the action

$$f(\varphi) := \int \langle \varphi(t), f(t) \rangle dt, \quad \text{for all } \varphi \in L_1(I, \mathbb{R}^m), f \in L_\infty(I, \mathbb{R}^m),$$

is equipped with the  $w^*$  topology.

**Lemma 2.**  $\Psi := L_\infty(I, \mathcal{C})$  is a convex  $w^*$ -compact. It is essential that here  $\mathcal{C}$  is the simplex defined earlier.

*Proof.* Obviously  $\Psi$  is convex, since for  $u_1, u_2 \in \Psi$ ,  $\lambda u_1 + (1 - \lambda)u_2$  is again a  $L_\infty$  function, and by that again  $\mathcal{C}$ -valued due to convexity of  $\mathcal{C}$ . Moreover,  $\Psi$  is bounded (in norm) with

$$\|u\|_\infty = \sup_{t \in I} \|u(t)\| \leq \max_{y \in \mathcal{C}} \|y\| \quad \text{for all } u \in \Psi.$$

For  $\Psi$  to be a  $w^*$ -compact it remains only to check it is  $w^*$ -closed, i.e. <sup>1</sup>, any  $u^0 \in L_\infty(I, \mathbb{R}^m) \setminus \Psi$  is  $w^*$ -separable from  $\Psi$ . In order to  $w^*$ -separate  $u^0$  from  $\Psi$ , we need a suitable functional  $\varphi \in L_1 \hookrightarrow (L_\infty)^*$ , with which

$$\varphi(u^0) \leq B - \varepsilon \quad \text{and} \quad \forall u \in \Psi : \varphi(u) \geq B.$$

---

<sup>1</sup>В учебника на Hermes допускат, че има  $u^0 \in \partial \Psi \setminus \Psi$ , и стигат до противоречие със същия аргумент нататък. Този стил ми се вижда излишно утежняващ, затова подходах с отвореност на допълнението.

Since  $u^0 \notin \Psi$ , then in a non-null  $E \subseteq I$ ,  $u^0(t) \notin \mathcal{C}$ . Then by lemma 1 there exist  $\varepsilon > 0, B > 0, \eta \in \mathbb{R}^m$  and a non-null  $E' \subset E$  with the property

$$\begin{aligned} \langle \eta, x \rangle &\geq B && \text{for all } x \in \mathcal{C} \\ \langle \eta, u^0(t) \rangle &\leq B - \varepsilon && \text{for all } t \in E' \end{aligned}.$$

We set  $\varphi(t) = \begin{cases} \frac{1}{\mu(E')} \eta, & t \in E' \\ 0, & \text{else} \end{cases}$ , and then  $\varphi \in L_1$  and fulfills the required conditions:

$$\begin{aligned} \varphi(u^0) &= \int_I \langle \varphi(t), u^0(t) \rangle dt = \frac{1}{\mu(E')} \int_{E'} \langle \eta, u^0(t) \rangle dt \\ &\leq \frac{1}{\mu(E')} \int_{E'} (B - \varepsilon) = B - \varepsilon \end{aligned}$$

and analogously  $\varphi(u) = \int_I \langle \varphi, u \rangle = \frac{1}{\mu(E')} \int_{E'} \langle \eta, u \rangle \geq B$  for all  $u \in \Psi$ .

Since  $\Psi$  is  $w^*$ -separated by any  $u^0 \notin \Psi$ , then  $\Psi$  is  $w^*$ -closed, and since it is  $\|\cdot\|_\infty$ -bounded – it is a  $w^*$ -compact.  $\square$

*Remark.* For  $u \in L_\infty(I, \mathbb{R}^m)$ , the product  $Y \cdot u$  is a function from  $L_1(I, \mathbb{R}^m)$ , because  $Y_{ij} \in L_1$  and

$$(Y \cdot u)_i = \sum_{j=1}^m Y_{ij}(s) u_j(s) \leq \sum_{j=1}^m Y_{ij}(s) \|u\|_\infty.$$

Then it makes sense to talk about the integral  $\int_I Y \cdot u$  that we will look at from here on.

**Definition.** We say that the control  $u \in \Psi$  leads to the point  $\xi = Tu$ , where  $T : L_\infty(I, \mathbb{R}^m) \rightarrow \mathbb{R}^n$  is the linear operator

$$Tu := \int_I Y(s) \cdot u(s) ds \quad \text{for all } u \in L_\infty(I, \mathbb{R}^m).$$

*Remark.*  $T$  is continuous (in norm), since for  $\varepsilon > 0$ , with  $\delta := \frac{\varepsilon}{\mu(I) \int_I Y}$  and  $\|h\|_\infty < \delta$  we have

$$|T(u+h) - Tu| = \left| \int_I Y(s) \cdot h(s) ds \right| \leq \mu(I) \|h\|_\infty \left| \int_I Y \right| < \varepsilon.$$

Moreover, it is  $w^*$ -continuous, since its coordinate functionals  $T_i : L_\infty(I, \mathbb{R}^m) \rightarrow \mathbb{R}$ ,  $i = 1, \dots, n$  are defined by  $T_i(u) = \sum_j \int_I Y_{ij} u_j$  where  $Y_{ij}$  are  $L_1$ -functions.

**Theorem** (*Bang-bang principle for controls in a simplex*).

$$T(\Psi) = T(\Psi_e).$$

Moreover, the two sets are convex compacts.

*Proof.*  $T\Psi_e \subseteq T\Psi$  is obvious, we need to prove the reverse inclusion. The proof goes in the following steps:

1. According to the last remark,  $T$  is  $w^*$ -continuous, hence the image  $T\Psi \subset \mathbb{R}^n$  of the  $w^*$ -compact  $\Psi$  is again a convex compact (the second statement in this theorem).
2.  $T$  is a continuous map between compacts, then the possible controls  $T^{-1}(\xi) \cap \Psi$  leading to any point  $\xi \in \mathbb{R}^n$  form a convex  $w^*$ -compact, necessarily having an extreme point  $u \in T^{-1}(\xi) \subset \Psi$  (Krein-Milman).
3. It turns out that this extremum  $u$  is a *bang-bang* control.

If we assume the opposite, then in some non-null set  $E \subset I$  the controls are far from  $\text{Ext } \mathcal{C}$  and from the lemma below (3) it follows  $u$  is a midpoint of a proper line segment of controls  $[u - h, u + h] \subset \Psi$ , leading to the same  $\xi$ , contradicting extremality of  $u$ .

□

**Lemma 3.** *Let  $n \in \mathbb{N}$ ,  $Y \in L_1(I, M_{n \times m})$ ,  $E \subset I$  is non-null, and the control  $u \in \Psi$  leads to  $\xi := Tu$  and satisfies  $\text{dist}(u(E), \text{Ext } \mathcal{C}) > \varepsilon$  for some  $\varepsilon > 0$ .*

*Then there exists a displacement  $h \in L_\infty(I, \mathbb{R}^m)$  with the following properties:*

1.  $u(t) \pm h(t) \in \mathcal{C}$  for all  $t \in I$ , where  $h(t) \neq 0$  for all  $t \in E$  ( $u$  lies in a proper line segment);
2.  $\int_I Y \cdot h = 0$  (that is,  $u \pm h$  leads again to  $\xi$ ).

In other words,  $u \in [u - h, u + h] \subset T^{-1}(\xi) \cap \Psi$ .

*Proof.* The proof goes in the following steps

1. For each  $t \in E$ , according to the next lemma (4), we can embed  $u(t)$  in a segment with endpoints  $u(t) \pm \tilde{h}(t) \in \mathcal{C}$  so that  $\|\tilde{h}(t)\|_\infty > \varepsilon^2$ . Then  $\|\tilde{h}\|_\infty > \varepsilon^2$  too. Unfortunately, however,  $\int_I Y \cdot \tilde{h} \neq 0$  in the general case.
2. From the non-atomicity of the Lebesgue measure we can partition for any  $k \in \mathbb{N}^+$  (the suitable  $k$  will be determined later)  $E = \bigcup_{j=1}^k E_j$ , so that  $E_i \cap E_j = \emptyset$  for all  $i \neq j$  and  $\mu(E_j) \geq 0$  for all  $j$ .
3. Now in each  $E_j, j = 1, \dots, k$  we will weight the restriction of  $\tilde{h}$  with a suitable  $\alpha_j \in \mathbb{R}$  so that  $\int_I Y \cdot h = 0$ :

$$h(t) := \begin{cases} \alpha_j \tilde{h}(t), & t \in E_j \\ 0, & \text{otherwise} \end{cases}.$$

The condition  $\int_I Y \cdot h = 0$  is a homogenous linear system with  $n$  equations and  $k$  unknowns

$$\int_I Y \cdot h = \sum_{j=1}^k \alpha_j \int_{E_j} Y \cdot \tilde{h} = 0.$$

A non-trivial solution for the coefficients  $\alpha_j$  always exists whenever  $k \geq n + 1$  – fix one with  $|\alpha_j| < 1$  for all  $j$ .

4. Let us convince ourselves that  $h$  satisfies property 1 (from the lemma statement). Obviously  $h(t) \neq 0$  for all  $t \in E$ , and  $u(t) \pm h(t)$  lies in  $\mathcal{C}$ , since it lies (for all  $t \in E$ ) in the line segment with endpoints  $u(t) \pm \tilde{h}(t) \in \mathcal{C}$  (due to  $|\alpha_j| < 1$ ), and  $\mathcal{C}$  is convex.

Therefore  $u \pm h \in \Psi = L_\infty(I, \mathcal{C})$  and  $u \in [u - h, u + h]$  from convexity of  $\Psi$ .

□

**Lemma 4.** *Let  $\xi \in \mathcal{C}$  satisfy  $\text{dist}(\xi, \text{Ext } \mathcal{C}) > \varepsilon$  for some  $\varepsilon > 0$ . Then  $\xi$  can be put in a line segment  $[\xi - h, \xi + h] \subset \mathcal{C}$  with  $\|h\|_\infty \geq \varepsilon^2$ .*

*Moreover, the mapping  $\xi \mapsto h$  is measurable.*

*(interestingly, we cannot state  $\|h\|_\infty \geq \varepsilon$ , the largest lower bound is  $\frac{\varepsilon^2}{1-\varepsilon}$ )*

*Proof.* Geometrically it is natural to choose  $h$  parallel to  $r_j - \xi$ , where  $r_j \in \text{Ext } \mathcal{C}$  is the closest to  $\xi$  corner with smallest index  $j$ . We put  $h := \beta(r_j - \xi)$ . Clearly  $\xi \pm h \in \text{affine span } \mathcal{C}$ . We will show that for sufficiently small  $\beta > 0$  even  $\xi \pm h \in \mathcal{C}$ .

$$\xi \pm h = \xi \pm \beta(r_j - \xi) = (1 \mp \beta)\xi \pm \beta r_j.$$

Since  $\xi \in \mathcal{C}$ , then  $\xi = \sum_i \beta_i r_i$  with some  $\beta_i \geq 0$  and  $\sum_i \beta_i = 1$ . Then

$$\begin{aligned} \xi \pm h &= (1 \mp \beta) \sum_i \beta_i r_i \pm \beta r_j \\ &= \underbrace{(\pm\beta + (1 \mp \beta)\beta_j)}_{\beta'_j} r_j + \sum_{i \neq j} \underbrace{(1 \mp \beta)\beta_i}_{\beta'_i} r_i \end{aligned}$$

and we have to check that the RHS is a convex combination of  $\{r_i\}_{i=1}^m$ .  $\sum_i \beta'_i = 1$  is equivalent to  $\xi \pm h \in \text{affine span } \mathcal{C}$ , so it remains to check  $0 < \beta'_i$  for all  $i$ .

If  $|\beta| \leq 1$  then obviously  $\beta'_i \geq 0$  for all  $i \neq j$ , and

$$\beta'_j = \pm(1 - \beta_j)\beta + \beta_j$$

and hence  $\beta'_j \geq 0$  holds always if  $\beta \leq \frac{\beta_j}{1-\beta_j}$ . We choose namely the maximal  $\beta = \beta_j/(1 - \beta_j)$ .

Since  $\text{dist}(\xi, \text{Ext } \mathcal{C}) > \varepsilon$  and  $r_j$  is a closest corner to  $\xi$ , then  $\beta_j > \varepsilon$ ,  $\beta > \frac{\varepsilon}{1-\varepsilon}$  and

$$\|h\|_\infty = \beta \|r_j - \xi\|_\infty > \beta\varepsilon = \frac{\varepsilon^2}{1-\varepsilon} > \varepsilon^2.$$

For measurability of the mapping  $\xi \mapsto h$  it suffices to observe that  $\xi \mapsto j$  is constant on sets

$$F_j = \left\{ \xi \in \mathcal{C} : \begin{array}{l} d(\xi, r_j) = \text{dist}(\xi, \text{Ext } \mathcal{C}) \\ \text{and } d(\xi, r_j) > d(\xi, r_i) \forall i < j \end{array} \right\}, \quad j = 1, \dots, m,$$

which partition  $\mathcal{C}$  and are measurable due to continuity of  $d(\cdot, \cdot)$ . □

#### Sources:

- Functional analysis and time optimal control, Volume 56 (Mathematics in Science and Engineering) by Hermes (Editor)
- Geometric Functional Analysis and its Applications Richard B. Holmes