

Name: _____ Date: _____

Period: _____

Keep in Mind

This is a practice test. It mirrors the real Unit 2 test in length, format, and the ideas it covers, but the numbers and contexts differ. Work every problem with no notes, then check your answers against the key.

Part A — Vocabulary (8 pts)

Write the letter of the best description next to each term.

- | | |
|-----------------------------------|---|
| 1. Pivot _____ | (A) $A = LU$: a lower-triangular L times an upper-triangular U . |
| 2. Multiplier _____ | (B) The first nonzero entry in a row, used to clear the entries below it. |
| 3. Upper triangular (U) _____ | (C) A matrix with no inverse — its determinant $ad - bc$ is 0. |
| 4. Inverse matrix _____ | (D) $\ell = (\text{entry to clear}) \div (\text{pivot})$; how many pivot rows you subtract. |
| 5. Singular matrix _____ | (E) The identity with two rows swapped; it records a row exchange. |
| 6. LU factorization _____ | (F) The matrix A^T formed by flipping A across its diagonal (rows $\leftrightarrow$ columns). |
| 7. Permutation matrix _____ | (G) A matrix with all zeros below the main diagonal. |
| 8. Transpose _____ | (H) A^{-1} , the matrix that undoes A : $A^{-1}A = I$. |

Part B — Multiple Choice (12 pts)

Circle the best answer.

- To clear the entry below a pivot p , you subtract ℓ times the pivot row, where ℓ is _____ (A) $p \times (\text{entry})$ (B) $(\text{entry}) \div p$ (C) $p \div (\text{entry})$ (D) $p - (\text{entry})$
- In the factorization $A = LU$, the matrix L holds _____ (A) the pivots (B) the solution $\mathbf{x}$ (C) the multipliers, with 1's on the diagonal (D) all zeros
- A 2×2 matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is **singular** (no inverse) exactly when (A) $ad - bc = 0$ (B) $ad - bc = 1$ (C) $a = d$ (D) it is upper triangular
- If P is a permutation matrix, the product PA (A) scales the rows of A (B) swaps rows of A (C) transposes A (D) inverts A
- For any two matrices that can be multiplied, $(AB)^T =$ _____ (A) $A^T B^T$ (B) AB (C) BA (D) $B^T A^T$
- Two equations in two unknowns have **infinitely many** solutions when the two lines (A) cross once (B) are parallel but different (C) are the same line (D) are perpendicular

Part C — Short Answer & Computation (30 pts)

Show your work.

1. Solve the system by **elimination**, then back-substitute and check. State the pivot and the multiplier you used.

$$x + 2y = 4, \quad 3x + 4y = 10.$$

2. Row-reduce $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 5 & 3 \\ 4 & 10 & 8 \end{bmatrix}$ to upper-triangular U . List the three multipliers and the pivots.

3. Let $A = \begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix}$. Use the $ad - bc$ formula to find A^{-1} , then check that $A^{-1}A = I$.

4. Using the inverse from the previous problem, solve $A\mathbf{x} = \mathbf{b}$ for $\mathbf{b} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$ (compute $\mathbf{x} = A^{-1}\mathbf{b}$).

5. Factor $A = \begin{bmatrix} 2 & 4 \\ 6 & 7 \end{bmatrix}$ as $A = LU$. Write L and U , then verify $LU = A$.

6. Using the L and U from the previous problem, solve $A\mathbf{x} = \mathbf{b}$ for $\mathbf{b} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ in **two sweeps**: first $L\mathbf{c} = \mathbf{b}$ (forward), then $U\mathbf{x} = \mathbf{c}$ (back).

7. **(a)** The matrix $A = \begin{bmatrix} 0 & 2 \\ 1 & 3 \end{bmatrix}$ has a 0 in the top-left pivot spot. Write the permutation matrix P that swaps the two rows, and give PA . **(b)** Write the transpose of $B = \begin{bmatrix} 1 & 2 \\ 2 & 5 \end{bmatrix}$ and state whether B is symmetric.

Part D — Extended Response (10 pts)

Explain your reasoning in full sentences.

1. Let $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$. **(a)** Compute $ad - bc$. Does A have an inverse? What does that say about solving $A\mathbf{x} = \mathbf{b}$? **(b)** How many solutions does $A\mathbf{x} = \begin{bmatrix} 3 \\ 6 \end{bmatrix}$ have? **(c)** How many does $A\mathbf{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ have? Justify (b) and (c) using elimination or the row/line picture.

2. Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 1 \end{bmatrix}$. **(a)** Compute $A^T A$ and check that it is symmetric. **(b)** Explain why $A^T A$ is *always* symmetric for any matrix A , using the reversal rule $(AB)^T = B^T A^T$.