

Kvanttilaskenta, kevät 2015 – Viikko 4

Rodion “rodde” Efremov

February 2, 2015

edx Problem 1

False. By definition of reversibility we should have x at the output of R_f .

edx Problem 2

False. The quantum circuit should modify each α_x .

edx Problem 3

True. Straight from the slides.

edx Problem 4

$$\begin{aligned} H^{\otimes 2} \frac{1}{\sqrt{2}} |00\rangle &= \frac{1}{\sqrt{2}} |++\rangle \\ &= \frac{1}{\sqrt{2}} \left(\frac{1}{\sqrt{2}} |0\rangle + \frac{1}{\sqrt{2}} |1\rangle \right) \left(\frac{1}{\sqrt{2}} |0\rangle + \frac{1}{\sqrt{2}} |1\rangle \right) \\ &= \frac{1}{2\sqrt{2}} \left(|00\rangle + |01\rangle + |10\rangle + |11\rangle \right). \end{aligned}$$

Also

$$\begin{aligned} H^{\otimes 2} \frac{1}{\sqrt{2}} |11\rangle &= \frac{1}{\sqrt{2}} |--\rangle \\ &= \frac{1}{\sqrt{2}} \left(\frac{1}{\sqrt{2}} |0\rangle - \frac{1}{\sqrt{2}} |1\rangle \right) \left(\frac{1}{\sqrt{2}} |0\rangle - \frac{1}{\sqrt{2}} |1\rangle \right) \\ &= \frac{1}{2\sqrt{2}} \left(|00\rangle - |01\rangle - |10\rangle + |11\rangle \right), \end{aligned}$$

so

$$\begin{aligned}
H^{\otimes 2}\psi &= H^{\otimes 2}\left(\frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle\right) \\
&= \frac{1}{2\sqrt{2}}\left(2|00\rangle + 2|11\rangle\right) \\
&= \frac{1}{\sqrt{2}}\left(|00\rangle + |11\rangle\right) \\
&= \psi.
\end{aligned}$$

edx Problem 5

Let

$$|\psi\rangle = \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle).$$

$$\begin{aligned}
H^{\otimes 2}\frac{1}{\sqrt{2}}|01\rangle &= \frac{1}{\sqrt{2}}|+-\rangle \\
&= \frac{1}{\sqrt{2}}\left(\frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}|1\rangle\right)\left(\frac{1}{\sqrt{2}}|0\rangle - \frac{1}{\sqrt{2}}|1\rangle\right) \\
&= \frac{1}{2\sqrt{2}}\left(|00\rangle - |01\rangle + |10\rangle - |11\rangle\right).
\end{aligned}$$

$$\begin{aligned}
H^{\otimes 2}\frac{1}{\sqrt{2}}|10\rangle &= \frac{1}{\sqrt{2}}|-+\rangle \\
&= \frac{1}{\sqrt{2}}\left(\frac{1}{\sqrt{2}}|0\rangle - \frac{1}{\sqrt{2}}|1\rangle\right)\left(\frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}|1\rangle\right) \\
&= \frac{1}{2\sqrt{2}}\left(|00\rangle + |01\rangle - |10\rangle - |11\rangle\right).
\end{aligned}$$

Now we see that $H^{\otimes 2}|\psi\rangle$ is

$$\begin{aligned}
\frac{1}{2\sqrt{2}}\left(|00\rangle - |01\rangle + |10\rangle - |11\rangle\right) + \frac{1}{2\sqrt{2}}\left(|00\rangle + |01\rangle - |10\rangle - |11\rangle\right) &= \frac{1}{2\sqrt{2}}(2|00\rangle - 2|11\rangle) \\
&= \frac{1}{\sqrt{2}}(|00\rangle - |11\rangle).
\end{aligned}$$

edx Problem 6

Yes. Both may yield $|00\rangle$ with probability $\frac{1}{2}$ and $|11\rangle$ with probability $\frac{1}{2}$.

edx Problem 7

Apply circuit A and then D.

edx Problem 8

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

edx Problem 9

Suppose

$$H^{\otimes 3} |\psi\rangle = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle).$$

Since $H^{\otimes 3}$ is reversible, if we apply it again to $H^{\otimes 3} |\psi\rangle$, we will obtain $|\psi\rangle$. Let us calculate that ket by ket:

$$\begin{aligned} H^{\otimes 3} \frac{1}{\sqrt{2}} |000\rangle &= \frac{1}{\sqrt{2}} |+++ \rangle \\ &= \frac{1}{4} \left(|0\rangle + |1\rangle \right)^3 \\ &= \frac{1}{4} \left(|0\rangle + |1\rangle \right) \left(|00\rangle + |01\rangle + |10\rangle + |11\rangle \right) \\ &= \frac{1}{4} \left(|000\rangle + |001\rangle + |010\rangle + |011\rangle + |100\rangle + |101\rangle + |110\rangle + |111\rangle \right). \end{aligned}$$

$$\begin{aligned} H^{\otimes 3} \frac{1}{\sqrt{2}} |111\rangle &= \frac{1}{\sqrt{2}} |-- - \rangle \\ &= \frac{1}{4} \left(|0\rangle - |1\rangle \right)^3 \\ &= \frac{1}{4} \left(|0\rangle - |1\rangle \right) \left(|00\rangle - |01\rangle - |10\rangle + |11\rangle \right) \\ &= \frac{1}{4} \left(|000\rangle - |001\rangle - |010\rangle + |011\rangle - |100\rangle + |101\rangle + |110\rangle - |111\rangle \right). \end{aligned}$$

Now

$$\begin{aligned}
 |\psi\rangle &= H^{\otimes 3} H^{\otimes 3} |\psi\rangle \\
 &= H^{\otimes 3} \frac{1}{\sqrt{2}} |000\rangle + H^{\otimes 3} \frac{1}{\sqrt{2}} |11\rangle \\
 &= \frac{1}{4} (2 |000\rangle + 2 |011\rangle + 2 |101\rangle + 2 |110\rangle) \\
 &= \frac{1}{2} (|000\rangle + |011\rangle + |101\rangle + |110\rangle).
 \end{aligned}$$

edx Problem 10

(a)

$$\frac{1}{2^{n-1}}.$$

(b) We see a uniformly random string $y \in \{0, 1\}^n$.

edx Problem 11

$$\frac{1}{\sqrt{2^n}} \sum_{x \in \{0,1\}^n} (-1)^{x \cdot f(x)} |x\rangle |f(x)\rangle.$$

edx Problem 12

$$\frac{1}{\sqrt{2^n}} \sum_x |x\rangle |0101\rangle.$$

edx Problem 13

1111.

edx Problem 14

Suppose Alice starts with two qubits in the Bell state $\frac{1}{\sqrt{2}} |00\rangle + \frac{1}{\sqrt{2}} |11\rangle$ and teleports these qubits to Bob by applying the quantum teleportation protocol to each qubit separately.

As we are speaking about teleportation, Bob sees the same state and receives exactly 2 bits of information as there is only 4 Bell states.

QCE 7.1

$$\vec{\sigma} = \sigma_x \hat{x} + \sigma_y \hat{y} + \sigma_z \hat{z}, \quad \vec{n} = \sin \theta \cos \phi \hat{x} + \sin \theta \sin \phi \hat{y} + \cos \theta \hat{z}.$$

Now

$$\begin{aligned} \vec{\sigma} \cdot \vec{n} &= (\sigma_x \hat{x} + \sigma_y \hat{y} + \sigma_z \hat{z})(\sin \theta \cos \phi \hat{x} + \sin \theta \sin \phi \hat{y} + \cos \theta \hat{z}) \\ &= \end{aligned}$$

QCE 7.2

Let

$$|\psi\rangle = \frac{|0\rangle|1\rangle - |1\rangle|0\rangle}{\sqrt{2}}.$$

Now we have

$$|0\rangle = \frac{|+\rangle + i|-\rangle}{\sqrt{2}}, \quad |1\rangle = \frac{|+\rangle - i|-\rangle}{\sqrt{2}},$$

so

$$\begin{aligned} |\psi\rangle &= \frac{|0\rangle|1\rangle - |1\rangle|0\rangle}{\sqrt{2}} \\ &= \frac{1}{2\sqrt{2}} \left((|+\rangle + i|-\rangle)(|+\rangle - i|-\rangle) - (|+\rangle - i|-\rangle)(|+\rangle + i|-\rangle) \right) \\ &= \frac{1}{2\sqrt{2}} \left(|++\rangle - i|+-\rangle + i|-+\rangle + |--\rangle + |++\rangle + i|+-\rangle - i|-+\rangle + |--\rangle \right) \\ &= \frac{1}{2\sqrt{2}} (2|++\rangle + 2|--\rangle) \\ &= \frac{1}{\sqrt{2}} (|++\rangle + |--\rangle). \end{aligned}$$

QCE 7.3

Let

$$|\beta_{00}\rangle = \frac{|00\rangle + |11\rangle}{\sqrt{2}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \quad |\beta_{01}\rangle = \frac{|01\rangle + |10\rangle}{\sqrt{2}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ 1 \\ 0 \end{pmatrix}.$$

Also we know that

$$Z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

so

$$\begin{aligned} Z \otimes Z &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \end{aligned}$$

Now

$$\begin{aligned} Z \otimes Z |\beta_{00}\rangle &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix} \\ &= \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix} \\ &= \beta_{00} \\ &= (-1)^y |\beta_{xy}\rangle, \end{aligned}$$

where $y = 0$ and $x = 0$. Also

$$\begin{aligned} Z \otimes Z |\beta_{01}\rangle &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ 1 \\ 0 \end{pmatrix} \\ &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ -1 \\ -1 \\ 0 \end{pmatrix} \\ &= -\beta_{01} \\ &= (-1)^y |\beta_{xy}\rangle, \end{aligned}$$

where $y = 1$ and $x = 0$.

QCE 7.4

Show that $X \otimes X |\beta_{xy}\rangle = (-1)^x |\beta_{xy}\rangle$. From the book we know that

$$|\beta_{xy}\rangle = \frac{|0y\rangle + (-1)^x |1\bar{y}\rangle}{\sqrt{2}} = \frac{1}{\sqrt{2}} \begin{pmatrix} \bar{y} \\ y \\ (-1)^x y \\ (-1)^x \bar{y} \end{pmatrix}.$$

Also

$$\begin{aligned} X \otimes X &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \end{aligned}$$

so

$$\begin{aligned} X \otimes X |\beta_{xy}\rangle &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \bar{y} \\ y \\ (-1)^x y \\ (-1)^x \bar{y} \end{pmatrix} \\ &= \frac{1}{\sqrt{2}} \begin{pmatrix} (-1)^x \bar{y} \\ (-1)^x y \\ y \\ \bar{y} \end{pmatrix} \\ &= \frac{(-1)^x |0y\rangle + |1\bar{y}\rangle}{\sqrt{2}} \\ &= (-1)^x \beta_{xy}. \end{aligned}$$

QCE 7.5

Show that $Y \otimes Y |\beta_{xy}\rangle = (-1)^{x+y} |\beta_{xy}\rangle$. From the book we know that

$$|\beta_{xy}\rangle = \frac{|0y\rangle + (-1)^x |1\bar{y}\rangle}{\sqrt{2}} = \frac{1}{\sqrt{2}} \begin{pmatrix} \bar{y} \\ y \\ (-1)^x y \\ (-1)^x \bar{y} \end{pmatrix}.$$

Also

$$\begin{aligned} Y \otimes Y &= \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \\ &= \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}, \end{aligned}$$

so

$$\begin{aligned}
Y \otimes Y |\beta_{xy}\rangle &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \bar{y} \\ y \\ (-1)^x y \\ (-1)^x \bar{y} \end{pmatrix} \\
&= \frac{1}{\sqrt{2}} \begin{pmatrix} -(-1)^x \bar{y} \\ (-1)^x y \\ y \\ -\bar{y} \end{pmatrix} \\
&= (-1)^{x+\bar{y}} |\beta_{xy}\rangle.
\end{aligned}$$

The book has a typo...

QCE 7.6

Show that $X \otimes X$ commutes with $Z \otimes Z$. They commute if and only if $X \otimes XZ \otimes Z = X \otimes XZ \otimes Z$:

$$\begin{aligned}
X \otimes X &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}^{\otimes 2} \\
&= \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}.
\end{aligned}$$

$$\begin{aligned}
Z \otimes Z &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\otimes 2} \\
&= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.
\end{aligned}$$

Now

$$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

Also

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix},$$

which proves that the two matrices commute.

QCE 7.7

Consider the eigenvectors in Example 7.4. Show that $[H_I, \vec{\sigma}_A \cdot \vec{\sigma}_B] = 0$, and hence show that the eigenvectors of the Hamiltonian are eigenvectors of the $\vec{\sigma}_A \cdot \vec{\sigma}_B$ operator. In particular, show that $\vec{\sigma}_A \cdot \vec{\sigma}_B |\phi_i\rangle = |\phi_i\rangle$ for $i = 1, 2, 3$ and $\vec{\sigma}_A \cdot \vec{\sigma}_B |\phi_4\rangle = -3 |\phi_4\rangle$.

From the Example 7.4 we have that

$$H_I = \frac{\mu^2}{r^3} \begin{pmatrix} -2 & 0 & 0 & 0 \\ 0 & 2 & 2 & 0 \\ 0 & 2 & 2 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix}, \quad \vec{\sigma}_A \cdot \vec{\sigma}_B = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 2 & 0 \\ 0 & 2 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Condition $[H_I, \vec{\sigma}_A \cdot \vec{\sigma}_B] = 0$ means that the two matrices commute and that is the case as a routine calculation may show. Wolframalpha tells me that the eigenvectors of H_I are

$$\begin{pmatrix} 0 \\ -1 \\ -1 \\ 0 \end{pmatrix}, \quad \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \quad \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad \begin{pmatrix} 0 \\ -1 \\ 1 \\ 0 \end{pmatrix},$$

and that is not in accord with the eigenvectors of $\vec{\sigma}_A \cdot \vec{\sigma}_B$, thank you, book. :(

Now

$$\begin{aligned} \vec{\sigma}_A \cdot \vec{\sigma}_B |\phi_1\rangle &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 2 & 0 \\ 0 & 2 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ 1 \\ 0 \end{pmatrix} \\ &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ 1 \\ 0 \end{pmatrix} \\ &= |\phi_1\rangle, \end{aligned}$$

$$\begin{aligned} \vec{\sigma}_A \cdot \vec{\sigma}_B |\phi_2\rangle &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 2 & 0 \\ 0 & 2 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} \\ &= \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} \\ &= |\phi_2\rangle, \end{aligned}$$

$$\begin{aligned}
\vec{\sigma}_A \cdot \vec{\sigma}_B |\phi_3\rangle &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 2 & 0 \\ 0 & 2 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \\
&= \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \\
&= |\phi_3\rangle,
\end{aligned}$$

$$\begin{aligned}
\vec{\sigma}_A \cdot \vec{\sigma}_B |\phi_4\rangle &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 2 & 0 \\ 0 & 2 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ -1 \\ 0 \end{pmatrix} \\
&= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ -3 \\ 3 \\ 0 \end{pmatrix} \\
&= -3 |\phi_4\rangle.
\end{aligned}$$

QCE 7,8

Is the state $X \otimes Z |\beta_{00}\rangle$ entangled?

$$\begin{aligned}
X \otimes Z |00\rangle &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix} \\
&= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix} \\
&= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ -1 \\ 1 \\ 0 \end{pmatrix} \\
&= \frac{-|01\rangle + |10\rangle}{\sqrt{2}} \\
&= -|\beta_{11}\rangle,
\end{aligned}$$

so the state is entangled but is not a Bell state.

QCE 7.9

Find the Pauli representation of

$$\rho = \begin{pmatrix} \sin^2 \theta & e^{-i\phi} \sin \theta \cos \theta \\ e^{i\phi} \sin \theta \cos \theta & \cos^2 \theta \end{pmatrix}.$$

First we need to compute

$$\begin{aligned} c_0 &= \text{Tr}(\rho\sigma_0) \\ &= \text{Tr}(\rho) \\ &= \sin^2 \theta + \cos^2 \theta \\ &= 1, \end{aligned}$$

$$\begin{aligned} c_1 &= \text{Tr}(\rho\sigma_1) \\ &= \text{Tr}(\rho\sigma_x) \\ &= \text{Tr} \begin{pmatrix} \sin^2 \theta & e^{-i\phi} \sin \theta \cos \theta \\ e^{i\phi} \sin \theta \cos \theta & \cos^2 \theta \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \\ &= \text{Tr} \begin{pmatrix} e^{-i\phi} \sin \theta \cos \theta & \sin^2 \theta \\ \cos^2 \theta & e^{i\phi} \sin \theta \cos \theta \end{pmatrix} \\ &= e^{-i\phi} \sin \theta \cos \theta + e^{i\phi} \sin \theta \cos \theta \\ &= (\sin \theta \cos \theta)(\cos \phi - i \sin \phi + \cos \phi + i \sin \phi) \\ &= 2 \sin \theta \cos \theta \cos \phi, \end{aligned}$$

$$\begin{aligned} c_2 &= \text{Tr}(\rho\sigma_2) \\ &= \text{Tr}(\rho\sigma_y) \\ &= \text{Tr} \begin{pmatrix} \sin^2 \theta & e^{-i\phi} \sin \theta \cos \theta \\ e^{i\phi} \sin \theta \cos \theta & \cos^2 \theta \end{pmatrix} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \\ &= \text{Tr} \begin{pmatrix} ie^{-i\phi} \sin \theta \cos \theta & -i \sin^2 \theta \\ i \cos^2 \theta & -ie^{i\phi} \sin \theta \cos \theta \end{pmatrix} \\ &= ie^{-i\phi} \sin \theta \cos \theta - ie^{i\phi} \sin \theta \cos \theta \\ &= \sin \theta \cos \theta (ie^{-i\phi} - ie^{i\phi}) \\ &= \sin \theta \cos \theta (i(\cos \phi - i \sin \phi) - i(\cos \phi + i \sin \phi)) \\ &= \sin \theta \cos \theta (i \cos \phi - i^2 \sin \phi - i \cos \phi - i^2 \sin \phi) \\ &= 2 \sin \theta \cos \theta \sin \phi. \end{aligned}$$

$$\begin{aligned}
c_3 &= \text{Tr}(\rho\sigma_3) \\
&= \text{Tr}(\rho\sigma_z) \\
&= \text{Tr} \begin{pmatrix} \sin^2 \theta & e^{-i\phi} \sin \theta \cos \theta \\ e^{i\phi} \sin \theta \cos \theta & \cos^2 \theta \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \\
&= \text{Tr} \begin{pmatrix} \sin^2 \theta & -e^{-i\phi} \sin \theta \cos \theta \\ e^{i\phi} \sin \theta \cos \theta & -\cos^2 \theta \end{pmatrix} \\
&= \sin^2 \theta - \cos^2 \theta.
\end{aligned}$$

The Pauli representation for the system is

$$\rho = \sum_{i=0}^3 \text{Tr}(\sigma_i)\sigma_i.$$

QCE 7.10

Use (7.36) to show that $|\beta_{10}\rangle$ is entangled. Apply the same criterion to test $X \otimes Z |\beta_{00}\rangle$.

Here we have

$$|\beta_{00}\rangle = \frac{|00\rangle + |11\rangle}{\sqrt{2}}, \quad |\beta_{10}\rangle = \frac{|00\rangle - |11\rangle}{\sqrt{2}}.$$

The density operator for $|\beta_{10}\rangle$ is

$$\rho = \frac{1}{2}(|00\rangle - |11\rangle)(\langle 00| - \langle 11|) = \frac{1}{2}(|00\rangle\langle 00| - |00\rangle\langle 11| - |11\rangle\langle 00| + |11\rangle\langle 11|),$$

and its density matrix is

$$\frac{1}{2} \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix}.$$

Next we find

$$\begin{aligned}
c_{11} &= \text{Tr}(\rho X \otimes X) = \text{Tr} \frac{1}{2} \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \\
&= \frac{1}{2} \text{Tr} \begin{pmatrix} -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & -1 \end{pmatrix} \\
&= -1
\end{aligned}$$

$$\begin{aligned}
c_{22} &= \text{Tr}(\rho Y \otimes Y) = \text{Tr} \frac{1}{2} \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix} \\
&= \frac{1}{2} \text{Tr} \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix} \\
&= 1
\end{aligned}$$

$$\begin{aligned}
c_{33} &= \text{Tr}(\rho Z \otimes Z) = \text{Tr} \frac{1}{2} \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\
&= \frac{1}{2} \text{Tr} \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix} \\
&= 1.
\end{aligned}$$

Now we see that $|c_{11}| + |c_{22}| + |c_{33}| = 3$, so the state is entangled.

$$\begin{aligned}
X \otimes Z |\beta_{00}\rangle &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix} \\
&= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix} \\
&= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ -1 \\ 1 \\ 0 \end{pmatrix} \\
&= \frac{-|01\rangle + |10\rangle}{\sqrt{2}}.
\end{aligned}$$

Now the density operator/matrix for the above state is

$$\begin{aligned}
\rho &= \frac{1}{2}(-|01\rangle + |10\rangle)(-\langle 01| + \langle 10|) \\
&= \frac{1}{2}(|01\rangle\langle 01| - |01\rangle\langle 10| - |10\rangle\langle 01| + |10\rangle\langle 10|) \\
&= \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.
\end{aligned}$$

Next we find

$$\begin{aligned}
c_{11} &= \text{Tr}(\rho X \otimes X) = \frac{1}{2} \text{Tr} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \\
&= \frac{1}{2} \text{Tr} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} = -1,
\end{aligned}$$

$$\begin{aligned}
c_{22} &= \text{Tr}(\rho Y \otimes Y) = \frac{1}{2} \text{Tr} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix} \\
&= \frac{1}{2} \text{Tr} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} = -1,
\end{aligned}$$

$$\begin{aligned}
c_{33} &= \text{Tr}(\rho Z \otimes Z) = \frac{1}{2} \text{Tr} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\
&= \frac{1}{2} \text{Tr} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} = -1.
\end{aligned}$$

Once again we have $|c_{11}| + |c_{22}| + |c_{33}| = 3$ so this state is entangled as well.

QCE 7.11

Derive

$$|\beta_{00}\rangle\langle\beta_{00}| = \frac{1}{4}(I \otimes I + X \otimes X - Y \otimes Y + Z \otimes Z).$$

We have that

$$|\beta_{00}\rangle = \frac{|00\rangle + |11\rangle}{\sqrt{2}},$$

so

$$\begin{aligned} |\beta_{00}\rangle \langle \beta_{00}| &= \frac{1}{2}(|00\rangle + |11\rangle)(\langle 00| + \langle 11|) \\ &= \frac{1}{2}(|00\rangle \langle 00| + |00\rangle \langle 11| + |11\rangle \langle 00| + |11\rangle \langle 11|) \\ &= \frac{1}{2} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}. \end{aligned}$$

Now

$$\begin{aligned} I \otimes I + X \otimes X - Y \otimes Y + Z \otimes Z &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \\ &\quad - \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix} + \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & -1 & -1 & 0 \\ 0 & -1 & -1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 2 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 2 & 0 & 0 & 2 \end{pmatrix} \\ &= 4 |\beta_{00}\rangle \langle \beta_{00}|, \end{aligned}$$

which yields the desired result.

QCE 7.12

Can the following state be written in diagonal form in terms of the Bell basis?

$$\rho = \begin{pmatrix} \frac{1}{2} & 0 & 0 & -\frac{1}{8} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -\frac{1}{8} & 0 & 0 & \frac{1}{2} \end{pmatrix}$$

Using (7.43), determine if this state is a separable state.

Now

$$\begin{aligned}
\rho &= \frac{1}{2} \left(|00\rangle \langle 00| + |11\rangle \langle 11| \right) - \frac{1}{8} \left(|00\rangle \langle 11| + |11\rangle \langle 00| \right) \\
&= \frac{1}{2} \left(|\beta_{00}\rangle \langle \beta_{00}| + |\beta_{10}\rangle \langle \beta_{10}| \right) - \frac{1}{8} \left(|\beta_{00}\rangle \langle \beta_{00}| - |\beta_{10}\rangle \langle \beta_{10}| \right) \\
&= \frac{3}{8} \left(|\beta_{00}\rangle \langle \beta_{00}| \right) + \frac{5}{8} \left(|\beta_{10}\rangle \langle \beta_{10}| \right).
\end{aligned}$$

Since $c_{00} = \frac{3}{8} \leq \frac{1}{2}$, this state is separable.