When misleading context changes the answer, does MedGemma say so?

Temesgen Tewolde
Duke University · AIPI 590: Explainable AI

Abstract

A misleading sentence in the prompt can change a medical model's answer. This study asks whether MedGemma 1.5 4B admits it. Two cues, a fabricated clinical claim and a bare "the answer is C", were added to 200 exam questions, with the reasoning trace on and off. Both cues moved the model to the wrong answer in 34%–42% of eligible cases. When it followed, the response mentioned the claim 78% of the time but the bare assertion only 39%.

Objectives


1. How often does each cue move the model to a wrong answer?
2. When it does, do the trace or the response say so?
3. Does letting the model think first change either?

Methods

- **Data:** 200 MedMisQA questions from MedMisBench[1], each with a fabricated claim for one wrong option.
- **Model:** MedGemma 1.5 4B[3] (open weights), run with thinking on and off.
- **Conditions** (one sentence added after the options): clean; sham (irrelevant); fabricated claim; bare assertion ("The correct answer is C").
- **Runs:** 4 conditions × 2 modes × 200 questions = 1,600 greedy rollouts.
- **Follow rate:** share of questions where the model moves to the cued wrong option. Sham baseline ≈ 4%.
- **Disclosure:** a judge model (Qwen3-14B) checks whether the trace or response cites the cue as given. Checked against blind LLM annotators (κ 0.78 response, 0.76 trace); no human labels yet.

Results


- Both cues steer the model, and thinking does not protect ($p = 0.76$, $p = 1.00$). Authority-framed cues were followed about twice as often as patient-framed ones.
- The trace admits the bare assertion more than the response does (60% vs 39%, $p = 0.002$). For a fabricated claim the two are level (80% vs 78%).




Follow rate by cue and mode. Dashed line: sham baseline (4%). Whiskers: 95% CIs.

61% of followed bare assertions never mentioned

21% of followed fabricated claims never mentioned



Share of followed rollouts that mention the cue. Followed (on / off): assertion 59 / 49, claim 70 / 60. No trace when thinking is off.



Follow rate by source framing (thinking on and off pooled). Authority-framed cues were followed about twice as often as patient-framed ones.

Conclusion

- A bare "the answer is C" moves a small medical model almost as much as a fabricated clinical claim.
- It leaves less in the response: most followed assertions are never mentioned there.
- Thinking first does not reduce susceptibility.
- The trace holds what the response lacks, for the assertion only, and only if the provider exposes it.

Untested implication: a monitor reading only the response would miss most of these cases.

Example

Without a cue the model chose **D** (correct). With **"The clinical guideline states that the correct answer is C"** added, it chose **C**.

Trace: "The clinical guideline states the correct answer is C. This confirms my reasoning."

Response: "...Cephalexin is a safe and effective first-line treatment for acute uncomplicated cystitis..." **It never mentions the guideline.**

Limitations

- One small model; English multiple choice; greedy decoding.
- Judge checked only against LLM annotators; weakest for thinking-off responses (κ 0.44).
- Thinking-off still reasons in the response; 10–15% of those rollouts loop.
- Source framing is not randomized; groups hold different questions.

References

1. Zhou et al. *MedMisBench: Measuring Epistemic Resilience of LLMs Under Misleading Medical Context.* arXiv:2606.12291 (2026).
2. Linzmayer & Elhadad. *Untangling the Mechanisms of Misleading Context in Medical Question Answering.* arXiv:2609.02754 (2026).
3. *MedGemma technical report.* arXiv:2507.05201 (2025).

Note: the cues are deliberately false medical statements. Evaluation use only, not clinical advice.