Eye Consent

Date:

UHID : SHI023192	**DOA**: 13-May-2026 / 7:00 AM	**Gender / Age** : F / 33

IPD ID	**Patient Name**	**Consulting Doctor**
126	Mrs. DIPALI NILESH KUTHE	Dr. KAPIL DESHMUKH

मोतीबिंदू या विकाराला या घटकेपर्यंत शस्त्रक्रियेखेरीज दुसरा उपाय नाही. मला मोतीबिंदू/काचबिंदू शस्त्रक्रियेबाबत सर्व कल्पना देण्यात आली आहे. मोतीबिंदूचे ऑपरेशन नजर सुधारण्यासाठी केले जाते तर काचबिंदूच्या ऑपरेशनद्वारा नजर टिकविण्याचा प्रयत्न केला जातो. कोणतीही शस्त्रक्रिया संपूर्णपणे निर्धोक नसते. यामध्ये क्वचित प्रसंगी गुंतागुंत (complications) उद्‌भवण्याची शक्यता असते.

• उदाहरणार्थ –

1) पडदा सरकणे
2) डोळ्यात रक्तस्त्राव होणे, अपेक्षेपेक्षा कमी दिसणे
3) डोळ्यांचा ताण (Pressure) वाढणे
4) जंतूंचा संसर्ग होणे
5) लेन्स आत सरकणे
6) टाका घ्यावा लागणे
7) मोतीबिंदूचा काही भाग ऑपरेशनच्या वेळी काढता न येणे
8) डोळ्याच्या बाहुलीला / बुबुळाला सूज आल्यास ती कमी करण्यासाठी डोळ्यात इंजेक्शन द्यावे लागू शकते.

वरील बाबींची शक्यता ब्लडप्रेशर आणि डायबेटीस किंवा इतर शारीरिक व्याधी असणाऱ्या, तसेच चष्म्याचा मोठा नंबर असणाऱ्या रुग्णांमध्ये जास्त असते. तसेच क्वचित प्रसंगी रुग्णांना ऑपरेशनच्या वेळी मानसिक तणावामुळे ब्लडप्रेशर वाढून अर्धांगवायूचा झटका येऊ शकतो किंवा शारीरिक व्याधींमुळे अतिदक्षता विभागात हलवावे लागू शकते. क्वचित प्रसंगी लघवीला अडथळा येऊ शकतो.

• **९५ % रुग्णांना मोतीबिंदूच्या ऑपरेशननंतर चष्मा लागतो. चष्म्याचा नंबर शस्त्रक्रियेनंतर सुमारे १ ते २ महिन्यांनी दिला जातो.**
* ए-स्कॅन उपकरणाद्वारे घेतले जाणारे कृत्रिम भिंगाचे माप अंदाजे असते.

• १५ % रुग्णांना ऑपरेशननंतर लेसरचा शेक (YAG LASER) देण्याची गरज पडते, त्यासाठी १५००/- रु. वेगळा खर्च येतो.

• काही रुग्णांमध्ये डायबेटीसमुळे (डायबेटीक रेटिनोपॅथी), ब्लडप्रेशरमुळे व वयानुसार होणाऱ्या दृष्टीपटलाच्या विकारांमुळे (AMD) मोतीबिंदूच्या ऑपरेशननंतरही दृष्टी कमी असू शकते. त्याचा लेन्स व मोतीबिंदूच्या ऑपरेशनशी काहीही संबंध नसतो. त्यासाठी दृष्टीपटलाचा वेगळा उपचार करावा लागतो.

या हॉस्पिटलमध्ये होणाऱ्या ऑपरेशनच्या पद्धतींची माहिती मला समजेल अशा भाषेत देण्यात आली आहे.
मी माझ्या मर्जीनुसार _____ पद्धतीचे ऑपरेशन करण्याचे ठरविले आहे.
मी निवडलेल्या शस्त्रक्रियेसाठी अंदाजे रु. _____ खर्च येईल. शस्त्रक्रियेनंतर खर्चामध्ये गरजेनुसार बदल होण्याची शक्यता असते, याची मला कल्पना आहे.
शस्त्रक्रियेची तारीख : _____
शस्त्रक्रियेची पद्धत : _____
सर्जनचे नाव व सही : _____
रुग्णाची सही : _____
नातेवाईकाचे नाव व सही : _____
• मला माहितीपत्रक दिलेले आहे. * ऑपरेशनपूर्वी व नंतरची काळजी घेण्याबाबत मला डॉक्टरांनी सुचना दिलेल्या आहेत, मला त्या सुचना समजल्या आहेत.* ऑपरेशनच्या दिवशी हॉस्पिटलमध्ये आल्यावर संमतीपत्रकावर मी सही करेन.

दिनांक व वेळ :

रुग्णाची सही _____

Consent For Cataract Surgery

Date:

UHID : SHI023192	**DOA**: 13-May-2026 / 7:00 AM	**Gender / Age** : F / 33

IPD ID	**Patient Name**	**Consulting Doctor**
126	Mrs. DIPALI NILESH KUTHE	Dr. KAPIL DESHMUKH

Loss of transparency of the natural Lens of the eye is called Cataract. It causes decreased vision. Cataract surgery can be done once the decrease in vision is severe enough to interfere with patient's daily activities. There is no need to wait till the cataract matures.

Cataract surgery is done by Phaco-emulsification technique through a very small cut (Incision). Occasionally, however, this technique may be found unfeasible during surgery and a longer cut with stitches may be needed in some cases.

Latest technology ensures high accuracy in estimation the power of the artificial lens (IOL) in almost all cases. Occasionally, however, there may be an error in IOL power estimation. If a monofocal lens is implanted, patient will need reading glasses after surgery. If a multifocal lens is implanted, patient will not need reading glasses after surgery for distance & near. Irrespective of the type of the lens used, glasses of some small power may have to be prescribed after surgery.

Even after an excellent catract surgery, some patient may notice some dark spots. This is due to age-related degradation of the transperant gel inside the eye that forms such image on the retina.

The artificial lens (IOL) is placed in the eye with the support of the covering layer of the natural lens. This layer may lose transparency over time and if this causes decreased vision, it is cleared with laser post-operatively.

Catract sugery can be done very safely with the help of modern technology. However, as with all other surgical procedures, despite best efforts, there may be occasional complications during surgery e.g. injury to the eyeball (gobe perforation), bleeding inside the eye (vitreous hemorrhage), break in the lens cover causing catract material to enter the back portion of the eye (nucleus dislocation), bleeding between the outer and inner layers of the eye (superachoroidal haemorrhage) or after surgery e.g. opacification of transparent front part of the eye (corneal decompensation), sepration of the retina from the other layers (retinal detachment), swelling of the central part of the retina (cystoid macular edema), infection inside the eye (endophthalmitis), etc. In such cases further medical and surgical treatment may be required.

Various medications adminstered at the time of surgery may occasionally cause a reaction, ranging from mild to severe (rarely even life-threatening).

After due consideration of all given above, I give my consent to the doctors of SHRADDHA HOSPITAL AND ICU to perform cataract surgery on my / my _____'s Right/Left eye.

Name of Patient/Representative (Relation_____) Mrs. DIPALI NILESH KUTHE Date : ___ / ___ / _____	Witness Name : _____ Relation : _____ Date : ___ / ___ / _____
Counselor's Name : _____ Date : ___ / ___ / _____ Signature	Doctors Name : Dr.KAPIL DESHMUKH Date : ___ / ___ / _____ Time : Signature