



DPP - 1

Video Solution on Website:-

https://physicsaholics.com/home/courseDetails/57

Video Solution on YouTube:-

https://youtu.be/Ye7mcKHdp1I

- Q 1. The postulates of kinetic theory will be true if the number of molecules be -
 - (a) Any

- (b) Very large
- (c) Very small
- (d) Avogadro's number
- Q 2. Which of the following statements about kinetic theory of gases is wrong
 - (a) The molecules of a gas are in continuous random motion
 - (b) The molecules continuously undergo inelastic collisions
 - (c) The molecules do not interact with each other except during collisions
 - (d) The collisions amongst the molecules are of short duration
- Under which of the following conditions is the law RV = RT obeyed most closely by a Q 3. real gas
 - (a) High pressure and high temperature
 - (b) Low pressure and low temperature
 - (c) Low pressure and high temperature
 - (d) High pressure and low temperature
- A certain sample of gas has a volume of 0.2 litre measured at 1 atm pressure and 0 °C. Q 4. At the same pressure but at 273 °C, its volume will be
 - (a) 0.4 litres

(b) 0.8 litres

(c) 27.8 litres

(d) 55.6 litres

Figure shows two flasks connected to each other. The volume of the flask 1 is twice Q 5. that of flask 2. The system is filled with an ideal gas at temperature 100 K and 200 K respectively. If the mass of the gas in 1 be m then what is the mass of the gas in flask 2



- (a) m
- (c) m/4
- (d)m/8
- Q 6. If the pressure of an ideal gas contained in a closed vessel is increased by 0.5%, the increase in temperature is 2 K. The initial temperature of the gas is
 - (a) 27° C
- (b) 127° C
- (c) 300°*C*
- (d) 400° C



Physicsaholics





- Q 7. Air is pumped into an automobile tube upto a pressure of $200 \, kPa$ in the morning when the air temperature is $22^{\circ}C$. During the day, temperature rises to $42^{\circ}C$ and the tube volume expands by 2%. The pressure of the air in the tube at this temperature, will be approximately
 - (a) 212 *kPa*
- (b) 206 kPa
- (c) 209 *kPa*
- (d) 200 kPa
- Q 8. A vessel is filled with an ideal gas at a pressure of 10 atmospheres and temperature 27 °C. Half of the mass of the gas is removed from the vessel and temperature of the remaining gas is increased to 87°C. Then the pressure of the gas in the vessel will be
 - (a) 5 atm

(b) 6 atm

(c) 7 atm

- (d) 8 atm
- Q 9. The pressure P, volume V and temperature T of a gas in the jar A and the other gas in the jar B at pressure P, volume P04 and temperature P27, then the ratio of the number of molecules in the jar P4 and P6 will be-
 - (a) 1:1
- (b) 2:1
- (c) 1:2
- (d) 4:1
- Q 10. Which one of the following graphs represents the behavior of an ideal gas when temperature is constant?





- Q 11. During an experiment an ideal gas is found to obey an additional law $VP^2 = constant$. The gas is initially at temperature T and volume V, when it expands to volume 2V, the resulting temperature is -
 - (a) T/2

(b) $T/\sqrt{2}$


(c) $T\sqrt{2}$

- (d) 2T
- Q 12. A vessel has 6 g of hydrogen at pressure P and temperature 500 K. A small hole is made in it so that hydrogen leaks out. How much hydrogen leaks out if the final pressure is P/2 and temperature falls to 300 K -
 - (a) 2 g
- (b) 4 g
- (c) 3 g
- (d) 1g
- Q 13. The figure shows pressure versus density graph for an ideal gas at two temperature T_1 and T_2 :



hysicsaholics






- (a) $T_1 > T_2$ (c) $T_1 = T_2$

- (b) $T_1 < T_2$ (d) None of these
- Q 14. A vessel is filled with a gas at a pressure of 76 cm of mercury at a certain temperature. The mass of the gas is increased by 50 % by introducing more gas in the vessel at the same temperature. The resultant pressure, in cm of Hg, is -
 - (a) 76

(b) 114

(c) 152

- (d) 1117
- Q 15. Pressure versus temperature graph of an ideal gas is as shown in figure. Density of the gas at point A is ρ_0 . Density at B will be



- (d) $2\rho_0$

Q.1 b	Q.2 b	Q.3 c	Q.4 a	Q.5 c
Q.6 b	Q.7 с	Q.8 b	Q.9 d	Q.10 d
Q.11 c	Q.12 d	Q.13 a	Q.14 b	Q.15 b