



# SIR PRATEEK JAIN

- . Founder @Physicsaholics
- . Top Physics Faculty on Unacademy (IIT JEE & NEET)
- . 8+ years of teaching experience in top institutes like FIITJEE (Delhi, Indore) , CP (KOTA) etc.
- . Produced multiple Top ranks.
- . Research work with HC Verma sir at IIT Kanpur
- . Interviewed by International media.

PLUS

ICONIC <sup>\*\*</sup>

✓ India's Best Educators

✓ Interactive Live Classes

✓ Structured Courses & PDFs

✓ Live Tests & Quizzes

✗ Personal Coach

✗ Study Planner

24 months

No cost EMI

₹2,333/mo

₹56,000

>

18 months

No cost EMI

₹2,625/mo

₹47,250

>

12 months

No cost EMI

₹3,208/mo

₹38,500

>

6 months

No cost EMI

₹4,667/mo

₹28,000

>

To be paid as a one-time payment

View all plans

 Add a referral code

APPLY

# PHYSICSLIVE

PLUS

ICONIC <sup>\*\*</sup>

✓ India's Best Educators

✓ Interactive Live Classes

✓ Structured Courses & PDFs

✓ Live Tests & Quizzes

✗ Personal Coach

✗ Study Planner

24 months

No cost EMI

₹2,100/mo

+10% OFF ₹50,400

>

18 months

No cost EMI

₹2,363/mo

+10% OFF ₹42,525

>

12 months

No cost EMI

₹2,888/mo

+10% OFF ₹34,650

>

6 months

No cost EMI


₹4,200/mo

+10% OFF ₹25,200

>

To be paid as a one-time payment

View all plans

 Awesome! PHYSICSLIVE code applied

×

Use code **PHYSICSLIVE** to get 10% OFF on Unacademy PLUS.

For Video Solution of this DPP, Click on below link

Video Solution  
on Website:-

<https://physicsaholics.com/home/courseDetails/57>

Video Solution  
on YouTube:-

<https://youtu.be/Ye7mcKHdp1I>

# Physics DPP

**DPP- 1 kinetic theory of gases**  
**By Physicsaholics Team**

Q) The postulates of kinetic theory will be true if the number of molecules be -

(a) Any

(b) Very large

(c) Very small

(d) Avogadro's number

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. b



Solution:

According to Kinetic theory all directions are equally favourable for motion of molecules. This assumption is true only if there are large no of molecules

ANS(b)

Q) Which of the following statements about kinetic theory of gases is wrong

- (a) The molecules of a gas are in continuous random motion
- (b) The molecules continuously undergo inelastic collisions
- (c) The molecules do not interact with each other except during collisions
- (d) The collisions amongst the molecules are of short duration

**Join Unacademy PLUS Referral Code :**

**Physicslive**



Ans. b

Solution:

In KTG the assumption of collision is elastic not inelastic.

Q) Under which of the following conditions is the law  $PV = RT$  obeyed most closely by a real gas

- (a) High pressure and high temperature
- (b) Low pressure and low temperature
- (c) Low pressure and high temperature
- (d) High pressure and low temperature

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. c

Solution:

At low pressure and high temperature real gas obey  $PV = RT$  i.e. they behave as ideal gas because at high temperature we can assume that there is no force of attraction or repulsion works among the molecules and the volume occupied by the molecules is negligible in comparison to the volume occupied by the gas.

Q) A certain sample of gas has a volume of 0.2 litre measured at 1 atm pressure and  $0^{\circ}\text{C}$ . At the same pressure but at  $273^{\circ}\text{C}$ , its volume will be

(a) 0.4 litres

(b) 0.8 litres

(c) 27.8 litres

(d) 55.6 litres

**Join Unacademy PLUS Referral Code :**

**Physicslive**



Ans. a

$$\text{at; } P = \text{const.}$$

$$P \cancel{V} = nRT$$

$$\frac{V}{T} = \text{const}$$


$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$

$$\frac{0.2}{273} = \frac{V_2}{(273+273)}$$

$$\frac{0.2}{273} = \frac{V_2}{2 \times 273}$$

$$\boxed{V_2 = 0.4 \text{ Ltr}}$$

Q) Figure shows two flasks connected to each other. The volume of the flask 1 is twice that of flask 2. The system is filled with an ideal gas at temperature 100 K and 200 K respectively. If the mass of the gas in 1 be  $m$  then what is the mass of the gas in flask 2



(a)  $m$

(b)  $m/2$


(c)  $m/4$

(d)  $m/8$

Join Unacademy PLUS Referral Code :

Physicslive

Ans. c



$$V_2 = 2V_1 \text{ (given)}$$

as both containers/flask are connected

$\Rightarrow P = \text{same in both}$

$$PV = nRT$$

$$P = \frac{nRT}{V}$$

$$P = \text{const}$$

$$\frac{n_1 RT_1}{V_1} = \frac{n_2 RT_2}{V_2}$$

$$\frac{n_1 R (100)}{(2V_2)} = \frac{n_2 R (200)}{V_2}$$

$$\boxed{n_2 = \frac{n_1}{2}}$$

multiply this equation by molecular mass of gas  $M$

$$n_2 \times M = \frac{n_1 M}{2}$$

$$\boxed{m_2 = \frac{m_1}{2}}$$

$m_1 = \text{mass of gas in flask 1}$   
 $m_1 = m$

$$\therefore \boxed{m_2 = \frac{m}{2}}$$

**Q)** If the pressure of an ideal gas contained in a closed vessel is increased by 0.5%, the increase in temperature is 2 K. The initial temperature of the gas is

(a)  $27^{\circ}\text{C}$

(b)  $127^{\circ}\text{C}$

(c)  $300^{\circ}\text{C}$

(d)  $400^{\circ}\text{C}$

**Join Unacademy PLUS Referral Code :**

**Physicslive**



Ans. b

Closed vessel  $\Rightarrow$  Volume is constant - &  $n = \text{constant}$

$$pV = nRT$$

$$\Rightarrow \frac{nRT}{p} = \text{constant}$$

$$\text{when } p_2 \rightarrow p_1 + \frac{0.5}{100} p_1$$

$$p_2 = p_1 + 0.005 p_1 = 1.005 p_1$$

$$\& \quad T_2 = T_1 + 2$$

$$\therefore \frac{nRT_1}{p_1} = \frac{nRT_2}{p_2} \Rightarrow \frac{T_1}{p_1} = \frac{T_1 + 2}{1.005 p_1}$$

$$1.005 T_1 = T_1 + 2$$

$$0.005 T_1 = 2$$

$$T_1 = \frac{2}{5 \times 10^{-3}} = \frac{2000}{5}$$

$$\boxed{T_1 = 400 \text{ K}}$$

$$\therefore T_1 = (400 - 273)^\circ \text{C}$$

$$\boxed{T_1 = 127^\circ \text{C}}$$

Q) Air is pumped into an automobile tube upto a pressure of  $200 \text{ kPa}$  in the morning when the air temperature is  $22^\circ\text{C}$ . During the day, temperature rises to  $42^\circ\text{C}$  and the tube volume expands by 2%. The pressure of the air in the tube at this temperature, will be approximately

(a)  $212 \text{ kPa}$

(b)  $206 \text{ kPa}$

(c)  $209 \text{ kPa}$

(d)  $200 \text{ kPa}$

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. c

$$PV = nRT$$

$$n = \text{const.}$$

$$\therefore \frac{PV}{T} = \text{const.}$$

$$P_1 = 200 \text{ kPa}$$

$$T_1 = 22 + 273 = 295 \text{ K}$$

$$V_1 = V$$

$$\text{Now } P_2 = ? ; T_2 = 42^\circ\text{C} = 42 + 273 \\ = 315 \text{ K}$$

$$V_2 = V + \frac{2}{10}V = 1.02V$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \Rightarrow \frac{(200 \text{ kPa}) \times V}{295} = \frac{P_2 (1.02V)}{315}$$

$$P_2 = 209 \text{ kPa}$$

Q) A vessel is filled with an ideal gas at a pressure of 10 atmospheres and temperature  $27^{\circ}\text{C}$ . Half of the mass of the gas is removed from the vessel and temperature of the remaining gas is increased to  $87^{\circ}\text{C}$ . Then the pressure of the gas in the vessel will be

(a) 5 atm

(b) 6 atm

(c) 7 atm

(d) 8 atm

**Join Unacademy PLUS Referral Code :**

**Physicslive**



Ans. b

$$P_1 = 10 \text{ atm.}$$

$$T_1 = 27^\circ\text{C} = 300 \text{ K}$$

$n_1 = \text{no. of moles}$

Now half gas is removed

$$\therefore n_2 = \frac{n_1}{2}$$

$$T_2 = 87^\circ\text{C} = 360 \text{ K.}$$

$$P_2 = ?$$

$$PV = nRT$$

$$\frac{P_1 V_1}{n_1 R T_1} = \frac{P_2 V_2}{n_2 R T_2}$$

$$\frac{10 \times V}{n_1 \times R \times 300} = \frac{P_2 \times V}{\left(\frac{n_1}{2}\right) R (360)}$$

$$2P_2 = \frac{10}{300} \times 360$$

$$P_2 = \frac{1800}{300}$$

$$\boxed{P_2 = 6 \text{ atm}}$$

Q) The pressure  $P$ , volume  $V$  and temperature  $T$  of a gas in the jar  $A$  and the other gas in the jar  $B$  at pressure  $2P$ , volume  $V/4$  and temperature  $2T$ , then the ratio of the number of molecules in the jar  $A$  and  $B$  will be-

(a) 1 : 1

(b) 2 : 1

(c) 1 : 2

(d) 4 : 1

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. d

$$n = \frac{PV}{RT}$$

$$\frac{n_1}{n_2} = \left( \frac{P_1 V_1}{RT_1} \right) / \left( \frac{P_2 V_2}{RT_2} \right) = \left( \frac{PV}{RT} \right) / \left( \frac{2P \times \frac{V}{2}}{R \times 2T} \right)$$





$$\frac{n_1}{n_2} = \frac{4}{1} \quad \left[ n_1 \text{ \& } n_2 \text{ are no. of moles of gas in jar A \& B} \right]$$

$$\frac{N_1/N_A}{N_2/N_A} = \frac{4}{1} \quad \left[ N_1 \text{ \& } N_2 \text{ are no. of molecules in jar A \& B} \right]$$

and  $N_A = \text{Avogadro number,}$

$$\boxed{\frac{N_1}{N_2} = \frac{4}{1}}$$

Q) Which one of the following graphs represents the behavior of an ideal gas when temperature is constant ?



Join Unacademy PLUS Referral Code :

Physicslive




Ans. d

$$PV = nRT$$

When;  $T = \text{constant}$

$$\Rightarrow PV = \text{constant}$$



Q) During an experiment an ideal gas is found to obey an additional law  $VP^2 = \text{constant}$ . The gas is initially at temperature  $T$  and volume  $V$ , when it expands to volume  $2V$ , the resulting temperature is -

(a)  $T/2$

(b)  $T/\sqrt{2}$

(c)  $T\sqrt{2}$

(d)  $2T$

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. c

$$VP^2 = \text{constant}$$

$$PV = nRT$$

$$P = \frac{nRT}{V}$$

$$\Rightarrow V \left( \frac{nRT}{V} \right)^2 = \text{constant}$$

$$(nR)^2 \frac{T^2}{V} = \text{const}$$

$$\Rightarrow \frac{T^2}{V} = \text{constant}$$

$$\frac{T_1^2}{V_1} = \frac{T_2^2}{V_2} \Rightarrow \frac{T^2}{V} = \frac{T_2^2}{2V}$$

$$T_2^2 = 2T^2$$

$$\boxed{T_2 = \sqrt{2} T}$$

Q) A vessel has 6 g of hydrogen at pressure  $P$  and temperature 500 K. A small hole is made in it so that hydrogen leaks out. How much hydrogen leaks out if the final pressure is  $P/2$  and temperature falls to 300 K -

(a) 2 g

(b) 4 g

(c) 3 g

(d) 1 g

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. d

$$\Rightarrow \because PV = nRT$$

$$\text{initially} \Rightarrow P(V) = 3 \times R \times 500 \quad - (1)$$

$$\text{finally} \Rightarrow \frac{P}{2}(V) = n_2 \times R \times 300 \quad - (2)$$

$$\frac{(1)}{(2)} \Rightarrow \frac{PV}{\frac{P}{2}V} = \frac{3R \times 500}{n_2 R \times 300}$$


$$2 = \frac{5}{n_2} \Rightarrow n_2 = 2.5 \text{ mole}$$

$$\begin{aligned} \text{mass of H}_2 \text{ gas in } 2.5 \text{ mole} &= 2.5 \times 2 \\ &= 5 \text{ gm.} \end{aligned}$$

$$\therefore \text{mass of leaked H}_2 \text{ gas} = 6 - 5 = \underline{\underline{1 \text{ gm}}}$$



Q) The figure shows pressure versus density graph for an ideal gas at two temperature  $T_1$  and  $T_2$ :




- (a)  $T_1 > T_2$                       (b)  $T_1 < T_2$   
(c)  $T_1 = T_2$                       (d) None of these

Join Unacademy PLUS Referral Code :

Physicslive

Ans. a



$$P = \frac{SRT}{M}$$

at constant  $S$

$$P \propto T \quad \text{or} \quad T \propto P$$

as from diagram!  $P_1 > P_2$

$$\Rightarrow \boxed{T_1 > T_2}$$

Q) A vessel is filled with a gas at a pressure of 76 cm of mercury at a certain temperature. The mass of the gas is increased by 50 % by introducing more gas in the vessel at the same temperature. The resultant pressure, in cm of Hg, is -

(a) 76

(b) 114


(c) 152

(d) 1117

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. b




Pr. = 76 cm of Hg

Temp. = T

mass of gas = m

M = molecular mass of gas.

$$\therefore n = \frac{m}{M}$$



Now  $m \rightarrow 1.5m$ .

$$\therefore n_2 = \frac{1.5m}{M} = 1.5n$$

Temp. = T

$P_1 = P_2 = ?$

[V = Volume = constant]

at  $T = \text{constant} \Rightarrow \frac{PV}{n} = \text{constant}$

$$\frac{(76 \text{ cm of Hg}) \times V}{n} = \frac{P_2 \times V}{1.5n}$$

$$P_2 = 114 \text{ cm of Hg}$$


Q) Pressure versus temperature graph of an ideal gas is as shown in figure. Density of the gas at point A is  $\rho_0$ . Density at B will be

(a)  $\frac{3}{4}\rho_0$

(b)  $\frac{3}{2}\rho_0$

(c)  $\frac{4}{3}\rho_0$

(d)  $2\rho_0$




Join Unacademy PLUS Referral Code :

Physicslive

Ans. b





$$P = \frac{\gamma R T}{M}$$

$$\Rightarrow \frac{P}{\gamma T} = \frac{R}{M} = \text{constant}$$

$$\left( \frac{P_0}{\gamma T} \right)_A = \left( \frac{P}{\gamma T} \right)_B \Rightarrow \frac{P_0}{\gamma_A T_0} = \frac{3P_0}{\gamma_B (2T_0)}$$

$$\frac{1}{\gamma_0} = \frac{3}{2\gamma_B}$$

$$\boxed{\gamma_B = \frac{3}{2} \gamma_0}$$

For Video Solution of this DPP, Click on below link

Video Solution  
on Website:-

<https://physicsaholics.com/home/courseDetails/57>

Video Solution  
on YouTube:-

<https://youtu.be/Ye7mcKHdp1I>

Chalo Niklo