



## SIR PRATEEK JAIN

- . Founder @Physicsaholics
- . Top Physics Faculty on Unacademy (IIT JEE & NEET)
- . 8+ years of teaching experience in top institutes like FIITJEE (Delhi, Indore) , CP (KOTA) etc.
- . Produced multiple Top ranks.
- . Research work with HC Verma sir at IIT Kanpur
- . Interviewed by International media.



## NEET UG subscription

PLUS

ICONIC \*\*

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✗ Personal Coach
- ✗ Study Planner

24 months ₹2,100/mo  
No cost EMI +10% OFF ₹50,400 >

18 months ₹2,363/mo  
No cost EMI +10% OFF ₹42,525 >

12 months ₹2,888/mo  
No cost EMI +10% OFF ₹34,650 >

6 months ₹4,200/mo  
No cost EMI +10% OFF ₹25,200 >

To be paid as a one-time payment

[View all plans](#)



Awesome! **PHYSICSLIVE** code applied



# PHYSICSLIVE

Use code **PHYSICSLIVE** to get 10% OFF on Unacademy PLUS and learn from India's Top Faculties.



## NEET UG subscription

PLUS

ICONIC \*\*

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✗ Personal Coach
- ✗ Study Planner

24 months ₹2,100/mo  
No cost EMI +10% OFF ₹50,400 >

18 months ₹2,363/mo  
No cost EMI +10% OFF ₹42,525 >

12 months ₹2,888/mo  
No cost EMI +10% OFF ₹34,650 >

6 months ₹4,200/mo  
No cost EMI +10% OFF ₹25,200 >

To be paid as a one-time payment

[View all plans](#)



Awesome! **PHYSICSLIVE** code applied



For Video Solution of this DPP, Click on below link

Video Solution  
on Website:-

<https://physicsaholics.com/home/courseDetails/58>

Video Solution  
on YouTube:-

<https://youtu.be/uUgK5PMvkDo>

# **JEE Main & Advanced, NSEP, INPhO, IPhO**

## **Physics DPP**

**DPP- 3 KTG: Kinetic Energy of Gas, Degree of freedom of gas molecules**

**By Physicsaholics Team**

Q) A gas mixture consists of 2 moles of oxygen and 4 moles of argon at temperature  $T$ . Neglecting all vibrational modes, the total internal energy of the system is:

- (a)  $4 R T$
- (b)  $5 R T$
- (c)  $15 R T$
- (d)  $11 R T$

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. d



Solution:

Internal energy of a gas  $U = \frac{f}{2} nRT$

$$U_{\text{oxygen}} = \frac{5}{2} \times 2 RT = 5RT$$

$$U_{\text{oxygen}} = \frac{3}{2} \times 4 RT = 6RT$$

$$U_{\text{total}} = 11RT$$

ANS (d)

Q) The molecules of an ideal gas have 6 degrees of freedom. The temperature of the gas is T. The average translational kinetic energy of its molecules is:

- (a)  $\frac{3}{2} k T$
- (b)  $\frac{6}{2} k T$
- (c)  $k T$
- (d)  $\frac{1}{2} k T$

**Join Unacademy PLUS Referral Code :**

**Physicslive**



Ans. a

Solution:

Av. energy associated with each  
degree of freedom =  $\frac{1}{2} kT$

degree of freedom of K.E. = 3

average translational K.E. =  $\frac{3}{2} kT$

ANS (a)

Q) The average translational kinetic energy of  $O_2$  (molar mass 32) molecules at a particular temperature is 0.048 eV. The translational kinetic energy of  $N_2$  (molar mass 28) molecules in eV at the same temperature is –

- (a) 0.0015
- (b) 0.003
- (c) 0.048
- (d) 0.768

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. c

Solution:

Average translational KE of a molecule does not depend on molar mass. At same temperature it is same for all gases.

ANS(c)

Q) A gas sample is enclosed in a closed container, temperature of gas is continuously increasing. Match the correct options in column-II corresponding to column-I

Column I	Column II
(a) Internal energy of gas	(P) Increases
(b) Average momentum of gas molecules	(q) Decreases
(c) Number of molecules moving with most probable speed	(r) Zero
(d) $\frac{V_{avg}}{V_{rms}}$	(s) Remains constant

Join Unacademy PLUS Referral Code :

Physicslive



Ans.  $a(p)$  ,  $b(r, s)$  ,  $c(q)$  ,  $d(s)$

Solution:


(a)  $U = \frac{f}{2} nRT \rightarrow U$  increases on increasing  $T$   
 $a \rightarrow p$

(b) Since  $\overline{V_{av}} = \vec{0} \Rightarrow \overline{P_{av}} = \vec{0}$   
 $b \rightarrow r \neq$

(c) maxima of graph shows  
no. of molecules having  
most probable velocity.

It decreases on increasing temperature

$c \rightarrow q$



$$(d) \quad \frac{V_{av}}{V_{rms}} = \frac{\sqrt{8RT/\pi M}}{\sqrt{3RT/M}} = \sqrt{\frac{8}{3\pi}}$$
$$= \text{Constant}$$

$d \rightarrow s$

Q) Temperature of an ideal gas is 300 K. The change in temperature of the gas when its volume changes from  $V$  to  $2V$  in the process  $P = aV$  (Here,  $a$  is a positive constant) is:

- (a) 900 K
- (b) 1200 K
- (c) 600 K
- (d) 300 K

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. a

Solution:

$$T = \frac{PV}{nR} = \frac{av^2}{nR} \propto v^2$$

$v \rightarrow 2 \text{ times}$

$\Rightarrow T \rightarrow 4 \text{ times}$

$$\begin{aligned} \text{final temperature} &= 300 \times 4 \\ &= 1200 \text{ K} \end{aligned}$$

$$\Delta \text{change in temperature} = 900 \text{ K}$$

ANS (a)




Q) In the  $p$ - $T$  graph shown in figure, match the following:

Table-1

- (a) Process a-b
- (b) Process b-c
- (c) Process c-d
- (d) Process d-a

Table-2

- (p) Constant volume
- (q)  $\Delta U = 0$
- (r)  $P$  increasing
- (s)  $P$  decreasing



Join Unacademy PLUS Referral Code :

Physicslive

Ans.  $a(q, r)$  ,  $b(p, r)$  ,  $c(q, s)$  ,  $d(p, s)$

Solution:

(a)  $T = \text{Constant} \Rightarrow \Delta U = 0$  ;  $P$  is increasing

$$P = \frac{PM}{RT} \Rightarrow P = \frac{PRT}{M} \Rightarrow P \text{ is increasing}$$

(b)  $P = \text{Constant}$ ,  $T$  is increasing  
 $a \rightarrow v, \gamma$

$$P = \frac{PRT}{M} \Rightarrow P \text{ is increasing}$$

$$P = \text{Constant} \Rightarrow \frac{P}{T} = \text{Constant} \Rightarrow V = \text{Constant}$$

$b \rightarrow P, \gamma$

(c)  $P$  is decreasing,  $T$  is constant

$\Downarrow$

$P$  is decreasing

$\Downarrow$

$$\Delta U = 0$$

$C \rightarrow Q, S$

(d)  $P$  is constant,  $T$  is decreasing

$\Downarrow$

$V$  is constant

$\Downarrow$

$$\Delta U \neq 0 \text{ \& } P = \frac{PRT}{m}$$

$\Downarrow$

$P$  is decreasing

$d \rightarrow P, S$

Q) One mole of an ideal gas undergoes a process  $P = \frac{P_0}{1 + \left(\frac{V_0}{V}\right)^2}$ . Here,  $P_0$  and  $V_0$  are constants. Change in temperature of the gas when volume is changed from  $V = V_0$  to  $V = 2V_0$  is :

(a)  $-\frac{2P_0V_0}{5R}$

(b)  $\frac{11P_0V_0}{10R}$

(c)  $-\frac{5P_0V_0}{4R}$

(d)  $P_0V_0$

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. b



Solution:

$$P = \frac{P_0}{1 + \left(\frac{V_0}{V}\right)^2}$$

$$\text{at } V = V_0, \quad P = \frac{P_0}{1+1} = \frac{P_0}{2} \Rightarrow T_1 = \frac{P_0 V_0}{2R}$$

$$\text{at } V = 2V_0, \quad P = \frac{P_0}{1+\frac{1}{4}} = \frac{4P_0}{5} \Rightarrow T_2 = \frac{4P_0 \times 2V_0}{5R}$$

change in temperature

$$\Delta T = \frac{P_0 V_0}{R} \left[ \frac{8}{5} - \frac{1}{2} \right] = \frac{11 P_0 V_0}{10 R}$$

ANS (b)

Q) Two containers of equal volume contain the same gas at pressures  $p_1$  and  $p_2$  and absolute temperatures  $T_1$  and  $T_2$  respectively. On joining the vessels, the gas reaches a common pressure  $p$  and a common temperature  $T$ . The ratio  $P/T$  is equal to

(a)  $\frac{p_1}{T_1} + \frac{p_2}{T_2}$

(b)  $\frac{1}{2} \left[ \frac{p_1}{T_1} + \frac{p_2}{T_2} \right]$

(c)  $\frac{p_1 T_2 + p_2 T_1}{T_1 + T_2}$

(d)  $\frac{p_1 T_2 - p_2 T_1}{T_1 - T_2}$

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. b

PRATEEK JAIN  
PHYSICSAHOLICS

Solution:

$V \rightarrow$  Volume of one Container

$n_1$  &  $n_2$  are no of moles in Containers.


$$n_1 = \frac{P_1 V}{RT_1}, \quad n_2 = \frac{P_2 V}{RT_2}$$

$$\begin{aligned} \frac{P}{T} &= \frac{nR}{2V} = \frac{(n_1 + n_2)R}{2V} = \frac{n_1 R}{2V} + \frac{n_2 R}{2V} \\ &= \frac{P_1}{2T_1} + \frac{P_2}{2T_2} \end{aligned}$$

ANS (b)

## COMPREHENSION

Figure shows a cylindrical tube of volume  $V$  with adiabatic walls containing an ideal gas. The internal energy of this gas is given by  $1.5 nRT$ . The tube is divided into two equal parts by a fixed diathermic wall.



Initially the pressure and temperature on the two sides are  $p_1, T_1$  and  $p_2, T_2$  respectively. The system is left for sufficient time so that the temperature becomes equal on the two sides.

Q) What is the ratio of pressures on the left and right sides?

(a)  $p_2 T_2 / p_1 T_1$

(b)  $p_1 T_2 / p_2 T_1$

(c)  $\frac{p_1 + p_2}{T_1 + T_2}$

(d)  $\frac{p_1 T_1}{p_2 T_2}$

Join Unacademy PLUS Referral Code :

Physicslive



Ans. b

Solution:

$P_1$ $T_1$ $n_1$	$P_2$ $T_2$ $n_2$
-------------------------	-------------------------

$V/2$        $V/2$



$P_1 f$ $T$	$P_2 f$ $T$
----------------	----------------

$$n_1 = \frac{P_1 V}{2RT_1}$$

$$n_2 = \frac{P_2 V}{2RT_2}$$

$$\frac{P_1 f}{P_2 f} = \frac{\frac{n_1 RT}{V/2}}{\frac{n_2 RT}{V/2}} = \frac{n_1}{n_2} = \frac{P_1 T_2}{P_2 T_1}$$

Ans (b)

Q) What is the final equilibrium temperature?

(a)  $\frac{T_1 T_2 (p_1 + p_2)}{p_1 T_2 + p_2 T_1}$

(b)  $\frac{p_1 p_2 (T_1 + T_2)}{p_1 T_2 + p_2 T_1}$

(c)  $\frac{T_1 T_2 (p_1 + p_2)}{p_1 T_1 + p_2 T_2}$

(d)  $\frac{T_1^2 p_2^2}{p_1 T_2 + p_2 T_1}$

**Join Unacademy PLUS Referral Code :**

**Physicslive**

Ans. a

Solution:

Internal energy loss by left + Internal energy gain by right = 0

$$\Rightarrow \frac{f}{2} n_1 R (T - T_1) + \frac{f}{2} n_2 R (T - T_2) = 0$$

$$\Rightarrow \frac{n_1}{n_2} (T - T_1) + (T - T_2) = 0$$

$$\Rightarrow \frac{P_1 T_2}{P_2 T_1} (T - T_1) + (T - T_2) = 0$$

$$\Rightarrow T (P_1 T_2 + P_2 T_1) = \frac{P_1 T_1 T_2 + P_2 T_1 T_2}{(P_1 + P_2) T_1 T_2}$$

$$\Rightarrow T = \frac{P_1 T_2 + P_2 T_1}{P_1 + P_2}$$

Ans (a)

For Video Solution of this DPP, Click on below link

Video Solution  
on Website:-

<https://physicsaholics.com/home/courseDetails/58>

Video Solution  
on YouTube:-

<https://youtu.be/uUgK5PMvkDo>

Chalo Niklo