



SIR PRATEEK JAIN

- . IIT JEE & NEET FACULTY (KOTA)
- . TOP PHYSICS FACULTY ON UNACADEMY.
- . 8+ YEARS OF TEACHING EXPERIENCE
- . RESEARCH WORK WITH HC VERMA SIR AT IIT KANPUR
- . PRODUCED RANKS LIKE AIR 6, AIR 10 ETC.

PLUS

ICONIC ^{**}

✓ India's Best Educators

✓ Interactive Live Classes

✓ Structured Courses & PDFs

✓ Live Tests & Quizzes

✗ Personal Coach

✗ Study Planner

24 months

No cost EMI

₹2,333/mo

₹56,000

>

18 months

No cost EMI

₹2,625/mo

₹47,250

>

12 months

No cost EMI

₹3,208/mo

₹38,500

>

6 months

No cost EMI

₹4,667/mo

₹28,000

>

To be paid as a one-time payment

View all plans

 Add a referral code

APPLY

PHYSICSLIVE

PLUS

ICONIC ^{**}

✓ India's Best Educators

✓ Interactive Live Classes

✓ Structured Courses & PDFs

✓ Live Tests & Quizzes

✗ Personal Coach

✗ Study Planner

24 months

No cost EMI

₹2,100/mo

+10% OFF ₹50,400

>

18 months

No cost EMI

₹2,363/mo

+10% OFF ₹42,525

>

12 months

No cost EMI

₹2,888/mo

+10% OFF ₹34,650

>

6 months

No cost EMI


₹4,200/mo

+10% OFF ₹25,200

>

To be paid as a one-time payment

View all plans

 Awesome! PHYSICSLIVE code applied

×

Use code **PHYSICSLIVE** to get 10% OFF on Unacademy PLUS.

For Video Solution of this DPP, Click on below link

<https://physicsaholics.com/home/courseDetails/49>

JEE Main **Physics DPP**

DPP-1 Units & Measurements: Units & Dimensional Formula
By Physicsaholics Team

Q) Which of the following physical quantities has neither dimensions nor unit?
(Hint:- $f = \mu N$; where, μ = coefficient of friction, f = friction force & N = Normal force)

(a) Angle

(b) Luminous intensity

(c) Coefficient of friction

(d) Current

Join Unacademy PLUS Referral Code :

Physicslive

Ans. c

Luminous intensity

↳ cd

Current

↳ Amp.

Angle

↳ rad

Coefficient of friction

$$\mu = \frac{f}{N} = \frac{\text{Force}}{\text{Force}}$$

↳ No unit

No dimension.

Q) Dimensional formula for coefficient of viscosity (η) [use $F = 6\pi\eta rv$ (r=radius ; v=velocity; F=viscous force)]:

(a) $ML^{-2}T^{-1}$

(c) $M^1L^1T^{-2}$

(b) $M^{-1}L^1T^{-1}$

(d) $ML^{-1}T^{-1}$

Join Unacademy PLUS Referral Code :

Physicslive

Ans. d

$$F = G \lambda x y v$$

$$[F] = M L T^{-2}$$

$$[x] = L$$

$$[v] = L T^{-1}$$

$$[\lambda] = \frac{[F]}{[x][v]} = \frac{M L T^{-2}}{L \cdot L T^{-1}}$$

$$= \underline{M L^{-1} T^{-1}}$$

Q) The dimensions of radian per second are:

(a) $[M^0 L^0 T^0]$

(c) $[M^0 L^0 T^{-1}]$

(b) $[M^0 L^0 T^1]$

(d) $[M^0 L^2 T^{-1}]$

Join Unacademy PLUS Referral Code :

Physicslive

Ans. c

radian per sec = rad/sec

rad \rightarrow Angle \rightarrow dimensionless

$$[\text{Radian per sec}] = \frac{1}{[\text{sec}]} = \frac{1}{T} = \underline{T^{-1}}$$

Q) The dimensional formula of radius of gyration is:

(a) $[M^0 L^0 T^0]$

(b) $[M^0 L^0 T]$

(c) $[M^0 L T^0]$

(d) $[M L T^{-1}]$

Join Unacademy PLUS Referral Code :

Physicslive

Ans. c

Solution:

Radius of gyration is measure of distance.

(You will study this in the chapter “Rotational Motion.”)

Q) From the following pairs of physical quantities, in which group dimensions are not same:

[Hint:- Linear momentum = mass \times velocity, Torque = Force \times perpendicular distance, Impulse = Change in momentum]

- (a) Linear Momentum and impulse (b) Torque and energy
(c) Energy and work (d) Light year and minute

Join Unacademy PLUS Referral Code :

Physicslive

Ans. d

(a) Linear Momentum (P)

Impulse (I)

$$P = mv$$

$$I = \Delta P$$

$$\text{So, } [I] = [P]$$

(b) Torque & Energy

(τ)

(E)

$$\tau = F \times r_{\perp}$$

$$E = F \times x$$

Energy or work

$$[\tau] = [E]$$

(c) Energy & work

$$[E] = [W]$$

(d) Light year is length
minute is time

$$[L] \neq [T]$$

Q) The dimensional formula for Planck's constant (h) is
(Hint:- Unit of planks constant = J-sec)

(a) $[ML^{-2}T^{-3}]$

(c) $[ML^2T^{-1}]$

(b) $[M^0L^2T^{-2}]$

(d) $[ML^{-2}T^{-2}]$

Join Unacademy PLUS Referral Code :

Physicslive

Ans. c

$$[h] = ?$$

Unit of $h = \text{Joule} \cdot \text{sec}$

Joule \rightarrow Energy

$$[E] = M L^2 T^{-2}$$

$$[h] = [E] [t]$$

$$= M L^2 T^{-2} \cdot T$$

$$= M L^2 T^{-1}$$

Q) An atmosphere:

- (a) is a unit of pressure
- (b) is a unit of force
- (c) gives an idea of the composition of air
- (d) is the height above which there is no atmosphere

Join Unacademy PLUS Referral Code :

Physicslive

Ans. a

atmosphere (atm) is the unit
of pressure.

$$1 \text{ atm} = 10^5 \text{ pascal}$$

Q) The dimensions of wavelength (λ) is:
(Wavelength = Distance travelled by wave in one time period)

(a) $[M^0 L^0 T^0]$

(b) $[M^0 L T^0]$

(c) $[M^0 L^{-1} T^0]$

(d) none of these

Join Unacademy PLUS Referral Code :

Physicslive

Ans. b

wavelength = measure of
length

$$[\lambda] = L \text{ or } M^0 L^1 T^0$$

Q) State which of the following is correct?

(Hint:- When a charge q is accelerated by a Voltage V then its energy = qV)

(a) joule = coulomb \times volt

(b) joule = coulomb/volt

(c) joule = volt + coulomb

(d) joule = volt/coulomb

Join Unacademy PLUS Referral Code :

Physicslive

Ans. a

$$\text{Energy} = q \Delta V$$

$$\text{Joule} = \text{Coulomb} \times \text{Volt}$$

Q) Of the following quantities, which one has dimensions different from the remaining three?

(Hint:- Angular Momentum = mass \times velocity \times perpendicular distance,
& When a charge q is accelerated by a voltage V then its energy = qV)

- (a) Energy per unit volume
- (b) Force per unit area
- (c) Product of voltage and charge per unit volume
- (d) Angular momentum

Join Unacademy PLUS Referral Code :

Physicslive

Ans. d

$$(a) \left[\frac{E}{V} \right] = \frac{ML^2T^{-2}}{L^3} = ML^{-1}T^{-2}$$

$$(b) \left[\frac{F}{A} \right] = \frac{MLT^{-2}}{L^2} = ML^{-1}T^{-2}$$

$$(c) \left[\frac{\text{Volt} \times \text{Coulomb}}{V} \right] = \frac{\text{Joule}}{\text{Volume}} = \frac{ML^2T^{-2}}{L^3} = ML^{-1}T^{-2}$$

= Energy/Volume

$$(d) [L] = [mvr] = MLT^{-1}L = ML^2T^{-1}$$

(angular momentum is denoted by = L)

Q) The dimensions of frequency is:

(Hint:- frequency $(f) = \frac{1}{T}$; T = Time period)

(a) $[T^{-1}]$

(c) $[M^0 L^0 T^{-2}]$

(b) $[M^0 L^0 T^0]$

(d) None of these

Join Unacademy PLUS Referral Code :

Physicslive

Ans. a

frequency, $f = \frac{1}{T}$

T = Time period

$$[T] = T$$

$$[f] = \frac{1}{T} = T^{-1}$$

$$[f] = T^{-1} \text{ or } M^0 L^0 T^{-1}$$

Q) Young's modulus (Y) of a material has the same unit as
($Y = \frac{\text{Stress}}{\text{Strain}}$; where, $\text{Stress} = \frac{\text{Force}}{\text{Area}}$ & $\text{Strain} = \frac{\text{Change in length}}{\text{original length}}$)

(a) Pressure

(b) Strain

(c) Density

(d) Force

Join Unacademy PLUS Referral Code :

Physicslive

Ans. a

$$Y = \frac{\frac{F}{A}}{\frac{\Delta l}{l}} = \text{Youngs Modulus}$$

$$[Y] = \frac{\left[\frac{F}{A}\right]}{\left[\frac{\Delta l}{l}\right]} \quad \left[\because \frac{\Delta l}{l} \rightarrow \text{Dimensionless} \right]$$

$$\therefore [Y] = \left[\frac{F}{A}\right]$$

$$(1) \text{ Pressure } = P = \frac{F}{A}$$

$$[P] = \left[\frac{F}{A}\right]$$

$$\therefore [Y] = [P]$$

Q) The unit of impulse is the same as that of
(Hint:- Impulse = Force \times time, Momentum = mass \times velocity, Power = Energy per unit time)

(a) Energy

(b) Power

(c) Momentum

(d) Velocity

Join Unacademy PLUS Referral Code :

Physicslive

Ans. c

$$\begin{aligned}\text{Impulse} &= \text{Force} \times \text{time} \\ &= \text{MLT}^{-2} \times \text{T} = \text{MLT}^{-1}\end{aligned}$$

$$\begin{aligned}\text{Momentum} &= \text{mass} \times \text{velocity} \\ &= \text{M} \times \text{LT}^{-1} = \text{MLT}^{-1}\end{aligned}$$

For Video Solution of this DPP, Click on below link

<https://physicsaholics.com/home/courseDetails/49>

Chalo Niklo